




Institute of  
Medicinal Plants

## Journal of Medicinal Plants

Journal homepage: [www.jmp.ir](http://www.jmp.ir)



### Research Article

## Phytochemical, physicochemical and biological evaluation of *Colchicum kurdicum* (Bornm.) Stef.: a study on materia medica of Persian medicine

Mohammad Azadbakht<sup>1</sup>, Ali Davoodi<sup>1,\*</sup>, Seyed Jalal Hosseinimehr<sup>2</sup>, Saeed Emami<sup>3</sup>, Masoud Azadbakht<sup>4</sup>, Fatemeh Mirzaee<sup>1</sup>, Hossein Bakhshi Jouybari<sup>1</sup>

<sup>1</sup> Department of Pharmacognosy, Faculty of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

<sup>2</sup> Department of Radiopharmacy, Faculty of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

<sup>3</sup> Department of Medicinal Chemistry and Pharmaceutical Sciences Research Center, Faculty of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

<sup>4</sup> Department of Plant Systematics, High Educational of Sanna Institute, Sari, Iran

### ARTICLE INFO

#### Keywords:

*Colchicum kurdicum*  
Tropolone alkaloid  
Total flavonoid  
Chromatography  
Antioxidant  
Anti-inflammatory activity

### ABSTRACT

**Background:** The genus *Colchicum* (Colchicaceae) is a perennial and monocotyledon flowering plant that has more than 160 species in the world. In addition, this plant is an important medicinal plant in traditional and modern medicines. **Objective:** The aim of this study was to evaluate the phytochemical profiles and physicochemical properties, antioxidant and anti-inflammatory activities of *Colchicum kurdicum* (Bornm.) Stef. corm. **Methods:** Phytochemical profiles included total tropolone alkaloid, total phenolic/total tannin and total flavonoid contents were determined by spectrophotometric method. Moreover, tropolone alkaloid profiles was analyzed by HPLC method. Physicochemical properties including macroscopic and organoleptic properties, solubility, foreign matter, ash values and heavy metals were evaluated based on pharmacopeial protocols. In addition, in vitro anti-inflammatory and antioxidant activities of the plant were determined. **Results:** Total tropolone alkaloids, phenol, tannin and flavonoid contents of the corm were estimated to be 0.652, 0.426, 0.052 and 0.325 g/100 g corm. In HPLC analysis, N-deacetyl-N-formyl colchicine, colchifoline, colchicoside and cornigerine were calculated as highest bioactive tropolone alkaloids. Physicochemical properties were determined within an acceptable range for the pharmacopoeia. Moreover, the anti-inflammatory and antioxidant activities in 10 mg/ml of methanolic extract were estimated 84.7 % and 94.8 %, respectively. **Conclusion:** The corm of the *Colchicum kurdicum* had the main bioactive compounds. In addition, these compounds have the valuable biological activities as anti-inflammatory and antioxidant activities.

**Abbreviations:** HPLC, High Performance Liquid Chromatography; PVPP, Polyvinyl Polypyrrolidone; ACN, Acetonitrile; SD, Standard Deviation

\*Corresponding author: [ali.davoodi@mazums.ac.ir](mailto:ali.davoodi@mazums.ac.ir)

doi: 10.29252/jmp.19.76.36

Received 13 October 2019; Received in revised form 24 June 2020; Accepted 19 July 2020

© 2020. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>)

## 1. Introduction

The genus *Colchicum* (Colchicaceae) is a perennial and monocotyledon flowering plant with more than 160 species in the world. In addition, the bulb-like corms and underground ovaries are the individual morphological characterizations of this genus [1, 2].

*Colchicum kurdicum* (Bornm.) Stef. is the important medicinal plant of this family, which is named as "Sorenjân" and was used in many diseases such as osteoarthritis, gout, cancer, inflammations, jaundice and sexual impotencies in the different traditional medicines.

In addition, the main pharmaceutical organs of *C. kurdicum* are the seeds and corms, which are contained the important bioactive compounds, including alkaloids especially tropolone alkaloids and isoquinoline alkaloids, phenolic compounds, tannins, flavonoids and carbohydrates [1, 2].

Tropolone alkaloids especially colchicine as the main important bioactive compounds of this plant are used for treatment of some diseases such as gout and cancers. These alkaloids could be used in mitosis-dependent diseases by the tubulin polymerization inhibitory effects. In addition, these compounds have high potential toxicity and a narrow therapeutic index [3].

Furthermore, other bioactive compounds of *C. kurdicum* are included isoquinoline alkaloids with anticholinesterase inhibitory activities, coumaric acid, ferulic acid, caffeic acid, vanillic acid, 2-hydroxybenzoic acid, apigenin, luteolin and 4, 6-dimethoxy-3, 7-dimethylcoumarin [4].

According to the literatures, the bioactive compounds of the *C. kurdicum*, especially tropolone alkaloids, flavonoids and phenolic acids, have anti-inflammatory, anti-arthritic and anti-oxidant activity.

Phytochemical, physicochemical and biological evaluations of the medicinal plants by

different spectrophotometric, *in vitro* and *in vivo* methods are the critical stage for validation of the quality. Moreover, the findings of the phytochemical profiles and bioactive compounds could be helped to estimate the biological activities [5, 6].

In the present study, phytochemical profiles of *C. kurdicum* were evaluated by spectrophotometric and HPLC methods. Furthermore, physicochemical properties including macroscopic and organoleptic properties, solubility, foreign matter, ash values and heavy metals were determined. In addition, biological activities included anti-inflammatory and anti-oxidant activities of *C. kurdicum* were evaluated.

## 2. Materials and Methods

### 2.1. Materials

All materials were purchased from the Sigma and Merck companies. All acids and solvents were purchased from Merck Company (Germany).

### 2.2. Plant preparation

The plant specimen was collected from 1500 to 2500 regions of Tang-e-Rah area of the Golestan National Park (37.366335, 55.780033) in Golestan province in October to April 2018. Dr. Masoud Azadbakht performed taxonomic identification of plant samples, and representative voucher specimens were deposited in the herbarium of the Department of Pharmacognosy in Faculty of Pharmacy of the Mazandaran University of Medical Science (E<sub>1</sub>-11312) [7].

### 2.3. Preparation of corm extracts

The powdered corms of the *C. kurdicum* were extracted using methanol with percolation method (24 h maceration and 1:10 plant/solvent

ratio). Then, the methanolic extract was concentrated using rota-vapour and was dried using freeze-dryer. Finally, the extract was stored at 4 °C until analysis [8].

#### 2.4. Total assay of phytoconstituents

Total tropolone alkaloids of the corm methanolic extract was determined using the acidic potassium dichromate method. One g of corm extract was dispersed in the mixture of methanol/H<sub>2</sub>SO<sub>4</sub> solution (3 %, pH = 1) and was extracted using sonication during 1 hour. The final acidic extract was decanted with chloroform and aqueous phase was collected and was alkalinized using NH<sub>4</sub>OH solution (10 %, pH = 12). The solution was extracted three times with chloroform and was dried. Then, 1 ml solution of extract (1 mg/ml) and (2.5, 5, 10, 20 and 40 mg/ml) colchicine were mixed with 1 ml acidic potassium dichromate solution (pH = 2) and were incubated at room temperature for 1 hour. Finally, UV absorbance was measured at 352 nm and the graph of calibration was drawn [9, 10].

Total phenolic and tannin contents of the corm extract were determined using the Folin-Ciocalteu method. One ml of the extract (1 mg/ml) was mixed with 1.5 ml Folin-Ciocalteu and was incubated at room temperature for 10 minutes. Then, 1.5 ml sodium carbonate solution (20 %) was added to the mixture and was incubated at room temperature for an additional 1 hour and absorptions at 725 nm were recorded. In addition, calibration curve was created using a standard concentration of gallic acid and total phenolic compounds of extract were obtained by calibration curve. Moreover, 100 mg polyvinyl polypyrrolidone (PVPP), 1 ml distilled water and 1 ml of the extracts (1 mg/ml) were added into tube and was mixed using and kept in 4 °C for 15 minutes. Then, it

was centrifuged (3000 g for 15 min) and collected the supernatant. This supernatant has only simple phenolic compounds other than tannins. The phenolic content of the supernatant was measured as mentioned above. Finally, the total tannin was obtained by differences of two above amount [6].

Furthermore, the total flavonoid content the corm extract was determined using the AlCl<sub>3</sub> method. One ml of the extracts (1 mg/ml) was mixed with 1 ml of AlCl<sub>3</sub> solution (2 %) and 1 ml potassium acetate solution (10 %) and was incubated at room temperature for 30 minutes. Finally, the absorbance of the mixture was measured at 415 nm and the total flavonoid contents was determined using a standard curve with quercetin (2.5, 5, 10, 20 and 40 mg/ml) as the standard [5].

#### 2.5. HPLC analysis

The HPLC analysis of the corm extract of the *C. kurdicum* was done by HPLC-UV. The separation, detection and assay of tropolone alkaloids including colchicine, demecolcine, 2-demethyl colchicine, 3-demethyl colchicine, colchicoside, colchifoline, cornigerine and N-deacetyl-N-formyl colchicinewere performed by a HPLC Smartline Manager 5000 (Knauer, Germany) with Smartline pump 1000 and EC Nucleodur C<sub>18</sub> column (4.6 mm × 250 mm, 5 µm particle size) at 25°C with UV detector 2500 basic model in 245 nm. Moreover, data acquisition and integration were performed with EZchrom Elite 3.2.0 software. The confidence of the accuracy of the individuality of the peaks was obtained by standard addition. The extract obtained from the mentioned above method (1 mg/ml) and the HPLC grade solutions of the compounds (125, 250, 500, 1000 and 2000 ppm) were injected into the HPLC with 20 µl volume and flow rate of the mobile phase was maintained at 1 ml/min. The mobile phases

were HPLC grade acetonitrile (ACN) and water. Moreover, the gradient condition were: 0-5 min, ACN 10 %; 5-20 min, ACN 100 %; 20-25 min, ACN 10 %; 25-30 min, ACN 10 %. Finally, the calibration curves of all compounds were obtained by plotting the peak region against the concentration of each sample and the amount of the compounds was calculated by the obtained formulas. Moreover,  $R^2 > 0.9$  representing the linear measurements [2, 11].

### 2.6. Physicochemical properties of the corm

All physicochemical properties of the *C. kurdicum*, including macroscopic and organoleptic properties, solubility, foreign matter, ash values and heavy metals were separately done based on pharmacopeial protocols [12, 13].

### 2.7. In vitro anti-inflammatory activity

In this study, anti-denaturation activities (as a parameter for determination of anti-inflammatory activity) of the *C. kurdicum* extract were evaluated. Five ml of the extract (10 mg/ml) was mixed with 1 ml of 2 % bovine serum albumin and 1 ml of distilled water and was incubated for 1 hour at 37°C. Then, the mixture was heated at 60 °C for 10 min and was cooled. In addition, 3 ml of phosphate buffer saline (pH = 7.4) was added to the final mixture and was evaluated by UV spectrophotometric method in 600 nm. The anti-denaturation activities were calculated using the following formula:

$$\text{Percentage of anti-denaturation activities} = 100 \times (\text{Absorbance of control} - \text{Absorbance of treated sample}) / \text{Absorbance of control}$$

The control solution was contained the distilled water instead of extract. Moreover, the diclofenac sodium was used as standard drug [14].

### 2.8. Antioxidant activity

The radical scavenging activity of methanolic extract of *C. kurdicum* was evaluated by stable DPPH free radical. A solution of DPPH (250  $\mu$ M) was prepared by dissolving DPPH (5 ml) in the methanol (2 ml) and the solution was kept in the dark at room temperature. Furthermore, different concentrations of the extract were prepared and mixed with the same volume of DPPH solution and incubated and analysed at 517 nm. The percentage of inhibition of the extract was determined by comparison with the methanol control group. In addition, ascorbic acid was used as positive control [15].

### 2.9. Statistical analysis

All analysis procedures were done triplet and the data were analysed by SPSS and Excel softwares and were characterized with mean  $\pm$  SD.

## 3. Results

### 3.1. Phytochemical characteristics


Table 1 shows the results of the total assay of the main bioactive phytoconstituents in the corm extract of the *C. kurdicum*. The amounts of total alkaloid, total phenol, total tannin and total flavonoid contents were estimated 0.05 to 0.7%.

### 3.2. HPLC analysis

Fig. 1 and Table 2 show the chromatograms and HPLC results of the *C. kurdicum*. The amounts of the compounds were calculated by HPLC. According to the obtained chromatograms, N-deacetyl-N-formyl colchicine, colchifoline, colchicoside and cornigerine were the main compounds in *C. kurdicum*.

**Table 1.** Total tropolone alkaloid, phenol, tannin and flavonoid contents of the *C. kurdicum*

Compounds	Total tropolone alkaloid (g colchicine/100 g corm)	Total phenol contents (g gallic acid/100 g corm)	Total tannin contents (g gallic acid/100 g corm)	Total flavonoid contents (g quercetin/100 g corm)
<i>C. kurdicum</i>	0.692 ± 0.021	0.426 ± 0.034	0.052 ± 0.005	0.325 ± 0.028

**Fig. 1.** The HPLC curves of the tropolone alkaloid-rich extract of the *C. kurdicum* and the standard curves of the compounds (1: Colchicoside, 2: 2-demethyl colchicine, 3: 3-demethyl colchicine, 4: Demecolcine, 5: Colchifoline, 6: N-Deacetyl-N-formyl Colchicine, 7: Colchicine, 8: Cornigerine)**Table 2.** The amounts of the tropolone alkaloids in the *C. kurdicum* based on HPLC data (mg/100 g corm)

Compounds	Colchicoside	2-demethyl colchicine	3-demethyl colchicine	Demecolcine	Colchifoline	N-deacetyl-N-formyl colchicine	Colchicine	Cornigerine
<i>C. kurdicum</i>	135.56 ± 1.34	48.12 ± 0.98	22.04 ± 0.92	18.76 ± 0.91	136.93 ± 0.82	177.65 ± 0.81	66.34 ± 0.23	129.95 ± 0.92

### 3.3. Physicochemical characteristics

Table 4 shows the physicochemical properties of the corm of the *C. kurdicum*. According to obtained data, the shape of the corms has been varied from oblong to oval. The colour of the powdered corms was grey and odour and taste of them were spicy and bitter, respectively. In addition, solubility of dry powder of *C. kurdicum* corms was estimated sparingly soluble in different solvents. Other physicochemical properties of the corm were illustrated in table 3.

### 3.4. Anti-inflammatory and anti-oxidant activities

The results of the anti-denaturation and radical scavenging activities of the *C. kurdicum* are shown in the table 4. According to this study, the anti-denaturation and radical scavenging activities of the methanolic extract of *C. kurdicum* were estimated  $84.7 \pm 1.2\%$  and  $94.8 \pm 1.8\%$ .

**Table 3.** Physicochemical data of the *C. kurdicum*

Characteristics	<i>C. kurdicum</i>	
Organoleptic Properties	Shape	Oblong to Oval
	Powder Colour	White to Gray
	Odour	Mild Spicy
	Taste	Bitter
Solubility	Water	Sparingly Soluble
	KOH 10%	Sparingly Soluble
	H <sub>2</sub> SO <sub>4</sub> 5%	Freely Soluble
	Methanol	Sparingly Soluble
	Ethanol 96%	Soluble
	Chloroform	Freely Soluble
	Hexane	Freely Soluble
Foreign Matter (%)		$1.8 \pm 0.16$
Moisture Content (%)		$5.4 \pm 1.53$
Ash Content (%)	Total Ash	$7.8 \pm 1.54$
	Acid Insoluble Ash	$4.3 \pm 1.23$
	Sulphated Ash	$25.5 \pm 0.89$
	Water Insoluble Ash	$0.96 \pm 0.05$
	Water Soluble Ash	$3.2 \pm 0.23$
Heavy Metals ( $\mu\text{g}/\text{kg}$ )	Hg	$0.5 \pm 0.01$
	Cu	$0.3 \pm 0.02$
	Cd	$0.1 \pm 0.07$
	Pb	$0.1 \pm 0.06$

**Table 4.** The anti-inflammatory and antioxidant activities of the *C. kurdicum* methanolic extract.

Compounds	<i>C. kurdicum</i> (10 mg/ml)	Diclofenac Sodium (10 mg/ml)	Ascorbic Acid (10 mg/ml)
Anti-denaturation activity (%)	$84.7 \pm 1.2$	$94.4 \pm 1.5$	-
Radical scavenging activity (%)	$94.8 \pm 1.8$	-	$81.8 \pm 1.4$

### 3. Discussion

In this study, phytochemical and physicochemical properties and anti-inflammatory and anti-oxidant activities of the corm of *C. kurdicum* was evaluated. The preliminary phytochemical studies could be helped to estimate the constituents of the plants and potential and valuable biological activities. In the different studies, the bioactive compounds of the plants have been evaluated. Researchers determined the phytochemical profiles of *C. luteum* Baker. In addition, this study reported the alkaloids, phenolic compounds, flavonoids, sterols, tannins and saponins in the methanolic extracts of the plant corms [4]. Moreover, other researchers detected the similar compounds in the corm extract of *Gloriosa superba* Linn [16].

In the different studies phytoconstituents of the plants and other bioresources were assayed. In a study, the chemical constituents of different parts of the *C. baytopiorum* C.D.Brickell that total alkaloids of the plant were determined by the spectrophotometric method [17]. In this study, total alkaloids were estimated 5.27%, 2.96% and 1.96% in the perigon, perigon tube and the leaves, respectively. In addition, researchers determined the total phenolic compounds, total tannins and total flavonoids of the *C. autumnale*. In this study, the estimated amounts of total phenolic compounds, tannins and flavonoids were 138.7 mg/l, 82 mg/l and 66.3 mg/L [18]. One study reported the spectrophotometric method for alkaloid assay based on the backbone structures. In this study, the alkaloid compounds were determined by complex with metal [19].

A study evaluated the total phenolic compounds and flavonoids of some medicinal plants by the Folin-Ciocalteu and aluminium chloride methods [20].

In this study, tropolone alkaloid profiles of the *C. kurdicum* were reported. In one study the amounts of colchicine-like compounds of some *Colchicum* species have been determined by HPLC method [11]. According to this study, *C. speciosum* and *C. macrophyllum* had higher amounts of colchicine.

In the literature, physicochemical properties of the plants have been determined. In this study, shape, powder colour, odour and taste of corms, solubility in the several solvents, percent of foreign matter, moisture contents, ashes and the amount of heavy metals of the *C. kurdicum* were evaluated.

In different studies, the physicochemical properties of some plants were evaluated. Organoleptic properties of the corm of *Colchicum luteum* have been explained ovoid shape and bitter taste. In addition, moisture content of *C. luteum* have been estimated 6% and total Ash, acid insoluble ash, water soluble ash and water insoluble ash were 1.33%, 6%, 1.24% and 4.28%, respectively. In this study, the moisture contents of *C. kurdicum* were calculated  $5.4 \pm 1.53\%$ .

Various ash values, which are used for determining the total properties of the compounds of medicinal plant, have been evaluated in the powdered corms of *C. kurdicum*.

Amount of heavy metals in the medicinal plants are the important parameter, which are determining the quality control. Heavy metals especially Hg, Cu, Cd and Pb have the standard amounts in plants and functional foods. In addition, high amounts of them induce the several complications such as cerebral and cardiovascular disorders. Based on standardization references, maximum amount of Hg, Cu, Cd and Pb were estimated 4, 6, 0.1 and 0.6  $\mu\text{g}/\text{kg}$ , respectively [21-26].

One study evaluated a novel method for rapid determining anti-inflammatory activities of the compounds. According to this study, anti-denaturation compounds are effective for improving the inflammatory conditions such as rheumatoid arthritis and osteoarthritis [27].

Researchers evaluated the *in vitro* anti-arthritic activities of the *Oryza sativa* var. Joha rice [14]. According to the study, the anti-inflammatory and anti-arthritic activity with anti-denaturation effects of the plant was 40 % to 60 %. Other researchers reported the anti-denaturation effects of the *Ziziphus oenoplia* between 50 % to 98 % for aqueous, ethanolic and ethyl acetate extracts [28]. In the present study, the anti-denaturation activity of the methanolic extract of *C. kurdicum* was estimated  $84.7 \pm 1.2$  % and was significant against the negative control. In addition, the anti-oxidant activity of the extract was calculated  $94.8 \pm 1.8$ %.

## 5. Conclusion

According to this study, the corm of the *C. kurdicum* had the main bioactive compounds and the appropriated physicochemical parameters. In addition, these compounds have the valuable biological activities especially anti-inflammatory and antioxidant activities, which could be used for treatment of inflammatory and oxidative diseases such as gout, rheumatoid arthritis, cardiovascular diseases and premature aging.

## References

1. Gulsoy-Toplan G, Goger F, Yildiz-Pekoz A, Gibbons S, Sariyar G and Mat A. Chemical constituents of the different parts of *Colchicum micranthum* and *C. chalcedonicum* and their cytotoxic activities. *Nat. Pro. Commun.* 2018; 13(5): 1-5.

So, *C. kurdicum* is the appropriated natural source for this valuable compounds and played the important role for drug discovery and design studies.

## Author contributions

Mohammad Azadbakht was the main study investigator, contributed to the collection of the data, and critically revised the manuscript; Ali Davoodi the study investigator, contributed to the collection of the data, and critically revised the manuscript; Seyed Jalal Hosseinimehr participated in the data interpretation and revised the manuscript; Saeed Emami, Masoud Azadbakht, Fatemeh Mirzaee and Hossein Bakhshi Joubari participated in the data interpretation.

## Conflict of interest

The authors declare that there is no conflict of interest.

## Acknowledgements

This article was a part of PhD thesis of Dr. Ali Davoodi, PharmD, supervised by Professor Mohammad Azadbakht. We thank all pharmacy school personnel especially pharmacognosy laboratory personals. This study was supported by a grant from Mazandaran University of Medical Sciences, Sari, Iran (Funding code: 1396-1102).

2. Alali FQ, El-Alali A, Tawaha K and El-Elimat T. Seasonal variation of colchicine content in *Colchicum brachyphyllum* and *Colchicum tunicatum* (Colchicaceae). *Nat. Prod. Res.* 2006; 20(12): 1121-8.


3. Al-Fayyad M, Alali F, Alkofahi A and Tell A. Determination of colchicine content in



- Colchicum hierosolymitanum* and *Colchicum tunicatum* under cultivation. *Nat. Prod. Let.* 2002; 16(6): 395-400.
4. Khan H, Tariq SA and Khan MA. Biological and phytochemical studies on corms of *Colchicum luteum* Baker. *J. Med. Plant. Res.* 2011; 5(32): 7031-5.
  5. Shafiee F, Khoshvishkaie E, Davoodi A, Dashti Kalantar A, Bakhshi Jouybari H and Ataee R. The determination of blood glucose lowering and metabolic effects of *Mespilus germanica* L. hydroacetic extract on Streptozocin-induced diabetic BALB/c mice. *Medicines* 2018; 5(1): 1-7.
  6. Davoodi A, Jouybari HB, Fathi H and Ebrahimnejad P. Formulation and physicochemical evaluation of medlar (*Mespilus germanica* L.) and oak (*Quercus castaneifolia* C.A.Mey.) Effervescent tablet. *Inter. J. Pharmaceut. Sci. Res.* 2018; 9(9): 3870-5.
  7. Fakhri M, Davoodi A, Parviz M, Sadeghi Ghadi Z, Mousavinasab SN, Farhadi R, Azadbakht M and Azadbakht M. Characterization and HPLC analysis of manna from some cotoneaster species. *Int. J. Pharma. Sci. Res.* 2017; 8: 1000-06.
  8. Pramparo M, Gregory S and Mattea M. Immersion vs. percolation in the extraction of oil from oleaginous seeds. *J. Amer. Oil Chem. Soc.* 2002; 79(10):955-60.
  9. Becer E, Hanoğlu DY, Kabadayı H, Hanoğlu A, Vatansever S, Yavuz DÖ, et al. The effect of *Colchicum pusillum* in human colon cancer cells via Wnt/ $\beta$ -catenin pathway. *Gene.* 2019; 686: 213-9.
  10. Bayrak S, Sökmen M, Aytaç E and Sökmen A. Conventional and supercritical fluid extraction (SFE) of colchicine from *Colchicum speciosum*. *Ind. Crop. Prod.* 2019; 128: 80-4.
  11. Ondra P, Válka I, Vičar J, Sütülpinar N and Šimánek V. Chromatographic determination of constituents of the genus *Colchicum* (Liliaceae). *J. Chrom. A.* 1995; 704(2): 351-6.
  12. Organization WH. Quality control methods for medicinal plant materials. *WHO Library Cataloguing in Publication Data.* 1998; 1: 1-123.
  13. Shokrzadeh M, Fathalinezhad F and Khoshvishkaie E. Concentrations of heavy metals (Cr, Cd, Pb) in three types of rice in Astaneh-Ashrafieh, Iran. *J. Maz. Univ. Med. Sci.* 2017; 26(146): 196-200.
  14. Rahman H, Eswaraiah MC and Dutta A. *In-vitro* anti-inflammatory and anti-arthritic activity of *Oryza sativa* var. Joha rice (an aromatic indigenous rice of Assam). *Am. Eurasian J. Agric. Environ. Sci.* 2015; 15: 115-21.
  15. Sharma OP and Bhat TK. DPPH antioxidant assay revisited. *Food Chem.* 2009; 113(4): 1202-5.
  16. Senthilkumar M. Phytochemical screening and antibacterial activity of *Gloriosa superba* Linn. *Inter. J. Pharmacogn. Phytochem. Res.* 2013; 5(1): 31-6.
  17. Pırlıdar S, Sütülpinar N, Atasever B, Erdem-Kuruca S, Papouskova B and Šimánek V. Chemical constituents of the different parts of *Colchicum baytopiorum* and their cytotoxic activities on K562 and HL60 cell lines. *Pharmaceut. Biol.* 2010; 48(1): 32-9.
  18. Suica-Bunghez I-R, Ion R-M, Teodorescu S, Sorescu A-A, Stirbescu R-M and Stirbescu N-M. Fitochemical and antioxidant characterization of autumn crocus (*Colchicum autumnale*) flowers and roots plant extracts. *J. Sci. Art.* 2017; 17(3): 539-46.
  19. Sangster AW and Stuart KL. Ultraviolet spectra of alkaloids. *Chem. Rev.* 1965; 65(1): 69-130.
  20. Ebrahimzadeh MA, Enayatifard R, Khalili M, Ghaffarloo M, Saedi M and Charati JY.

- Correlation between sun protection factor and antioxidant activity, phenol and flavonoid contents of some medicinal plants. *IJPR*. 2014; 13(3): 1041-7.
21. Ahmed SN, Ahmad M, Shinwari ZK and Shinwari S. Taxonomic, pharmacognostic and physicochemical authentication of *Colchicum luteum* baker (suranjantalkh) from its commercial adulterant. *Pak. J. Bot.* 2016; 48(5): 2039-45.
22. Silva RJ, de Aguiar-Dias AC, do CF Faial K and de Mendonca MS. Morphoanatomical and physicochemical profile of *Piper callosum*: valuable assessment for its quality control. *Rev. Brasi. Farmacogn.* 2017; 27(1): 20-33.
23. Mohamad TAST, Naz H, Jalal RS, Hussin K, Rahman MRA, Adam A and Weber JFF. Chemical and pharmacognostical characterization of two Malaysian plants both known as Ajisamat. *Rev. Brasil. Farmacogn.* 2013; 23(5): 724-30.
24. Yazdinezhad A, Ramezanloo N and Mozaffari S. Pharmacognostic and phytochemical investigation of *Heracleum persicum* Desf. ex Fischer. *Res. J. Pharmacogn.* 2016; 3(2): 17-24.
25. Tüzen M. Determination of heavy metals in fish samples of the middle Black Sea (Turkey) by graphite furnace atomic absorption spectrometry. *Food Chem.* 2003; 80(1): 119-23.
26. Sivaperumal P, Sankar T and Nair PV. Heavy metal concentrations in fish, shellfish and fish products from internal markets of India vis-a-vis international standards. *Food Chem.* 2007; 102(3):612-20.
27. Williams LAD, O'Connar A, Latore L, Dennis O, Ringer S, Whittaker JA, Conrad J, Volger B, Rosner H and Kraus W. The in vitro anti-denaturation effects induced by natural products and non-steroidal compounds in heat treated (immunogenic) bovine serum albumin is proposed as a screening assay for the detection of anti-inflammatory compounds, without the use of animals, in the early stages of the drug discovery process. *West Ind. Med. J.* 2008; 57(4): 327-31.
28. Ramalingam R, Madhavi BB, Nath AR, Duganath N, Sri EU and Banji D. *In-vitro* anti-denaturation and antibacterial activities of *Zizyphus oenoplia*. *Pharm. Lett.* 2010; 2(1): 87-93.

How to cite this article: Azadbakht M, Davoodi A, Hosseinimehr SJ, Emami S, Azadbakht M, Mirzaee F and Bakhshi Jouybari H. Phytochemical, physicochemical and biological evaluation of *Colchicum kurdicum* (Bornm.) Stef.: a study on materia medica of Persian medicine. *Journal of Medicinal Plants* 2020; 19(76): 36-45.  
doi: 10.29252/jmp.19.76.36



## فصلنامه گیاهان دارویی

Journal homepage: [www.jmp.ir](http://www.jmp.ir)پژوهشکده گیاهان دارویی  
جهاد دانشگاهی

## مقاله تحقیقاتی

## ارزیابی فیتوشیمیایی، فیزیکوشیمیایی و بیولوژیکی گل حسرت کردی: یک مطالعه بر روی مفرده پزشکی در طب ایرانی

محمد آزادبخت<sup>۱</sup>، علی داوودی<sup>۱\*</sup>، سید جلال حسینی مهر<sup>۲</sup>، سعید امامی<sup>۳</sup>، مسعود آزادبخت<sup>۴</sup>، فاطمه میرزائی<sup>۱</sup>، حسین بخشی جویباری<sup>۱</sup><sup>۱</sup> گروه فارماکونوزی، دانشگاه علوم پزشکی مازندران، دانشکده داروسازی ساری، مازندران، ایران<sup>۲</sup> گروه داروسازی هسته‌ای، دانشگاه علوم پزشکی مازندران، دانشکده داروسازی ساری، مازندران، ایران<sup>۳</sup> گروه شیمی دارویی، دانشگاه علوم پزشکی مازندران، دانشکده داروسازی ساری، مازندران، ایران<sup>۴</sup> گروه سیستماتیک گیاهی، موسسه آموزش عالی سنا، مازندران، ایران

## چکیده

## اطلاعات مقاله

گل‌واژگان:

گل حسرت کردی

آلکالوئیدهای تروپولونی

فلاونوئید تام

کروماتوگرافی

آنتی‌اکسیدان

فعالیت ضدالتهابی

**مقدمه:** گیاهان جنس سورنجان (*Colchicum*)، گیاهانی تک‌لپه‌ای، گل‌دار و چند ساله بوده که بیش از ۱۶۰ گونه در جهان دارد. همچنین، این گیاه در طب سنتی و مدرن کاربرد زیادی دارد. **هدف:** هدف از این مطالعه ارزیابی فیتوشیمیایی و فیزیکوشیمیایی، فعالیت‌های آنتی‌اکسیدانی و ضدالتهابی بنه گل حسرت کردی می‌باشد. **روش بررسی:** مقادیر تام آلکالوئیدهای تروپولونی، فنول، تانن و فلاونوئید با روش اسپکتروفتومتری تعیین شد. همچنین، آلکالوئیدهای تروپولونی با روش HPLC آنالیز شدند. خصوصیات فیزیکوشیمیایی شامل خواص ماکروسکوپیکی و ارگانولپتیکی، حالیت، ماده خارجی، مقادیر خاکستر و فلزات سنگین بر اساس پروتکل‌های فارماکوپه‌ای مورد بررسی قرار گرفتند. فعالیت‌های ضدالتهابی و آنتی‌اکسیدانی گیاه نیز تعیین شدند. **نتایج:** آلکالوئیدها، ترکیبات فنلی، فلاونوئیدها، تانن‌ها، ترپنوئیدها و استروئیدها، کومارین‌ها، ساپونین‌ها و گلیکوزیدها در بنه سورنجان تشخیص داده شدند. علاوه بر این، میزان تام آلکالوئیدهای تروپولونی، فنل، تانن و محتوای کل فلاونوئید بنه به ترتیب  $0/052 \pm 0/005$ ،  $0/028 \pm 0/025$  و  $0/652 \pm 0/021$ ،  $0/34 \pm 0/426$ ،  $0/028 \pm 0/025$  و  $0/652 \pm 0/021$  گرم محاسبه شد. در آنالیز HPLC، N-داستیل-N-فرمیل کلشی‌سین، کلشی‌فولین، کلشی‌کوزید و کورنی‌ژرین بیشترین میزان محاسبه شدند. خصوصیات فیزیکوشیمیایی در محدوده مناسبی تعیین شد. علاوه بر این، فعالیت‌های ضدالتهابی و آنتی‌اکسیدانی در ۱۰ میلی‌گرم در میلی‌لیتر عصاره متانولی به ترتیب  $1/2 \pm 84/7$  و  $1/8 \pm 94/8$  درصد تخمین زده شد. **نتیجه‌گیری:** بنه گل حسرت کردی حاوی ترکیبات فعال زیستی بود. همچنین، این ترکیبات دارای فعالیت‌های بیولوژیکی با ارزش بویژه فعالیت‌های ضدالتهابی و آنتی‌اکسیدانی بودند.

مخفف‌ها: HPLC، کروماتوگرافی مایع با کارایی بالا؛ PVPP، پلی‌وینیل‌پلی‌پیرولیدون؛ ACN، استونیتریل؛ SD، انحراف استاندارد

\* نویسنده مسؤول: [ali.davoodi@mazums.ac.ir](mailto:ali.davoodi@mazums.ac.ir)

تاریخ دریافت: ۲۱ مهر ۱۳۹۸؛ تاریخ دریافت اصلاحات: ۴ تیر ۱۳۹۹؛ تاریخ پذیرش: ۲۹ تیر ۱۳۹۹

doi: 10.29252/jmp.19.76.36

© 2020. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>)