

Protective Effects of Quercetin on Spermatogenesis in Streptozotocin-induced Diabetic Rat

Khaki A (D.V.M-Ph.D.)^{1*}, Nouri M (Ph.D.)², Fathiazad F (Ph.D.)³, Ahmadi-Ashtiani HR (Ph.D. student)⁴, Rastgar H (Ph.D.)⁵, Rezazadeh Sh (Ph.D.)⁶

1- Department of Veterinary Pathology (YRC), Islamic Azad University Tabriz Branch, Iran

2- Department of Biochemistry, Tabriz University of Medical Sciences, Iran

3- Department of Pharmacognosy Tabriz University of Medical Sciences, Iran

4- Clinical Biochemistry, Tarbiat-e-Modarres University, School of Medical Science, Department of Clinical Biochemistry, Tehran, Iran and Biochemistry & Nutrition Department of Zanjan Medical University

5- Ministry of Health and Medical Education, Food & Drug Laboratory Research center, Tehran, Iran

6- Department of Pharmacognosy and Pharmaceutics, Institute of Medicinal Plants, ACECR, Tehran, Iran

*Corresponding author: No. 1 Etekali La, South Shariati Street, 5138815941, Tabriz, Iran

Mobile Phone: +98-9143138-399

E - Mail: arashkhaki@yahoo.com

Receive: 22 Nov. 2008

Acceptance: 11 Mar. 2009

Abstract

Background: Quercetin is a strong antioxidant and long-term treatment of STZ-diabetic animals and it has been shown to reduce oxidative stress.

Objective: antioxidants have essential effect on spermatogenesis and sperm parameters. Enhanced oxidative stress and changes in antioxidant capacity are considered to play an important role in the pathogenesis of chronic diabetes mellitus.

Methods: Wistar male rat (n=40) were allocated into three groups, control group (n=10) and Quercetin (QR) group that received 15mg/kg (IP) QR, (n=10), and Diabetic group that received 55 mg/kg (IP) streptozotocin (STZ) (n=20) which was subdivided to two groups of 10; STZ group and treatment group. Treatment group received 55 mg/kg (IP) STZ plus 15mg/kg QR, daily for 4 weeks, respectively; however, the control group just received an equal volume of distilled water daily (IP). Diabetes was induced by a single intra peritonea injection of streptozotocin (55 mg/kg). Animals were kept in standard condition. In 28 day after inducing diabetic 5cc blood were collected for testosterone, TAC, MDA and Ox-LDL levels and testes tissues of Rat in whole groups were removed and sperm was collected from epididymis then prepared for analysis.

Results: Sperm population, percentage of sperm viability and motility significantly increased in group that has received 15 mg/kg (IP) Quercetin (p<0.05) in comparison to control and experimental groups.

Conclusion: Since in our study 15 mg/kg (IP) Quercetin have significantly Preventive effect on Sperm percentage of viability and motility by reducing level of Reactive Oxygen Species (ROS) in serum, so it seems that using it can be effective for sperm healthy parameters in Diabetic Rat.

Keywords: Quercetin, Streptozotocin, Spermatogenesis, Rat

Introduction

Diabetes has been associated with reproductive impairment in both men and women. About 90% of diabetic patients have disturbances in sexual function, including a decrease in libido, impotence and infertility [1, 2] Due attention has been paid to the search of effective drugs in the field of traditional Chinese medicine (TCM). Diabetes mellitus is a group of syndromes characterized by hyperglycemia, altered metabolism of lipids, carbohydrates and proteins and an increased risk of complications from vascular diseases [3]. Enhanced oxidative stress and changes in antioxidant capacity are considered to play an important role in the pathogenesis of chronic diabetes mellitus [4, 5]. Although the mechanisms underlying the alterations associated with diabetes mellitus are presently not well understood, hyperglycemia lead patients to increased oxidative stress because the production of several reducing sugars (through glycolysis and the polyol pathway) is enhanced [6]. These reducing sugars can easily react with lipids and proteins (nonenzymatic glycation reaction), increasing the production of reactive oxygen species (ROS) [6]. Diabetes is the most common endocrine disease that leads to metabolic abnormalities involving regulation of carbohydrate metabolism. In addition to imbalanced carbohydrate metabolism, yet another major concern in diabetes is increased oxidative stress. increased production of free radicals or ROS formation may induce oxidized LDL (Ox-LDL), which is key step in the sequence of events leading to atherosclerosis Sustained hyperglycemia and increased oxidative stress, are the major players in the development of secondary complications in diabetes. These abnormalities produce pathologies including vasculopathies, neuropathies, ophthalmopathies

and nephropathies, among many other medical derangements [7]. The balance of ROS and antioxidant is a major mechanism in preventing damage by oxidative stress. Therefore, the dietary supplement of antioxidants such as vitamins, flavonoids has been used to prevent the occurrence of many chronic diseases [8]. Quercetin is a well-known flavonoid and a strong antioxidant and long-term treatment of STZ-diabetic animals and it has been shown to reduce oxidative stress [9]. Since streptozotocin causes testicular dysfunction and degeneration under situations of experimentally induced diabetes in animal models [10], it is hypothesized that Quercetin can decreasing STZ harmful effect on testis and sperm parameters by reducing reactive oxygen species (ROS). According to our systematic study on effect of *Allium cepa* and other medicinal plant on spermatogenesis [11] we want to study the effect of Quercetin as a active component of *Allium cepa* on spermatogenesis to confirm wither the single component could be effective as much as the total extracts.

Materials and Methods

Animals

Forty adult Wistar albino male rats were 8 weeks old and weighing 250 ± 10 g, they were obtained from animal facility of pasture institute of Iran. Male rats were housed in temperature controlled rooms (25°C) with constant humidity (40 - 70%) and 12h/12h light/ dark cycle prior to use in experimental protocols. All animals were treated in accordance to the Principles of Laboratory Animal Care. The experimental protocol was approved by the Animal Ethical Committee in accordance with the guide for the care and use of laboratory animals prepared by Tabriz medical University. All Rats were fed a

standard diet and water. The daily intake of animal water was monitored at least one week prior to start of treatments in order to determine the amount of water needed per experimental animal. Thereafter, the rats were randomly selected and divided into control (n=10) and Quercetin (QR) group that received 15mg/kg QR (IP), (n=10), and Diabetic group that received 55mg/kg (IP) streptozotocin (STZ) (n=20) which was subdivided to two groups of 10; STZ group and treatment group. Treatment group received 55mg/kg (IP) STZ plus 15mg/kg QR (IP). The control group just received an equal volume of 1cc distilled water daily (IP). Diabetes was induced by a single intra peritoneal (I.P) injection of streptozotocin (STZ, Sigma- U.S.A.) in 0.1 M citrate buffer (pH 4.0) at a dose of 55 mg/kg body weight. Quercetin (QR) injections were continued to the end of the study (for 4 weeks), [12].

Induction of experimental type 1, Diabetes

Experimental type 1 diabetes was induced in rats by intra peritoneal (I.P) injection of 55 mg/kg streptozotocin (STZ) in distilled water. Control rats were received distilled water, only, [13].

Blood glucose determination

Blood samples were collected from the tail vein. Basal glucose levels were determined prior to STZ injection, using an automated blood glucose analyzer (Glucometer Elite XL). Sample collections were then made 48 h after STZ injection and blood glucose concentrations were determined and compared between groups. Rats with blood glucose concentrations above 300 mg/dl were declared diabetic and were used in the experimental group. One week after the induction of experimental diabetes, protocol was started.

Quercetin preparation

Quercetin powder was obtained from Sigma Chemical Company (St. Louis, MO, USA). It was dissolved and diluted with 20% glycerol in 0.9 % normal saline, mixed vigorously and stored in a dark bottle at 4°C. The quercetin solution was freshly prepared each week.

Surgical Procedure

In the 28th day, (at the end of the treatment period), the rats were killed with diethyl ether, and the testes in control & experimental groups were immediately removed. The weights of testes in all study groups were recorded.

Sperm analysis

Sperms from the cauda epididymis were released by cutting into 2 ml of medium (Hams F10) containing 0.5 % bovine serum albumin. After 5 min incubation at 37°C (with 5 % CO₂), the cauda epididymis sperm reserves were determined using the standard hemocytometric method and sperm motility was analyzed with microscope (Olympus IX70) at 10 field and reported as mean of motile sperm according to WHO method (WHO). The sperm abnormality was evaluated according to a standard method of Khaki et al [13]. Briefly the smears of sperm suspension were made on clean glass slides and stained with periodic acid-Schiff's reaction hematoxylin. The stained smears were observed under a light microscopic with 40X objective. Sperms were classified into normal and abnormal. The total sperm abnormality was expressed as percentage incidence.

Measurement of Serum Total Antioxidant capacity (TAS)

TAS was measured in serum by means of a

commercial kit (Randox Co-England). The assay is based on the incubation of 2, 2'-azino-di-(3-ethylbenzthiazoline sulphonate) (ABTS) with a peroxidase (methmyoglobin) and hydrogen peroxide to produce the radical cation ABTS⁺, which has a relatively stable blue-green color, measured at 600 nm. The suppression of the color is compared with that of the Trolox, which is widely used as a traditional standard for TAS measurement assays, and the assay results are expressed as Trolox equivalent (mmol/L) [14].

Measurement of Serum MDA

Tissue MDA levels were determined by the thiobarbituric acid (TBA) method and expressed as nmol MDA formed/mL. Plasma MDA concentrations were determined with spectrophotometer. A calibration curve was prepared by using 1, 1',3,3'-tetramethoxypropane as the standard [15].

Measurement of Ox-LDL

Oxidized LDL level was measured by using a Mercodia Oxidized LDL ELISA kit (Lot No. 15904; Mercodia, Uppsala, Sweden). Mercodia Oxidized LDL Competitive ELISA is based on the monoclonal antibody 4E6 [16, 17].

Statistical analysis

Statistical analysis was done using the ANOVA and test for comparison of data in the control group with the experimental groups. The results were expressed as mean \pm S.E.M (standard error of means). P-value less than 0.05 were considered significant and are written in the parentheses.

Results

Results of sperm motility, viability and count

Streptozotocin-induced diabetic model by 55 mg/kg significantly decreased sperm count, motility and viability in diabetic group as

compared with those observed in the control and other experimental groups. The sperm concentrations, motility and vitality were (23.81 ± 3.20 and 10.05 ± 6.88 and 43.26 ± 2.33) in STZ group and the corresponding values in QR group were (47.05 ± 5.70 , 35.42 ± 6.88 and 67.05 ± 5.11) and the corresponding values in treatment (STZ+QR) group were (42.03 ± 5.20 , 30.64 ± 3.01 and 57.25 ± 4.22). However, the sperm concentrations, motility and vitality in control group were (48.68 ± 7.70 , 33.75 ± 6.88 and 66.25 ± 4.73). There weren't any significant change in sperm abnormality in experimental groups as compared with control group, (Table 1).

Results of Total Antioxidant capacity (TAC) measurement in Serum

The mean concentration of Total Antioxidant capacity (TAC) showed significant increase ($p < 0.05$) in QR group (0.75 ± 0.03) as compared with control group (0.70 ± 0.03) and STZ (0.32 ± 0.04) and STZ+QR (0.61 ± 0.05) groups.

Results of Malondialdehyde (MDA) measurement in Serum

Malondialdehyde (MDA) level showed significant ($p < 0.05$) decrease in QR group (0.30 ± 0.212) and control group (0.25 ± 0.04) in comparison to STZ (4.1 ± 0.06) and STZ+QR (1.1 ± 0.08) groups.

Results of Ox-LDL concentration measurement in Serum

Ox-LDL level increased) in STZ (5.6 ± 0.85) and STZ+QR (4.9 ± 0.80) groups as compared with control (3.1 ± 0.05) and QR (3.0 ± 0.45) groups. (Table 1).

Discussion

It is noteworthy that diabetes-related alterations in Leydig cells are also related to



Table 1: The effect of the 55 mg/kg (IP) streptozotocin, 15 mg/kg (IP) Quercetin, 55 mg/kg (IP) streptozotocin plus 15 mg/kg Quercetin (Treatment), daily for 30 days on sperm parameters TAS, MDA, Ox-LDL level of control and experimental groups in the rats.

Groups	Control	Quercetin (15mg/kg(IP))	Streptozotocin (55mg/kg (IP))	Treatment 55mg/kg (IP) streptozotocin plus 15mg/kg Quercetin (n=10)
	(n=10)	(n=10)	(n=10)	(n=10)
Sperm Concentration (total count) (No of sperm/rat×10 ⁶)	48.68±7.70	47.05±5.70 (0.301)	23.81±3.20 * (0.001)	42.03±5.20 * (0.009)
Motility (%)	33.75±6.88	35.42±6.88 (0.231)	10.05±6.88 * (0.001)	30.64±3.01 * (0.008)
Viability (%)	66.25±4.73	67.05±5.11 (0.403)	43.26±2.33 * (0.007)	57.25±4.22 * (0.009)
Sperm Abnormality (%)	4.22±0.666	4.20±0.618	6.27±0.711	5.12±0.656
Total Antioxidant Capacity (TAS) (nmol/ml)	0.70±0.03	0.75±0.03 * (0.006)	0.32±0.04 * (0.003)	0.61±0.05 * (0.004)
Malondialdehyde (MDA) (nmol/ml)	0.25±0.04	0.30±0.212 * (0.004)	4.1±0.06 * (0.008)	1.1±0.08 * (0.005)
LDL-OX(u/l)	3.1±0.05	3.0±0.45 (0.108)	5.6±0.85 (0.061)	4.9±0.80 (0.055)

Data are presented as mean ± SE.

* P-value less than 0.05 were considered significant and are writing in the parentheses, (compared with the control group).

changes in the pituitary-testicular axis [18, 19, 20, 21, 22]. Thus, this disease induces a decrease in the serum levels of luteinizing hormone (LH), which is responsible for normal Leydig cell function [22, 23]. Diabetic testicular dysfunction might be transient or permanent depending on the degree and duration of the disease. Erectile dysfunction (ED) is a well-recognized complication of diabetes mellitus (DM). The low incidence of diabetes in infertile patients might be the reason for the limited amount of current research [24]. However, an altered testicular axis was noted in experimental studies Seethalakshmi et al. [25] found that testicular weight, sperm count and motility significantly decreased in diabetic rats. Moreover, Cameron et al. [26] defined increasing tubule wall thickness, germ cell depletion and Sertoli cell vacuolization in diabetic human testicular

biopsies and in diabetic rats. Results in same study showed that diabetes induces a clear impairment of reproductive performance in rats and tungstate treatment in these diabetic rats leads to a recovery of reproductive performance by increasing the number of Leydig cells [27]. Oxidative stress plays a role in the development of diabetic complications [7]. Oxidative damage was ascertained by measuring the malondialdehyde levels, reactive oxygen species (ROS) generation, alterations in antioxidant defences and specifies the level of oxidized-LDL. Our results showed sperm count and motility are significantly decreased and Expansion of interstitial space with vacuolization and Leydig cells had an abnormal fibroblast-like appearance. The measurement of TAC levels showed marked decrease in streptozotocin-induced diabetic group as compared with those

seen in the control and other experimental groups and these results were in agreement with Tang et al [28] research that showed testicular injury and apoptosis induced by diabetes are partially attributed to the augmented oxidative stress in testicular tissue. The dietary intake of flavonoids in humans has been estimated to be 16 – 1000 mg/day. Quercetin is regularly consumed by humans as it is the major flavonoid found in human diet [29, 30]. This flavonoid is reported to decrease capillary fragility, to protect against diabetic cataracts, to possess antiviral and anti-allergenic activities, to inhibit platelet aggregation and the oxidation of low density lipoproteins, and to act as an anti-inflammatory agent [31]. Quercetin an important flavonoid possesses beneficial effects on health due to its antioxidant function. One mechanism of the antioxidant action of quercetin was involved in scavenging free radicals, such as superoxide radicals generated by xanthine/ xanthine

oxidase. Previous studies indicated that reactive oxygen substances may be involved in possible testicular complications in streptozotocin-induced diabetic of rats [32]. Since, in our study quercetin could significantly improve epididymal sperm quality and decrease serum Reactive Oxygen Species (ROS) and ox-LDL in Streptozotocin-induced diabetic rats, and has beneficial effect on antioxidants and decreases the risk of degenerative diseases. We suggest using dietary plants, fruits, vegetables, onion, teas, and red wine rich of flavonoids and Quercetin which could have beneficial effects on diabetic persons and decrease risk of infertility in men.

Acknowledgments

We thank the staff at Islamic Azad University of Tabriz for their help and financial support in the preparation of this manuscript.

References

1. Jiang GY. Practical Diabetes. 1st Edition. Beijing: People's Health Publishing House. 1996: 295.
2. Shi-Liang FENG, Shu-Hua LI, Yan WANG, Chang-Chun CHEN, Bin GAO. Effect of ligustrum fruit extract on reproduction in experimental diabetic rats. *Asian J. Androl.* 2001; 3: 71 - 3.
3. Davis SN. Insulin, Oral Hypoglycemic Agents and the Pharmacology of the Endocrine Pancreas. In: Goodman and Gilman's the Pharmacological Basis of Therapeutics. Brunton, L. L. (Ed.). *McGraw-Hill, New York.* 2006; 1613 - 45.
4. Baynes JW, Thorpe SR. Role of oxidative stress in diabetic complications: a new perspective on an old paradigm. *J. of Diabetes.* 1999; 48: 1 - 9.
5. Wolff SP, Jiang ZY, Hunt JV. Protein glycation and oxidative stress in diabetes mellitus and ageing. *J. Free Radic Biol. Med.* 1991; 10: 339 - 52.
6. Palmeira CM, Santos DL, Seica R, Moreno AJ, Santos MS. Enhanced mitochondrial testicular antioxidant capacity in Goto-Kakizaki diabetic rats: role of coenzyme Q. *Am. J. Physiol. J. Cell Physiol.* 2001; 281: C1023 - 8.



7. Sexton WJ, Jarow JP. Effect of diabetes mellitus upon male reproductive function. *J. of Urology* 1997; 49: 508 - 13.
8. Peluso MR. Flavonoids attenuate cardiovascular disease, inhibit phosphodiesterase, and modulate lipid homeostasis in adipose tissue and liver. *J. of Exp. Biol. Med.* (Maywood). 2006; 231 (8): 1287 - 99.
9. Mahesh T, Menon VP. Quercetin alleviates oxidative stress in streptozotocin-induced diabetic rats. *J. of Phytother. Res.* 2004; 18: 123 - 7.
10. Shrilatha B, Muralidhara. Early oxidative stress in testis and epididymal sperm in streptozotocin-induced diabetic mice: its progression and genotoxic consequences. *J. of Reprod. Toxicol.* 2007; 23 (4): 578 - 87.
11. Khaki A, Fathiazad F, Nouri M, Khaki AA, Jabarikh H, Hammadeh M. Evaluation of Androgenic Activity of *Allium cepa* on Spermatogenesis in Rat. *J. of Folia Morphol.* (Warsz). 2009; 68: 45 - 51.
12. Coskun O, Kanter M, Korkmaz A, Oter S. Quercetin, a flavonoid antioxidant, prevents and protects streptozotocin-induced oxidative stress and beta-cell damage in rat pancreas. *J. of Pharmacol. Res.* 2005; 51 (2): 117 - 23.
13. Khaki A, Novin MG, Khaki AA, Nouri M, Sanati E, Nikmanesh M. Comparative study of the effects of gentamicin, neomycin, streptomycin and ofloxacin antibiotics on sperm parameters and testis apoptosis in rats. *Pak J. Biol. Sci.* 2008; 11 (13): 1683 - 9.
14. Feng R, He W, Ochi H. A new murine oxidative stress model associated with senescence. *J. of Mech Ageing Dev.* 2001; 122: 547 - 59.
15. Quintanilha AT, Packer L, Davies JM, Racanelli TL, Davies KJ. Membrane effects of vitamin E deficiency: bioenergetics and surface charge density studies of skeletal muscle and liver mitochondria. *J. of Ann. NY Acad Sci.* 1982; 393: 32 - 47.
16. Huang HFS, Linsenmeyer TA, Li MT, Giglio W, Anesetti R, von Hagen J, Ottenweller JE, Pogach L. Acute effects of spinal cord injury on the pituitary-testicular hormone axis and Sertoli cell functions: a time course study. *J. of Androl.* 1995; 16: 148 - 57.
17. Foglia VG, Rosner JM, Ramos M, Lema BE. Sexual disturbances in the male diabetic rat. *Horm Metab Res.* 1969; 1: 72 - 7.
18. Oksanen A. Testicular lesions of streptozotocin diabetic rats. *J. of Horm Res.* 1975; 6: 138 - 44.
19. Tesone M, Oliveira-Filho RM, Biella de Souza Valle L, Calvo JC, Baraño JLS, Foglia VG, Charreau EH. Androgen receptor in the diabetic rat. *J. of Diabetologia.* 1980; 18: 385 - 90.
20. Lin T, Haskell J, Vinson N, Terracio L. Characterization of insulin-like growth factor I receptors of purified Leydig cells and their role in steroidogenesis in primary culture: a comparative study. *J. of Endocrinol.* 1986; 119: 1641 - 7.
21. Feng HL, Jay PD, Sandlow JI, Sparks AET, Sandra A, Zheng LJ. Decreased expression of the c-kit receptor is associated with increased apoptosis in subfertile human testes. *J. of Fertil Steril.* 1999; 71: 85 - 9.
22. Steger RW, Rabe MB. The effect of diabetes mellitus on endocrine and reproductive function. *J. of Proc Soc. Exp. Biol. Med.* 1997; 214: 1 - 11.

- 23.** Benítez A, Pérez Díaz J. Effect of streptozotocin-diabetes and insulin treatment on regulation of Leydig cell function in the rat. *J. of Horm. Metab. Res.* 1985; 17: 5 - 7.
- 24.** Altay B, Cetinkalp S, Doganavsargil B, Hekimgil M, Semerci B. Streptozotocin-induced diabetic effects on spermatogenesis with proliferative cell nuclear antigen immunostaining of adult rat testis. *J. of Fertil Steril.* 2003; 80 (Suppl 2): 828 - 31.
- 25.** Seethalakshmi L, Menon M, Diamond D. The effect of streptozotocin-induced diabetes on the neuroendocrine-male reproductive tract axis of the adult rat. *J. of Urol.* 1987; 138: 190 - 4.
- 26.** Cameron DF, Murray FT, Drylie DD. Interstitial compartment pathology and spermatogenic disruption in testes from impotent diabetic men. *J. of Anat. Rec.* 1985; 213: 53 - 62.
- 27.** Ballester J, Domínguez J, Muñoz MC, Sensat M, Rigau T, Guinovart JJ, Rodríguez-Gil JE. Tungstate treatment improves Leydig cell function in streptozotocin-diabetic rats. *J. of Androl.* 2005; 26 (6): 706 - 15.
- 28.** Tang XY, Zhang Q, Dai DZ, Ying HJ, Wang QJ, Dai Y. Effects of strontium fructose 1,6-diphosphate on expression of apoptosis-related genes and oxidative stress in testes of diabetic rats. *Int. J. Urol.* 2008; 15 (3): 251 - 6.
- 29.** Skibola CF, Smith MT. Potential health impacts of excessive flavonoid intake. *J. of Free Radic. Biol. Med.* 2000; 29: 375 - 83.
- 30.** Manach C, Morand C, Crespy V, Demigne C, Texier O, Regeat F, Remesy C. Quercetin is recovered in human plasma as conjugated derivatives which retain antioxidant properties. *J. of FEBS Lett.* 1998; 426: 331 - 6.
- 31.** Bors W, Michel C, Stettmaier K. Antioxidant effects of flavonoids. *J. of Biofactors.* 1997; 6: 399 - 402.
- 32.** Naziroğlu M. Enhanced testicular antioxidant capacity in streptozotocin-induced diabetic rats: protective role of vitamins C and E and selenium. *J. of Biol. Trace Elem. Res.* 2003; 94 (1): 61 - 72.