

Journal of Medicinal Plants



Journal homepage: www.jmp.ir

Research Article

Identification of isoflavonoids in antioxidant effective fraction of *Arum rupicola* Boiss. leaves

Zahra Tofighi^{1,2}, Yasaman Shahpar², Alireza Taheri³, Saeed Tavakoli⁴, Raymond Asatouri⁵, Mahdieh Eftekhari⁶, Mahdi Vazirian², Mohammad Reza Shams Ardekani², Seyede Nargess Sadati lamardi^{7,*}

ARTICLE INFO

Keywords: Arum rupicola Antioxidant Cytotoxic effect Genistein Isoflavonoid

ABSTRACT

Background: Arum rupicola Boiss. (Araceae Family) is used by the native people of southern areas of Iran as a soup called "kardeh soup". Several flavonoids and phenol compounds have been identified from Arum species. Objective: The aim of this study was to evaluate antioxidant effect and total phenol contents as well as cytotoxic activity of the leaves of A. rupicola. Methods: Antioxidant activity of total methanol extract and fractions including n-hexane, chloroform, ethyl acetate and water residue were evaluated using FRAP and DPPH methods. Total phenol content was measured using Folin-Ciocalteu method. Cytotoxic activity of the extract and fractions were investigated against human breast cancer MCF-7, MDA-MB-231, and T47D cell lines by MTT assay. Further phytochemical isolation was done on the water residue using column chromatography. Results: According to the results, water residue showed the lowest IC₅₀ value (186.7 μg/ml) and the total methanol extract showed the most antioxidant power (163.62 mmol FeSO4/100 g extract) and phenol content (135 µmol Gallic acid/g extract). The hexane fraction also showed the highest cytotoxic effect against MCF-7 breast cancer cell line with IC₅₀ equal to 118.9 μg/ml. Phytochemical analysis of the water residue resulted in isolation and identification of three isoflavonoids named orobol, genistein and genistein 8-c-glucoside. Conclusion: Based on the identification of isoflavonoid compounds in this plant, its ability to be used as a phytoestrogenic supplement can be considered in future studies.

Abbreviations: FRAP, Ferric Reducing Ability of Plasma; DPPH, 2,2-Diphenyl-1-picrylhydrazyl; MTT, 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide

doi: 10.52547/jmp.20.79.14

Received 10 March 2021; Received in revised form 18 June 2021; Accepted 3 July 2021

© 2020. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (https://creativecommons.org/licenses/by-nc/4.0/)

¹ Medicinal Plants Research Center, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

² Department of Pharmacognosy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

³ Department of Medicinal Chemistry, Pharmaceutical Branch Islamic Azad University, Tehran, Iran

⁴ Medicinal Plants Research Center, Institute of Medicinal Plants, ACECR, Karaj, Iran

⁵ Department of Biology, Science and Research Branch Islamic Azad University, Tehran, Iran

⁶ Department of Pharmacognosy and Pharmaceutical Biology, Faculty of Pharmacy, Kermanshah University of Medical Sciences, Kermanshah, Iran

⁷Department of Traditional Pharmacy, School of Persian Medicine, Tehran University of Medical Sciences, Tehran, Iran

^{*} Corresponding author: n_sadati@tums.ac.ir

1. Introduction

Arum is a genus of flowering plants belonging to family Araceae, represented by 26 species and distributed in Northern Africa, Mediterranean Region, Western Asia, and Europe [1]. Several Arum species are growing in Lebanon, Cyprus, Syria and Palestine and consumed in folk medicine for treatment of cancer, cough, hemorrhoids, worms, constipation and urinary tract infections. The fresh leaves of Arum used in cooked or roasted form after washing several times with salted water [2]. Arum rupicola Boiss (Syn. Arum conophalloides Kotschy ex Schott) is used by the native people of southern areas of Iran as a soup called "kardeh soup" [3]. Also in some areas of Turkey the aerial parts of A. rupicola with local name of "kahri" were used as rice-vegetable dish [4]. Various pharmacological effects of Arum species have been reported such antioxidant, as antimicrobial effects [5-7]

A variety of flavonoids including, vitexin, caffeic acid, luteolin, vicenin and 3,6,8trimethoxy, 5,7,3',4'-tetrahydroxy flavone [8] and the alkaloid (S)-3,4,5-trihydroxy-1 H-pyrrol-2(5H)-one were isolated from A. palaestinum [9]. Also several flavonoids such as apigenin, luteolin, quercetin, vitexin, isoorientin, esculin, quercetin-3-O-β-glucoside, caffeic acid, and ferulic acid have been identified from A. dioscoridis [10, 11]. Flavonoids have antioxidant, anti-cancer, anti-allergic, antiinflammatory and anti-viral phytochemical compounds [12]. Since several flavonoids and phenolic compounds reported in the genus Arum, it can be expected that the aerial parts of the plant contain interesting biological active metabolites. The aim of this study was to evaluate phytochemicals, antioxidant effect and total phenolic contents of the leaves of Arum rupicola collected from southern areas of Iran.

2. Materials and Methods

2.1. Chemicals

2,2-Diphenyl 1-picrylhydrazyl (DPPH; Fluka, Switzerland); butylatedhydroxylanizole (BHA), methanol, hexane, chloroform, ethyl acetate (Merck, Germany) were purchased. 1 H and 13 C-NMR spectroscopy of compounds A, B and C were performed in DMSO-d₆ on a Bruker Avance DPX 400 spectrometer (Karlsruhe, Germany) [400 MHz, tetramethylsilane (TMS) as internal standard]. Column chromatography (CC) was done by Sephadex LH-20 (lipophilic Sephadex, 25–100 μ m; Sigma, Dorset, UK) columns.

2.2. Plant material

The leaves of *Arum rupicola* were collected from their growing habitats in Yasuj area, Iran, April 2016. The plant identified by Dr. Farideh Attar. A voucher specimen of the plant (46054 TUH) is deposited in the herbarium of the faculty of Sciences, Tehran University.

2.3. Extraction

The leaves of A. rupicola were air dried and grounded in the laboratory, and the obtained powder (1000 g) was extracted five times at room temperature with 80 % methanol for an interval of three days. The aqueous-methanol extract was evaporated under reduced pressure temperature to obtain a residue of 398.63 g. The total methanol extract was suspended in water and then fractionated using hexane (21.52 g), chloroform (3.03 g), and ethyl acetate (1.67 g) according to their polarity also water residue was The excessive solvents were evaporated with a rotary vacuum evaporator (60 rpm at 40 °C). All extracts were stored at 4 °C until use.

2.4. Total phenol contents assay

Total Phenol was determined colorimetrically using Folin-Ciocalteu reagent as described by Velioglu et al. 1998 and according to previous study [13]. The experiments were repeated three times. The phenol content calibration curves were plotted by measuring the absorption of certain concentraitions (25-150 mg/L) of gallic acid as a standard and the results were expressed as milligrams of gallic acid equivalents (GAE) per gram of dry matter (total extracts and fractions) as means ± SEM [13].

2.5. Antioxidant determination of DPPH assay

The total methanol extract was evaluated for its free radical scavenging activities using 2,2method diphenyl-1-picryl-hydrazyl (DPPH) according to Brand Williams et al. (1995) [14]. Different concentrations (25, 50, 100 µg/ml) of sample solutions (1 ml) in methanol were added to DPPH methanol solution (2 ml, 40 µg/ml). Butylated hydroxyanisole (BHA) (100 µg/ml) was used as positive control. After 30 min, the absorbance was measured at 517 nm. All tests were carried out in triple replicate. Percentage of radical scavenging activity of sample was calculated according to the following equation: Inhibition $\% = [(A0-As)/A0] \times 100$ that A0 is the absorbance of the control and as is the absorbance of the sample. Half maximal inhibitory concentration (IC₅₀) value (indicate the concentration of the sample (mg/ml), required to scavenge 50 % of DPPH) was calculated from the plotted graph of scavenging activity versus the concentration of extract, using linear regression analysis.

2.6. FRAP assay

Ferric reducing antioxidant power assay is based on reduction of a ferric-tripyridyl triazine complex to its ferrous colored form in the presence of antioxidants and was performed according to previous studies [15].

2.7. Cell culture

Human breast cancer cell lines including MCF-7 (human breast cancer cell line with estrogen, progesterone and glucocorticoid receptors), T47D (breast ductal carcinoma), and MDA-MB-231 (human breast adenocarcinoma), were supplied from the National Cell Bank of Iran Pasteur Institute, Tehran, Iran. Cancer cell lines were developed in RPMI-1640 medium accompanied with 10 % heat-inactivated fetal calf serum, 100 μg/ml streptomycin and 100 U/ml penicillin at 37 °C in a humidified atmosphere with 5 % CO₂.

2.8. Determination of cell viability by MTT assay

The in-vitro cytotoxic activity of all samples were assessed against three human breast cancer cell lines including MCF-7, T47D, and MDA-MB-231 cell lines using MTT colorimetric assay according to previous work [16].

Concisely, the concentration of 50, 100, 200, 400 and 500 µg/ml from all samples were prepared for each cell line. At first, samples were dissolved in DMSO (Dimethyl Sulfoxide) and further diluted with cell culture medium. The final DMSO concentration used was 0.1 % of total volume of the medium in all treatments, including the control group. In each plate, there were three control wells (cells without test extracts) and three blank wells (the medium with 0.1 % DMSO) for cell viability. Etoposide was used as a positive control for cytotoxicity. The plates were incubated for 72 h. At the end of the incubation period, the medium was removed and 200 µl phenol red free medium containing MTT (1 mg/ml), was added to wells, and followed by 4 h incubation. After incubation, the culture medium was then exchanged with 100 µl of DMSO and the absorbance of each well was measured by using a micro plate reader at 492 nm. For each sample, the concentration causing 50 % cell growth inhibition (IC₅₀) compared with the control was calculated from concentration response curves by regression analysis [16].

2.9. Purification and isolation of flavonoid compounds

In order to identify the main compounds in the plant, water residue was selected for purification due to proper weight and antioxidant effects. Fifteen grams of the water residue were loaded on a Sephadex LH-20 (25×5 cm) column and eluted with aqueous methanol (80 %). Fractions with a volume of 30 ml were collected from this column. The obtained fractions were compared using TLC method and the same fractions were combined. To gain more weight for each fraction, this step was performed several times, each time with about 5 g of water residue, and the same fractions were combined to afford 4 fractions. Compound **B** was isolated from the second column and for the final purification, Sephadex LH-20 column (80×1 cm) was used. The weight of compound B was about 20 mg. The R_f of compound **B** with the solvent system of formic acid/ acetic acid/water/ethyl acetate (11/11/26/100) on TLC sillicagel 60 F₂₅₄ was about 0.7.

To separate compound $\bf A$, the fractions No. 24-27 of the first column, the fractions 20-23 of the second column and the fractions 31-34 of the third column were combined, and first was loaded on the Sephadex LH-20 (70×2 cm) column and washed with aqueous methanol (80%). Then, the fraction No. 17-24 of this column was dried and for final purification, it was loaded on a Sephadex LH-20 (80×1 cm) column eluted with methanol (100 %) to afford compound $\bf A$ (30 mg). The Rf of compound $\bf A$ with the solvent

system of formic acid/ acetic acid/water/ethyl acetate (11/11/26/100) on TLC sillicagel 60 F₂₅₄ was about 0.8. To separate compound C, the fractions No. 35-37 of the first column, the fractions 28-33 of the second column and the fractions 37-39 of the third column were merged, and then was loaded twice on a Sephadex LH-20 (80×1 cm) column eluted with methanol (100 %) to afford compound C (3 mg). The R_f of compound C with the solvent system of formic acid/ acetic acid/water/ethyl acetate (11/11/26/100) on TLC sillicagel 60 F_{254} was about 0.6.

2.10. Chemical analysis

 1 H and 13 C-NMR spectroscopy of compounds **A, B** and **C** were recorded on a Brucker Avance 500 DRX (500 MHz) spectrometer. Chemical shifts are given in (ppm) DMSO using TMS as internal standard. Column chromatography (CC) was performed using Sephadex LH-20 (lipophilic Sephadex, 25–100 μ m; Sigma, Dorset, UK) columns.

Compound A (Orobol): $C_{15}H_{10}O_6$. ¹H-NMR (400 MHz, DMSO-d₆): $\delta = 8.47$ (1H, s, H-2), 7.34 (2H, m, H-5',6'), 6.83 (1H, d, J = 7.6 Hz, H-2'), 6.54 (1H, s, H-8), 6.37 (1H, s, H-6)

Compound B (Genistein): $C_{15}H_{10}O_5$. ¹H-NMR (400 MHz, DMSO-d₆): δ =8.07 (1H, s, H-2), 7.77 (2H, d, J = 7.2 Hz, H-2', 6'), 6.8 (2H, d, J = 7.2 Hz, H-3', 5'), 6.57 (1H, s, H-8), 6.46 (1H, s, H-6)

Compound C (**Genistein 8-***C***-glucoside**): $C_{21}H_{21}O_{10}$. UV (MeOH, λ_{max} , nm):265. ¹H-NMR (400 MHz, DMSO-d₆): δ =8.00 (1H, s, H-2), 7.93 (2H, d, J= 7.2 Hz, H-2',6'), 6.88 (2H, d, J= 7.2 Hz, H-3',5'), 6.59 (1H, s, H-6), 4.18 (1H, d, J=10 Hz, H-1")

2.11. Statistical analyses

The values were reported as mean \pm SD by SPSS and Excel 2010.

3. Results

3.1. DPPH radical scavenging activity

Free radical scavenging effects of total extract and different fractions from *Arum rupicola* leaves were assessed with 2, 2-diphenyl-1-picryl-hydrazyl (DPPH). IC₅₀ values were displayed in Table 1.

3.2. FRAP assay

Results of antioxidant effects of different fractions of the leaves extract using FRAP, are reported based on mmol Fe II/ 100 g of extract or

fraction. Ferric reducing antioxidant power of the extracts calculated using the calibration curve and regression equation of ferrous sulfate (R^2 = 0.996, y = 0.0008x-0.015). According to table 1, it was presented that antioxidant property of 100 grams of hexane, chloroform, ethyl acetate, and water residue as well as total methanol extract, had antioxidant activity equivalent to 140.17 \pm 4.03, 140.02 \pm 4.37, 110.01 \pm 3.9, 108.64 \pm 4.07 and 163.62 milimol FeSO₄, respectively.

3.3. Total phenol content

Total phenol content (μ mol of GAE/g of sample) as represented in table 2 varied from 116.55 to 135.00 μ mol of GAE/g sample using the standard curve of gallic acid ($R^2 = 0.9919$, y = 0.0067x - 0.0194).

Table 1. DPPH radical scavenging activity (IC₅₀ values), total phenolic content (TPC) and antioxidant power (mmol Fe/100 g sample) of different fractions from *Arum rupicola* leaves.

Sample	IC ₅₀ (μg/ml) ^a	TPC (μmol Gallic acid/g sample)	Antioxidant power (mmol Fe/100 g sample)
Hexane fraction	481.5 ± 2.26^{a}	120.6 ± 12.04	140.17 ± 4.03
Chloroform fraction	637.8 ± 11.3	133.2 ± 13.5	140.02 ± 4.37
Ethyl acetate fraction	441.2 ± 4.77	122.4 ± 12.38	110.01 ± 3.9
Water residue	186.7 ± 5.8	116.5 ± 9.7	108.64 ± 4.07
Total methanol extract	467.3 ± 2.19	135 ± 11.8	163.62 ± 4.42
ВНА	7.9 ± 0.06	-	
Vitamin A	14.2 ± 1.23	-	

^a IC₅₀ values represent an average of three independent experiments (mean \pm SD); BHA: Butylated hydroxyanisole.

3.4. In vitro cytotoxic assay

The cytotoxicity analysis of total methanolic extract, hexane, chloroform, ethyl acetate and water residue in breast cancer cell lines including MCF-7, MDA-MB, and T47D cell line was performed and the results were shown in Table 2.

Results showed that in MCF-7 cell line, hexane, chloroform fractions and total methanol extract showed cytotoxicity with IC_{50} equal to

 118.9 ± 0.38 , 258.5 ± 0.25 , 392.7 ± 0.25 µg/ml, respectively. In this cell line, etoposide showed cytotoxicity with IC₅₀ of 18.53 ± 0.24 µg/ml.

In the MDA-MB-231 cell line, hexane, chloroform fractions and total methanol extract have shown cytotoxicity with IC₅₀ of 137.2 \pm 0.22, 239.8 \pm 0.12, 357.3 \pm 0.11 µg/ml, respectively. Etoposide also showed cytotoxicity with IC₅₀ of 19.9 \pm 0.006 µg/ml.

In the T47D cell line, the fractions of chloroform, hexane, and total methanol extract as well as water residue showed cytotoxicity with IC₅₀ of 113.8 \pm 1.25, 122.8 \pm 0.2, 249.08 \pm 0.14, 399.3 \pm 0.6 μ g/ml), respectively. In this cell line, etoposide showed cytotoxicity with IC₅₀ of 23.3 \pm 0.096 μ g/ml.

3.5. Isolation of isoflavonoid compounds

Isolated compounds **A**, **B**, and **C** from the methanolic fraction of the total methanolic extract of leaves of *Arum rupicola* were identified by comparison of their NMR (¹H- & ¹³C-NMR) with those reported in the literatures [17-19]. Compounds **A**, **B** and **C** (Fig. 1) were identified as Orobol, Genistein and Genistein 8-C-glucoside, respectively.

Table 2. In vitro cytotoxic activity (IC₅₀, μg/ml) of total extract and fractions of *Arum rupicola* leaves against cancer and normal cell lines

Sample	IC ₅₀ (μg/ml) ^a MCF-7	IC ₅₀ (μg/ml) MDA-MB-231	IC ₅₀ (μg/ml) T-47D
Total methanol extract	392.7 ± 0.25	357.3 ± 0.11	249.08 ± 0.14
Hexane fraction	118.9 ± 0.38	137.2 ± 0.22	122.8 ± 0.2
Chloroform fraction	258.5 ± 0.25	239.8 ± 0.12	113.8 ± 1.25
Ethyl acetate fraction	> 500	> 500	> 500
Water residue	> 500	> 500	399.3 ± 0.6
Etoposide	18.53 ± 0.24	19.9 ± 0.006	23.3 ± 0.096

 $^{^{}a}$ IC₅₀ values represent an average of three independent experiments (mean \pm SD).

Fig. 1. Chemical structure of isoflavonos isolated from Arum rupicola leaves. A: Orobol, B: Genistein, and C: Genistein 8-C-glucoside

4. Discussion

Various flavonoids and phenolic compounds have been identified and reported in different species of *Arum* [20, 21], but no phytochemical studies have been performed on *A. rupicola*. In this study, antioxidant effects, cytotoxicity as well as purification and identification of secondary metabolites of *A. rupicola* leaves were

investigated using different chromatographic and spectroscopic methods.

According to the results, methanolic fraction showed higher radical scavenging effect and the total methanol extract of the plant exhibited higher antioxidant power and total phenol content than other fractions. In a study, antioxidant and antitumor activities of the ethanol extracts from different parts of six plants growing in Palestine, including *A. palaestinum* were evaluated and concluded that the studied plants showed different antioxidant abilities which were strongly associated with their phenolic contents [22].

The results of cytotoxicity study of plant samples on breast cancer cell lines showed that hexane fraction due to more toxicity in breast cancer cell lines could be considered for future studies. One study reported in vitro cytotoxic activity of different fractions and four flavonoid isolated (luteolin, compounds chrysoeriol, isoorientin, isovitexin) from the diethyl ether and ethyl acetate fraction of Arum palaestinum extract were investigated against four human carcinoma cell lines, epidermal carcinoma of larynx (Hep2), cervix (HeLa), liver (HepG2) and breast (MCF-7). Results showed that the fractionated extract and the isolated compounds showed significant antitumor activity against the four cell lines [1].

Phytochemical study of water residue of A. rupicola, using chromatographic methods, led to the isolation and identification of three isoflavonoid compounds, Orobol, Genistein and Genistein 8-C-glucoside. This is the first report on the occurrence of isoflavonoids in the genus Arum. Isoflavones, coumestans, stilbenes, and lignans important subclasses are of phytoestrogens. Isoflavones, which are more other phytoestrogens, studied than estrogen-like properties in mammals and are found in large quantities in soybean and its products [23, 24]. Isoflavonoids have also been reported in several plant families in addition to the leguminous family and attracted the attention of many specialists from phytochemistry and plant physiology to medicine and nutrition [25]. Several beneficial effects of isoflavones have cardioprotection, been reported, such as

osteoporosis prevention and antioxidant effects, which may be related to their phytoestrogenic effects [26]. Orobol is an isoflavone that is in small amount in soybean and is structurally similar to genistein, a largest isoflavone in soybean. Orobiol and genistein and their derivatives are known as multifunctional isoflavones with biological activities like neuroprotective, antiobesity and anti-cancer [27, 28].

The presence of these isoflavones in *A. rupicola* can be introduced it as a good source of phytoestrogens and antioxidant.

5. Conclusion

According to the results of this study, it can be concluded that the stronger antioxidant effect of water residue of *Arum rupicola* could be due to the presence of three isoflavonoid compounds called orobol, genistein and genistein 8-C-glucoside. Considering the biological effects of isoflavons, which are a class of phytoestrogens, it is recommended to study the effects of this plant on diseases related to estrogen deficiency and also to expand its use as a food source containing phytoestrogens.

Author contributions

Substantial contributions to design, analysis and interpretation of data: Z. T., M. V., M. E., and SN. SL; investigation: Y. S., A. T., S. T., and R. A.; drafting the article or revising: Y. S. and SN. SL.; final approval of the version to be published: Z. T. and MR. SA.

Conflict of interest

We declare that there is no conflict of interest.

Acknowledgment

This study was supported by Tehran University of Medical Sciences (TUMS) (Grant No: 95-01-56-31529).

References

- **1.** Farid MM, Hussein SR, Ibrahim LF, El Desouky MA, Elsayed AM, El Oqlah AA and Saker MM. Cytotoxic activity and phytochemical analysis of *Arum palaestinum* Boiss. *Asian Pac. J. Trop. Biomed.* 2015; 5(11): 944-7. doi: 10.1016/j.apjtb.2015.07.019.
- **2.** Jaradat N and Abualhasan M. Comparison of phytoconstituents, total phenol contents and free radical scavenging capacities between four *Arum* species from Jerusalem and Bethlehem. *Pharm Sci.* 2016; 22(2): 120-5.doi: 10.15171/PS.2016. 19.
- **3.** Derakhshanfar A, Moayedi J, Vahedi M and Valizadeh A. *Arum conophalloides* Aqueous Extract Induced Hepatotoxicity in Rat; Histopathological, Biochemical, and mir-122 Assessments. *Microrna*. 2020; 9(3): 224-231. doi: 10.2174/ 2211536608666191016142400. PMID: 31622226; PMCID: PMC7366011.
- **4.** Haghighi, H. Essential oil of the aleaves of *Arum conophalloides* (Araceae) from Iran. *IJPS*. 2016; 12(3): 11-16. doi: 10.22034/ijps.2016. 23837
- **5.** Farahmandfar R, Kenari RE, Asnaashari M, Shahrampour D and Bakhshandeh T. Bioactive compounds, antioxidant and antimicrobial activities of *Arum maculatum* leaves extracts as affected by various solvents and extraction methods. *Food Sci. Nutr.* 2019; 7: 465-475. doi: 10.1002/fsn3.815.
- **6.** Yabalak E. Radical scavenging activity and chemical composition of methanolic extract from *Arum dioscoridis* SM. var. *dioscoridis* and determination of its mineral and trace elements. *JOTCSA*. 2017; 5(1): 205-218. doi: 10.18596/jotcsa.350370.
- **7.** Al-Shmgani H, Kadri ZHM, Al-Halbosiy MMF and Dewir YH. Phytochemical analysis,

- cytotoxicity and antioxidant activity of cuckoo pint (*Arum maculatum*) leaf extract. *Acta Biol.* (*Szeged*) 2019; 63(2): 119-124.
- **8.** Afifi FU, Khalil E and Abdalla S. Effect of isoorientin isolated from *Arum palaestinum* on uterine smooth muscle of rats and guinea pigs. *J. Ethnopharmacol.* 1999 May; 65(2): 173-7. doi: 10.1016/s0378-8741(98)00147-0.
- **9.** El-Desouky SK, Kim KH, Ryu SY, Eweas AF, Gamal-Eldeen AM and Kim YK. A new pyrrole alkaloid isolated from *Arum palaestinum* Boiss. and its biological activities. *Arch. Pharm. Res.* 2007 Aug; 30(8): 927-31. doi: 10.1007/BF02993958.
- **10.** Afifi FU, Kasabri V, Litescu SC and Abaza IM. *In vitro* and *in vivo* comparison of the biological activities of two traditionally and widely used *Arum* species from Jordan: *Arum dioscoridis* Sibth & Sm. and *Arum palaestinum* Boiss. *Nat. Prod. Res.* 2016 Aug; 30(16): 1777-86. doi: 10.1080/14786419.2015.1072713.
- **11.** Karahan F, Kulak M, Urlu E, Gozuacik HG, Boyumez T, Sekeroglu N and Doganturk IH. Total phenolic content, ferric reducing and DPPH scavenging activity of *Arum dioscoridis*. *Nat. Prod. Res.* 2015; 29(17): 1678-83. doi: 10.1080/14786419.2014.991320.
- **12.** Panche AN, Diwan AD, Chandra SR. Flavonoids: an overview. *J. Nutr. Sci.* 2016 Dec 29; 5:e47. doi: 10.1017/jns.2016.41. PMID: 28620474; PMCID: PMC5465813.
- **13.** Velioglu Y, Mazza G, Gao L and Oomah B. Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *J. Agric. Food Chem.* 1998; 46(10): 4113-7. doi: 10.1021/jf9801973.

- **14.** Tofighi Z, Es-haghi A, Asl MM, Tajic AR, Navai MS, Tavakoli S, Hadjiakhoondi A and Yassa N. Investigation of chemical keys for relationship between plants and their unifloral honeys by hydrodistillation and SPME and biological activities of honeys. *Eur. Food Res. Tech.* 2014; 238(4): 665-73. doi: 10.1007/s00217-013-2147-y.
- **15.** Sadati N, Khanavi M, Mahrokh A, Nabavi S, Sohrabipour J and Hadjiakhoondi A. Comparison of antioxidant activity and total phenolic contents of some persian gulf marine algae. *J. Med. Plants* 2011; 1(37): 73-9. Available from: https://www.sid.ir/en/journal/ViewPaper.aspx?id=200886.
- **16.** Alipour E, Mousavi Z, Safaei Z, Pordeli M, Safavi M, Firoozpour L, Mohammadhosseini N, Saeedi M, Ardestani SK and Shafiee A. Synthesis and cytotoxic evaluation of some new [1,3] dioxolo [4,5-g] chromen-8-one derivatives. *DARU*. 2014; 22(1): 41. doi: 10.1186/2008-2231-22-41.
- **17.** Al-Mahweety JA. Phytochemical Studies on Medicinal Plants, Dracaenaceae resin, of Socotra Island, Yemen. *PSM Biological Res.* 2016; 1(2): 62-5. doi: 10.1016/j.fitote.2005.02.009.
- **18.** Fengqin W, JIANG K, Zuguang L. Purification and identification of genistein in *Ginkgo biloba* leaf extract. *Chin. J. Chromatogr.* 2007; 25(4): 509-13. doi: 10.1016/S1872-2059(07)60019-4.
- **19.** Sato S, Hiroe K, Kumazawa T and Jun-Ichi O. Total synthesis of two isoflavone C-glycosides: genistein and orobol 8-C-β-D-glucopyranosides. *Carbohydr Res.* 2006; 341(9): 1091-5. doi: 10.1055/s-0030-1257859.
- **20.** Afifi FU, Kasabri V, Litescu S, Abaza IF and Tawaha K. Phytochemical and Biological Evaluations of *Arum hygrophilum* Boiss.

- (Araceae). *Pharmacogn. Mag.* 2017 Apr-Jun; 13(50): 275-80. doi: 10.4103/0973-1296. 204551.
- **21.** Cole C, Burgoyne T, Lee A, Stehno-Bittel L and Zaid G. Erratum to: *Arum Palaestinum* with isovanillin, linolenic acid and beta-sitosterol inhibits prostate cancer spheroids and reduces the growth rate of prostate tumors in mice. *BMC Complement. Altern. Med.* 2015 Sep. 14; 15: 322. doi: 10.1186/s12906-015-0854-6.
- **22.** Husein AI, Ali-Shtayeh MS, Jondi WJ, Zatar NA, Abu-Reidah IM and Jamous RM. *In vitro* antioxidant and antitumor activities of six selected plants used in the traditional Arabic Palestinian herbal medicine. *Pharm. Biol.* 2014 Oct; 52(10): 1249-55. doi: 10.3109/13880209. 2014.886274.
- **23.** Rensen IV, Veit M, Wray V and Czygan F-C. Genistein-C-glucosides from *Genista cinerea*. *Nat. Prod. Lett.* 1995; 6(3): 203-7. doi: 10.1080/10575639508043160.
- **24.** King AMY, Young G. Characteristics and occurrence of phenolic phytochemicals. *J. Am. Diet. Assoc.* 1999; 99(2): 213-8. doi: 10.1016/S0002-8223(99)00051-6.
- **25.** Mackova Z, Koblovska R and Lapcik O. Distribution of isoflavonoids in non-leguminous taxa—an update. *Phytochemistry* 2006;67(9):849-55. doi: 10.1016/j.phytochem.2006.01.020.
- **26.** Rucinska A, Kirko S and Gabryelak T. Effect of the phytoestrogen, genistein-8-C-glucoside, on Chinese hamster ovary cells in vitro. *Cell Biol. Int.* 2007; 31(11): 1371-8.
- **27.** Hafidh RR. A comprehensive anticancer molecular study for genistein the promising anticancer drug. *J. Contemp. Med. Sci.* 2017; 3(11): 264-9. doi: 10.1016/j.cellbi.2007.05.012.
- **28.** Yang H, Lee SH, Ji H, Kim JE, Yoo R, Kim JH, Suk S, Huh CS, Park JHY, Heo YS, Shin HS,

[DOR: 20.1001.1.2717204.2021.20.79.2.7]

Kim BG and Lee KW. Orobol, an enzyme-convertible product of genistein, exerts antiobesity effects by targeting casein kinase 1 Epsilon. *Sci. Rep.* 2019; 9(1): 8942. doi: 10.1038/s41598-019-43950-9.

How to cite this article: Tofighi Z, Shahpar Y, Taheri AR, Tavakoli S, Asatouri R, Eftekhari M, Vazirian M, Shams Ardekani MR, Sadati lamardi SN. Identification of isoflavonoids in antioxidant effective fraction of *Arum rupicola* Boiss. leaves. *Journal of Medicinal Plants* 2021; 20(79): 14-23.

doi: 10.52547/jmp.20.79.14



فصلنامه گیاهان دارویی

Journal homepage: www.jmp.ir



مقاله تحقيقاتي

شناسایی ایزوفلاونوییدهای فراکسیون موثر آنتی اکسیدانی برگهای گونه ای شیپوری (Arum rupicola) زهرا توفیقی $^{','}$ ، یاسمن شهپر $^{'}$ ، علیرضا طاهری $^{'}$ ، سعید توکلی † ، رایموند آساتوری $^{\circ}$ ، مهدیه افتخاری $^{\circ}$ ، مهدی وزیریان $^{'}$ ، محمد رضا شمس اردکانی $^{'}$ ، سیده نرگس ساداتی لمردی $^{\circ}$.

ا مرکز تحقیقات گیاهان دارویی، دانشکده داروسازی، دانشگاه علوم پزشکی تهران، تهران، ایران

ٔ گروه فارماکوگنوزی، دانشکده داروسازی، دانشگاه علوم پزشکی تهران، تهران، ایران

" گروه شیمی دارویی، واحد علوم دارویی دانشگاه آزاد اسلامی، تهران، ایران

ً مرکز تحقیقات گیاهان دارویی، پژوهشکده گیاهان دارویی جهاد دانشگاهی، کرج، ایران

۵ گروه بیولوژی، واحد علوم و تحقیقات دانشگاه آزاد اسلامی، تهران، ایران

^۶ گروه فارماکو گنوزی و زیست فناوری دارویی، دانشکاه داروسازی، دانشگاه علوم پزشکی کرمانشاه، کرمانشاه، ایران

^۷گروه داروسازی سنتی، دانشکده طب ایرانی، دانشگاه علوم پزشکی تهران، تهران، ایران

اطلاعات مقاله چكيده

گلواژگان: شیپوری آنتیاکسیدان اثر سمیت سلولی جنیستئین ایزوفلاونویید

مقدمه: گیاه . Arum rupicola Boiss (خانواده گل شیپوری) گونهای شیپوری است که توسط مردم بومی مناطق مقدمه: گیاه . Arum rupicola Boiss (خانواده گل شیپوری) گونهای شیپوری است که توسط مردم بومی مناطق جنوبی ایران در آشی به نام "آش کارده" استفاده می شود. تاکنون ترکیبات فنلی متعددی از گونههای شیپوری شناسایی شده است. هدف از این مطالعه ارزیابی اثر آنتیاکسیدانی و تعیین محتوای تام فنلی و بررسی فعالیت سمیت سلولی عصاره برگ A. rupicola می باشد. روش بررسی: فعالیت آنتیاکسیدانی عصاره تام متانولی و فراکسیونهای هگزانی، کلروفرمی، اتیل استاتی و باقیمانده آبی به روش FRAP و PPH ارزیابی شد. محتوای تام فنلی با استفاده از روش فولین سیوکالتئو مورد سنجش قرار گرفت. اثر سمیت سلولی عصاره بر ردههای سلولی سرطان پستان انسان فیتوشیمیایی باقیمانده آبی با استفاده از کروماتوگرافی ستونی انجام شد. نتایج: بر اساس نتایج مشاهده شده، باقیمانده آبی کمترین مقدار ۱۵۶۵ برابر با ۱۸۶۷ میکروگرم در میلیلیتر را در تست PPPH نشان داد و عصاره متانولی تام فنلی برابر با بیکرومول گالیک اسید در گرم عصاره را نشان داد. فراکسیون هگزانی نیز بیشترین اثر سیتوتوکسیک را بر رده سلولی سرطان پستان پر استان یتر ایستان داد و محتوای تام فنلی برابر با ۱۳۵۸ میکرومول گالیک اسید در گرم عصاره را نشان داد. فراکسیون هگزانی نیز بیشترین اثر سیتوتوکسیک را بر رده سلولی سرطان پستان ۲-IC۶ MCF برابر با ۱۸۷۹ میکروگرم در میلیلیتر) نشان داد. بررسی فیتوشیمیایی باقیمانده آبی منجر به جداسازی و شناسایی سه ایزوفلاونوئیدی در این گیاه، میتوان قابلیت مصرف آن را به عنوان مکمل نیتراستروژنی در مطالعات آینده در نظر گرفت.

مخففها: FRAP، قدرت احیاکنندگی آهن پلاسما؛ DPPH، ۲،۲- دی فنیل ۱- پیکریل هیدرازیل؛ MTT، ۳-(۵،۴- دی متیل تیازول-۲- ایل)-۵،۲- دی فنیل تترازولیوم بروماید

تاریخ دریافت: ۲۰ اسفند ۱۳۹۹؛ تاریخ دریافت اصلاحات: ۲۸ خرداد ۱۴۰۰؛ تاریخ پذیرش: ۱۲ تیر ۱۴۰۰

doi: 10.52547/jmp.20.79.14

© 2020. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (https://creativecommons.org/licenses/by-nc/4.0/)

^{*} نویسنده مسؤول: n_sadati@tums.ac.ir