

Journal of Medicinal Plants

Journal homepage: www.jmp.ir



Review Article

A comprehensive review of the pharmacological properties and medicinal uses of *Allium jesdianum* Boiss. & Buhse

Zahra Shamsipour Nehzomi, Kobra Shirani*

Department of Toxicology, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

ARTICLE INFO	ABSTRACT
Keywords: Allium jesdianum Biological activity Phytochemical compounds Pharmacological effects Medicinal plants	Background: <i>Allium jesdianum</i> Boiss. & Buhse is a native herb from the <i>Amaryllidaceae</i> family found in Middle East regions. Due to its numerous medicinal and biological properties, it is used for the treatment of many diseases, including colds, infections, kidney stones, etc. Objective : In this review, we investigated the medicinal uses, phytochemical properties, and pharmacological properties of <i>A. jesdianum</i> . Methods: The medicinal uses and phytochemical and pharmacological properties of <i>A. jesdianum</i> , were searched databases such as Google Scholar, PubMed, Scopus, and the Web of Science using keywords such as <i>A. jesdianum</i> , biological activity, phytochemical compounds, pharmacological effects, medicinal plants. Results : Studies have shown that <i>A. jesdianum</i> contains various chemical compounds such as flavonoids, polyphenols, and Sulphur compounds. Moreover, the antioxidant, anticancer, analgesic, antidepressant, antifungal, antibacterial, kidney protective, liver protective, immune system enhancing, radioprotective, and hematopoietic effects of this plant have been demonstrated. Furthermore, <i>A. jesdianum</i> can be used as a natural preservative to improve the oxidative stability of foods rich in unsaturated fatty acids, control the growth of microorganisms, and enhance the immunity of foods. Conclusion : Based on the results obtained, <i>A. jesdianum</i> can be useful as a protective agent against chemical toxicity and overuse of drugs. However, further studies are needed to investigate the clinical effects of this extract on human health.

*Corresponding author: <u>k.shirani@modares.ac.ir</u>

doi: 10.61186/jmp.23.90.1

Received 29 May 2024; Received in revised form 29 September 2024; Accepted 1 October 2024

Abbreviations: ALT, Alanine Aminotransferase; AST, Aspartate Aminotransferase; Bax, Bcl-2-associated X protein; Bcl-2, B-cell lymphoma 2; B-CPAP, thyroid cancer cell line; CCL4, Carbon Tetrachloride; Caspase-3, cysteine-aspartic acid protease 3; CAT, catalase, COX-1, Cyclooxygenase-1; COX-2, Cyclooxygenase-2; CTGF, Connective Tissue Growth Factor; ESWL, Extracorporeal Shock Wave Lithotripsy; GABA, Gamma-aminobutyric acid; GBM, glioblastoma; HT-29, colon cancer cell line; IFN-γ, Interferon-gamma; IL-1β, Interleukin-1 Beta; IL-4, Interleukin-4; LDH lactate dehydrogenase; MDA Malondialdehyde; mg/kg, Milligrams per kilogram; MLKL, mixed lineage kinase domain-like protein; MMP-2, matrix metalloproteinase 2; MMP-9, matrix metalloproteinase 9; μ g/ml, micrograms per milliliter; NO, Nitric oxide; NSAID, Non-steroidal anti-inflammatory drug; p53, tumor protein p53; RAGE, Receptor For Advanced Glycation End Products; RIPK1, receptor-interacting serine/threonine-protein kinase 1; SOD, Superoxide Dismutase; Thr.C1.PI33, thyroid cancer cell line; TNF-α, Tumor Necrosis Factor Alpha

^{© 2023.} Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<u>https://creativecommons.org/licenses/by-nc/4.0/</u>)

1. Introduction

In recent years, the focus on plant foods and medicinal plants due to their potential impact on health has increased. There has been a global rise in the use of herbal medicines as supplements or alternatives to chemical medicines, primarily because of their lower cost and minimal side effects. Extensive studies have been conducted to investigate the bioactive compounds present in plants and their effects on improving health [1, 2].

The genus Allium is a diverse group of Amaryllidaceae family that comprises over 850 species, including 121 species found in Iran. This genus includes important horticultural products such as garlic, shallots, onions, leeks, and other species with unknown medicinal, spice, vegetable, or ornamental properties. Due to their beneficial effects on human health, numerous scientific research studies have examined the chemical and biological properties of Allium genus. The medical use of Allium genus dates back to the Ebers Codex book in Egypt around 1500 BC. Throughout history, notable figures like Hippocrates, Herodotus, Pasteur, and Albert Schweitzer have mentioned the use of Allium genus for medicinal purposes. Initially, they were likely used as protective agents against febrile

Z. Shamsipour Nehzomi & K. Shirani

diseases, including plague and other infectious diseases. In Europe during the Middle Ages, monks cultivated garlic to prevent and treat infectious diseases. In the 19th century, Pasteur described the antibacterial properties of garlic extract. During the First World War, garlic extract was used topically to prevent gangrene, and in the Second World War, it was referred to as Russian Penicillin. This effect is primarily attributed to two main categories of compounds: sulfur compounds and saponins. All types of Allium genus contain more than 200 components, including vitamins, sulfur-containing compounds, amino acids, proteins, and lipids, rare elements such as selenium, flavonoids, and antioxidants [3, 4].

One of the remarkable species within the *Allium* genus is *Allium jesdianum* Boiss. & Buhse, locally named Bowser, Sarpa, or Bon-Sorkh in different cities of Iran due to the red color of its onion. *A. jesdianum* found in Middle East regions, growing wild at altitudes of 1800-2600 meters in the western region of Iran. It is an herbaceous, perennial plant that grows from a bulbous rootstock. It produces a cluster of grass-like leaves, and a flowering scape that reaches a height of around 75-90cm (Figure 1).



Fig. 1. Allium jesdianum

Z. Shamsipour Nehzomi & K. Shirani

Local people have a long-standing tradition of gathering the *A. jesdianum* plant during the early spring and selling it in markets. This plant holds culinary value as it is commonly cooked and enjoyed alongside rice or used as an ingredient in soups. However, the distressing reality is that due to rampant illegal harvesting practices in recent years, the *A. jesdianum* plant has been designated as an endangered species.

Furthermore, the aerial parts of A. jesdianum have been deeply rooted in traditional medicine practices. They have been employed for centuries to alleviate various ailments such as abdominal pain, rheumatism, vomiting, kidney stones, and colds. The plant's medicinal properties have made it a valuable resource within local communities, where it has been relied upon as a natural remedy [5-7]. In light of the distinctive characteristics exhibited by this indigenous plant, our research endeavors led us to conduct a comprehensive review article aimed at exploring pharmacological effects and potential its applications. By shedding light on the scientific evidence surrounding this native plant and its different extracts, our review article seeks to contribute to the existing knowledge base. This work provides a solid foundation for further scientific exploration and facilitates the investigation of its potential for various therapeutic interventions.

2. Method

The scientific databases including Scopus, Web of Science, Google Scholar, and PubMed were searched to access all relevant books and papers in English and Persian until 2024. The keywords used in the search were A. jesdianum, biological activity, phytochemical compounds, pharmacological effects, and protective effects.

3. Results

3.1. Phytochemical compounds

Several studies on the essential oil and extract composition of A. jesdianum have highlighted notable variability in chemical constituents, influenced by factors such as ecotype, growth stage, and environmental conditions. For instance, analyses utilizing GC and GC/MS technologies revealed the presence of a range of key compounds. Key components identified in the essential oil include dimethyl trisulfide, hexadecanoic acid, and various terpenes. Specifically, one study found dimethyl trisulfide to comprise 22.34 % of the oil, while hexadecanoic acid accounted for 19.03 %. Other significant compounds included phytol, caryophyllene oxide, and curzerene. The essential oil features significant phytochemicals such as Trisulfide dimethyl and Hexadecanoic acid, both recognized for their bioactive properties. Other notable compounds include Disulfide methyl-1-(methylthio) ethyl and Beta-Ionone, which contribute to the oil's antioxidant, antimicrobial, and anti-inflammatory effects. The oil demonstrates strong antioxidant properties, characterized by a phenol content of 85 ± 3.2 mg GA/100 g and impressive free radical scavenging ability. These findings underscore therapeutic potential the of A. *jesdianum* essential oil, particularly in mitigating oxidative stress [8, 9].

In samples collected from different regions, such as Kohgiluyeh and Boyer-Ahmad province, hexadecanoic acid was frequently noted as the most abundant component. Studies analyzing the essential oil of *A. jesdianum* revealed that sulfide and sesquiterpene compounds, including allicin, diallyl disulfide, diallyl trisulfide, Curzerene, and Curzerenone, are key components of the oil.

Research on dried flower extracts has also revealed a diverse profile of monoterpenes,

Z. Shamsipour Nehzomi & K. Shirani

sesquiterpenes, and non-terpene compounds, with analyses identifying numerous constituents such as thymol, linalool, and nonanal among the major components. Additionally, comparative studies between wild and cultivated ecotypes demonstrated that cultivated plants generally yielded higher quantities and qualities of essential oils and other metabolites, suggesting that organic cultivation practices could enhance the chemical composition and nutrient content of the plant [10].

Phytochemical analyses of the fresh bulbs of *A. jesdianum* identified multiple steroidal glycosides, further contributing to the plant's bioactive potential. These compounds include newly identified compounds such as jesidin,

jesdoside, and jesdioside [11]. Additionally, computational methods have been applied to assess the corrosion resistance of certain phytochemicals found in *A. jesdianum* flowers. Notably, α -linolenic acid, palmitic acid, and 1-ecosanol displayed effective corrosion-resistant properties for metals such as iron, aluminum, and copper [12]. Furthermore, *A. jesdianum* contains phenols and flavonoids among its chemical compounds [13, 14].

3.2. Pharmacological effects

A list of pharmacological effects of *A*. *jesdianum* extract reported in the literature has been mentioned in Table 1.

Concentratio				
Anticancer effects	Model	Concentratio n (μg/ml)	Result	Ref
Methanolic extract	HeLa, K562	>2 (µg/ml)	Inhibitory effects on cell growth	[15]
Hydroalcoholic extract	GBM	250 (µg/ml)	Decreased cell survival, hindered cell migration, reduced cell invasion and adhesion	[16]
Hydroalcoholic extract	HT-29	25, 50, and 100 (μg/mL)	Decreased number of colon cancer cells, increased MDA levels, stimulated ROS production, reduced antioxidants, initiated necrosis, upregulated RIPK1, RIPK3, and MLKL	[17]
Hydroalcoholic extract	B-CPAP, Thr.C1.PI33	200 (g/ml)	Decreased cell growth, increased cytotoxicity, release of LDH, Induction of apoptotic cell death, downregulation of Bcl-2, upregulation of Bax, p53, and Caspase-3	[18]
		Renal prot	tective effects	
Hydroalcoholic extract	Mice	250 and 500 (mg/kg)	Reduced cell destruction, mitigated nephrotoxicity, exhibited antioxidant activity	[24]
Aqueous extract	Wistar rats	250 and 500 (mg/kg)	Reduced kidney stone formation, decreased concentration of urea, creatinine, and uric acid in blood	[6]
Hydroalcoholic extract	Rat	250 and 500 (mg/kg)	Reduced symptoms of diabetic nephropathy, reduced expression of CTGF and RAGE genes, improved oxidative stress conditions	[21]
Hydroalcoholic Extract	Human	250 (mg/kg) Twice Per Day	Improved symptoms, increased rate of stone passage, enhanced kidney function, reduced pain and other symptoms associated with kidney stones	[25]

Table 1. Pharmacological effects of A. jesdianum extract reported in the literature

Journal of Medicinal Plants

Z. Shamsipour Nehzomi & K. Shirani

Fable 1	I. Pharmacological	effects of A. jesdianun	<i>i</i> extract reported in the	literature (Continued)
----------------	---------------------------	-------------------------	----------------------------------	------------------------

Radio protective effects	Model	Concentratio n (µg/ml)	Result	Ref
Hydroalcoholic Extract	NMRI mice	500, 1,000, and 2,000 (mg/kg)	Reduced cell destruction, mitigated nephrotoxicity, exhibited antioxidant activity	[23]
Hepatoprotective eff	ects		*	
Hydroalcoholic Extract	NMRI mice	50 and 100 (mg/kg)	Improvement in the hepatic function and histopathological changes	[25]
Hydroalcoholic Extract	Rat	200 (mg/kg)	Increased antioxidant activity, Improved liver biochemical parameters, Reduced blood lipid levels	[26]
Hydroalcoholic Extract	Rat	200 and 400 (mg/kg)	Increased antioxidant enzymes activity, Reduced peroxidative enzyme levels, Decreased in inflammatory markers and liver fat deposition	[27]
Hydroalcoholic Extract	Rat	250 and 500 (mg/kg)	Improved liver markers, Decreased levels of inflammatory cytokines, Reduced oxidative damage, and mitigated apoptotic	[28]
Antioxidant activity				
Ethanolic Extract	DPPH assay	8 to 500 (µg/ml)	Scavenged free radical	[31]
Ethanolic Extract	DPPH assay	0.1, 0.5, 1, and (2 mg/ml)	Scavenged free radical	[32]
Methanol Extract Aqueous Extract	DPPH and ABTS radical scavenging activity	50,100, 200 mg/ml	Exhibited higher antioxidant activity	[35]
Neuroprotective effect	ets			
Aqueous Extract	Rat	50, 100, 200, and 400 (mg/kg)	Anxiolytic and anti-depressant activity	[30]
Hydroalcoholic Extract	Rat	100, 200, and 400 t (mg/kg/day)	Downregulation of Bax gene expression and upregulation of miR- 330, miR-132, and Bcl-2 gene expression Reversed mitochondrial toxicity parameters	[37]
Analgesic Effects		0.5		
Hydroalcoholic Extract	- Human fresh blood	0.5 to 6 mg/ml	Reduced the activity of the COX-1 enhanced the activity of the COX-2	[39]
Essential oil		0 to 100 (mg/ml)	Enhanced the activity of COX-1 but had no effect on COX-2	
Hydroalcoholic Extract	Rat		Increased Tail Flick and Hot Plate latencies	[40]
Blood Effects				
Hydroalcoholic		0.5 to 6		
Extract	Human fresh	(mg/ml)	Inhibit platelet aggregation	[39]
Essential oil	Essential oil blood			r 1
Methanolic Extract	Human fresh blood	?	Decreased in platelet aggregation	[41]

Z. Shamsipour Nehzomi & K. Shirani

Immune protective effects	Model	Concentratio n (µg/ml)	Result	Ref
Ethanolic Extract	Mice	50,100, 200 mg/kg	Increased number of lymphocytes and elevated levels of antibodies	[45]
Aqueous Extract	Rat (bone stem cells)	500 and 1000 (µg /ml)	Reduced IFN-γ and IL-17 Increased IL-4	[46]
Aqueous Extract	Mouse peritoneal macrophages	(0.5, 1, 5 and 10 mg/mL)	Increased NO production at non-toxic concentrations.	[47]
Non-medical Use				
-	Mild Steel in Acidic Media	200 ppm	Enhanced the corrosion resistance	[52]
Palmitic acid a-Linolenic acid 1-Ecosanol	Electron Density Functional Method	-	Prevented corrosion in metals	[12]
Aqueous Extract	As the nano- catalyst for electrocatalytic oxidation	200 mg/liter	Exhibited exceptional electrochemical activity	[53]
Alcoholic Extract	Shelf life of white fish fillets	(0.5%, 1%, 1.5%)	improved the shelf life of white fish fillets dose dependently	[54]
Antibacterial activity	y			
Aqueous Extract	Pathogenic bacteria	230 (mg/ml)	Inhibition of bacterial growth	[7]
Methanol Extract	antibiotics	130 (mg/ml)		
Bacterial extract derived from the endophyte of the A. <i>jesdianum</i>	Gram-positive bacteria	100 mg/ml	Decrease in the growth of pathogenic bacteria and fungi	[34]
Methanol Extract Aqueous Extract	P. aeruginosa, E. coli, E. aerogenes, L. innocua, and S. pyogenes	50,100, 200 mg/ml	Treatment for antibiotic-resistant bacteria	[35]
Antifungal activity				
Methanol Extract Ethanolic Extract	Trichophyton mentagrophytes	50 and 100 mg/ml	Decrease keratinase activity	[50]
Methanol Extract Ethanolic Extract	Human vaginal Candida glabrata isolates		Antifungal efficacy	[49]
Hydroalcoholic Extract	Candida albicans	(0.5, 1, 5 and 10 mg/mL)	Anticandidal activity.	[42]

Table 1. Pharmacological effects of A. jesdianum extract reported in the literature (Continued)

3.2.1. Anticancer effects

Zarabi et al., (2017) found that concentrations higher than 2 μ g/ml of the methanolic extract derived from the *A. jesdianum* plant exhibited significant inhibitory effects on cell growth in both HeLa and K562 cell lines. The presence of chemical compounds such as phenols, flavonoids, and anthocyanins in the plant strongly indicates their potential contribution to its anticancer activity [15]. The findings of Rashidi et al., (2022) study showed that the *A*. *jesdianum* extract (250 μ g/ml) cause a

Z. Shamsipour Nehzomi & K. Shirani

concentration- and time-dependent decrease in cell survival by hindering cell migration, reducing cell invasion and adhesion, and downregulating the expression of matrix metalloproteinase 2 and 9 (MMP-2 & 9) in glioblastoma (GBM) cells [16]. In another study the hydroalcoholic extract of A. jesdianum exposure led to a reduction in the number of colon cancer cells (HT-29) by elevating the levels of MDA, stimulating the production of reactive oxygen species (ROS), decreasing antioxidants, initiating necrosis, and upregulating receptor-interacting serine/threonine-protein kinase 1 (RIPK1), RIPK3, and mixed lineage kinase domain-like protein (MLKL) [17]. Moreover, exposure to the hydroalcoholic extract of A. jesdianum (200 mg/ml) resulted in decreased cell growth, increased cytotoxicity, the release of lactate dehydrogenase (LDH), and induction of apoptotic cell death in thyroid cancer cells (B-CPAP and Thr.C1.PI33). These effects were accompanied by downregulation of B-cell lymphoma 2 (Bcl-2) expression and upregulation of Bcl-2-associated X protein (Bax), tumor protein p53 (p53), and cysteine-aspartic acid protease 3 (Caspase-3) expression [18]. Further in our recent publication, the hydroalcoholic extract of A. jesdianum has shown significant cytotoxic effects on human melanoma cell lines [19]. Based on these findings, it may be concluded that A. jesdianum possesses anticancer properties, stimulates apoptosis, and may serve as a natural source for further clinical experimental and study for the development of anticancer drugs.

3.2.2. Renal protective effects

In a study conducted by Kalantari et al. (2018), the hydroalcoholic extract of *A. jesdianum* (250 and 500 mg/kg) was investigated for its protective effects against carbon

tetrachloride (CCl4) -induced nephrotoxicity in mice. The results demonstrated that A. jesdianum destruction, mitigated extract reduced cell nephrotoxicity, and exhibited antioxidant activity [20]. Another study examined the effect of the aqueous extract of A. jesdianum (250 and 500 mg/kg) on kidney stones induced by ethylene glycol in male Wistar rats. The findings revealed that the consumption of A. jesdianum extract at a dose of 500 mg/kg by rats significantly reduced kidney stone formation and decreased blood level of urea, creatinine, and uric acid [6].

In another study, the hydroalcoholic extract of *A. jesdianum* at doses of 250 and 500 mg/kg reduced diabetic nephropathy symptoms and lowered the expression of connective tissue growth factor (CTGF) and receptor for advanced glycation end products (RAGE) genes. The extract also improved oxidative stress by reducing MDA levels and increasing superoxide dismutase (SOD) activity [21]. The study demonstrated that the combination of tamsulosin and *A. jesdianum* extract improved symptoms, increased stone passage rate, enhanced kidney function, and reduced pain and other related symptoms compared to tamsulosin alone [22].

In general, the research results indicate that *A. jesdianum* is an effective herbal compound with increased antioxidant activity, which can serve as an adjunctive treatment in reducing symptoms associated with kidney disorders such as diabetic nephropathy, kidney stones, and risks caused by toxin exposure.

3.2.3. Protective effects against radiation

The study conducted by RoshanKhah et al. (2021) indicates that *A. jesdianum* extract (250 and 500 mg) has protective effects against gamma radiation-induced damage in the pancreas. The extract was found to decrease the expression of apoptotic proteins Caspase-3 and 9, increase the antioxidant activity of SOD and

catalase (CAT), and reduce the levels of inflammatory markers tumor necrosis factor alpha (TNF- α) and interleukin-1 beta (IL-1 β) in the pancreas of mice. These findings suggest that *A. jesdianum* extract has potential as a radioprotective agent [23].

3.2.4. Hepatoprotective effects

The liver is a vital metabolic organ involved in drug and chemical metabolism and is susceptible to damage from toxins. Liver damage is associated with systemic oxidative stress, leading to cell necrosis, fibrosis, tissue lipid peroxidation, and reduced glutathione levels. As reliable hepatoprotective drugs are lacking in modern medicine, compounds derived from plants represent a significant option [24]. Research suggests A. jesdianum may have hepatoprotective properties. According to the study by Sohrabinjad et al. (2019), mice that received alcoholic extract of A. jesdianum (50 and 100 mg/kg) experienced improvements in hepatotoxicity induced by acetaminophen [25]. Another study by Kalantari et al. in 2018 showed that the hydroalcoholic extract of A. jesdianum (200 mg/kg) reduced bromobenzene-induced hepatotoxicity in rats through increased antioxidant activity and decreased oxidative stress in the liver. Additionally, the extract significantly improved liver biochemical parameters such as alanine aminotransferase (ALT), aspartate aminotransferase (AST), and reduced blood lipid levels [26]. In another study same authors, consumption by the of A. jesdianum extract (200 and 400 mg/kg) hepatotoxicity induced against by CCl4 increased the activity of antioxidant enzymes, peroxidative enzyme levels. reduced inflammatory markers and liver fat deposition [27]. Furthermore, the use of A. jesdianum extract at both doses of 250 and 500 mg/kg

Z. Shamsipour Nehzomi & K. Shirani

improved liver markers, decreased levels of inflammatory cytokines, reduced oxidative damage, and mitigated apoptotic changes in the livers of rats exposed to mercury chloride [28]. Based on the obtained results, it can be concluded that Α. jesdianum may serve as а hepatoprotective substance against chemical toxicity and drug overdose by reducing inflammatory cytokines, oxidative damage, and apoptotic changes. However, further studies are required to investigate the clinical effects of this extract on human liver health.

3.2.5. Antioxidant activity

Oxidative stress arises when there is an imbalance between generation the and accumulation of ROS within cells and tissues and the body's ability to eliminate them. ROS are produced as a natural byproduct of oxygen metabolism, but various environmental stressors, such as UV radiation, ionizing radiation, pollutants, heavy metals, and anticancer drugs, can greatly increase ROS production, resulting in damage to cells and tissues. In recent years, there has been increasing interest in investigating the and potential of medicinal plants their compounds, particularly flavonoids and polyphenols, as a means of combating oxidative stress [29, 30]. In Ghasemi et al., (2019), various extracts of onion and leaves from different populations of A. jesdianum were found to exhibit significant antioxidant activity [31]. Furthermore, Ahmadzadeh et al.'s research in 2014 showed that the ethanolic extract of A. *jesdianum* leaves exhibited significant antioxidant activity all tested across concentrations (0.1, 0.5, 1, and 2 mg/ml) in a dose-dependent manner [32-34]. In 2021, Hojjati et al. conducted a study to investigate the antioxidant properties of A. jesdianum extracts using two extraction methods: aqueous and

methanolic. The results indicated that the methanolic extract (200 mg/l) exhibited higher antioxidant activity compared to the aqueous extract. As a result, both methanolic and aqueous extracts of *A. jesdianum* could serve as natural preservatives, enhancing the oxidative stability of food products rich in unsaturated fatty acids and improving the safety of food products [35].

3.2.6. Neuroprotective effects

Plants in the *Allium* genus, such as garlic and onions, contain bioactive compounds like allicin, known for their protective effects on the nervous system. Allicin offers antioxidant and antiinflammatory properties, helping to combat oxidative stress and reduce inflammation. This has led to growing interest in *Allium* genus, particularly *A. jesdianum* as potential natural therapies for supporting neurological health and preventing neurodegenerative diseases [36].

The results of Mousavi et al., (2013) study showed that administering the aqueous extract of the A. jesdianum plant (50, 100, 200, and 400 mg/kg) significantly reduced anxiety and depression behaviors in rats. Furthermore, changes in the expression of genes related to anxiety and depression were observed in the rats' brains after the extract was administered. These findings suggest that the A. jesdianum extract holds promise as a natural agent with the potential to alleviate anxiety and depression in humans by affecting the Gamma-aminobutyric acid (GABA) receptor system and serotonin system in the brain [33]. Kamranfar et al., (2023) evaluated the protective effects of a hydroalcoholic extract of A. jesdianum in a mouse model of Alzheimer's disease. Extracted from A. jesdianum markedly reduced cognitive dysfunction and mitochondrial toxicity induced by intracerebroventricular streptozotocin (icv-STZ) administration. After STZ injection, there

Z. Shamsipour Nehzomi & K. Shirani

was upregulation of Bax gene expression and downregulation of miR-330, miR-132, and Bcl-2 gene expression. Treatment with *A. jesdianum* extract led to a reversal of the expression of these microRNAs and genes, suggesting its potential in ameliorating Alzheimer and reducing neuronal apoptosis. This study demonstrates the neuroprotective ability of *A. jesdianum* against STZ-induced oxidative stress and cognitive impairment in mice, highlighting its therapeutic potential in the management of Alzheimer [37].

3.2.7. Analgesic Effects

Pain and sensitivity often result from inflammation caused by tissue injuries and infections, which stimulate nerves and release substances like prostaglandins that promote pain. Due to the side effects of long-term synthetic drugs for pain relief, many studies have investigated the analgesic and anti-inflammatory properties of plant extracts as natural alternatives to alleviate pain without the adverse effects of synthetic medications [38]. In Khaksarian et al., (2016) study, the effects of A. jesdianum extract doses ranging from 0.5 to 6 mg/ml and essential oil doses ranging from 0 to 100 mg/ml extracted investigated on the activity were of Cyclooxygenase-1 (COX-1) and Cyclooxygenase-2 (COX-2) enzymes. The study revealed that the A. jesdianum extract reduced the activity of the COX-1 enzyme while enhancing the activity of the COX-2 enzyme. On the other hand, the essential oil from the plant enhanced the activity of COX-1 but had no effect COX-2 [39]. Another on study using hydroalcoholic extract from A. jesdianum leaves demonstrated significant analgesic properties, indicating that these effects are mediated by various brain regions and the opioid system [40]. Overall, the results suggest that A. jesdianum can serve as a medicinal resource with anti-

Z. Shamsipour Nehzomi & K. Shirani

inflammatory and pain-relieving properties. Phenolic compounds, alkaloids, and saccharides present in the extract have been proposed as potentially effective substances contributing to these analgesic properties. However, further studies are needed to fully understand the precise analgesic mechanisms of *A. jesdianum*.

3.2.8. Blood Effects

In 2017, Khaksarian and his colleagues showed that, in addition to its anti-inflammatory and pain-relieving properties, A. jesdianum is able to inhibit platelet aggregation in a dosedependent fashion. This effect is also observed in drugs such as aspirin, which is a useful nonsteroidal anti-inflammatory drug (NSAID) in cardiovascular diseases [39]. Additionally, a 2015 study by Lorigooini et al. evaluated the anti-platelet aggregation effect of different species from the Allium genus. The extract of A. jesdianum, administered at a dose of 200 micrograms, caused a significant decrease in platelet aggregation [41]. Therefore, this plant, as an herbal composition, can be used as an alternative treatment for diseases associated with blood clot formation. Immune protective effects.

3.2.9. Immune System Effects

The immune system protects the body from pathogens and cancers, and even small declines in its function can increase disease risk. While there are various methods to boost immunity, some may be costly with limited effects. In contrast, many herbal medicines are costeffective, have good immunomodulatory properties, and cause fewer side effects, making them suitable as adjuncts or alternatives to conventional treatments [42]. Allium plants, which are found extensively around the globe, have the potential to enhance immune responses to different diseases through their ability to modulate immunity. This has led to significant interest in exploring the potential of A *A. jesdianum* as a natural immunomodulatory [43, 44].

The study by Rahimi and Madani in 2015 demonstrated the potential beneficial effects of the ethanolic extract of A. jesdianum on the humoral response of mice infected with bacteria but also highlighted the importance of finding the appropriate dosage to avoid any adverse effects [45]. In another study, Radjabian et al. (2021) showed that the aqueous extract of A. jesdianum acts as an immunomodulatory agent by positively influencing the response of spinal cord stem cells in terms of the production of Interleukin-4 (IL-4) cytokine and reducing the production of Interferon-gamma (IFN-y) and [46]. Interleukin-17 (IL-17) cytokines Furthermore, the aqueous extract of A. jesdianum has shown the ability to stimulate macrophage cells to increase Nitric oxide (NO) production at non-toxic concentrations [47]. Our recent study seems that A. jesdianum improves various factors of the immune system. Its effects include increasing the proliferation of spleen lymphocytes, delayed proliferative response, and antibody production. In general, A. jesdianum was able to strengthen various immune responses, including humoral immunity and cellular immunity [13]. Consequently, the use of A. jesdianum, at appropriate doses, may be considered a natural and harmless method to modulate the immune response. According to our latest unpublished research A. jesdianum exerts protective effects on the immune system of cyclophosphamide -treated animals by boosting both cellular and humoral immunity, with no observed hepatoxicity [48].

3.2.10. Antibacterial Effects

The antibacterial effects of both aqueous and methanolic extracts of *A. jesdianum* were

evaluated against several pathogenic bacteria resistant to antibiotics, such as Staphylococcus aureus, Pseudomonas aeruginosa, Escherichia coli, and Klebsiella pneumoniae. The results revealed that the methanolic extract of A. jesdianum exhibited better inhibition of bacterial growth compared to the aqueous extract. Notably, at a concentration of 500 µg/ml, the methanolic extract completely inhibited the growth of the tested bacteria [7]. Esmaeili et al. (2021) demonstrated the potential of the bacterial extract from A. jesdianum endophytes as a novel source for inhibiting the growth of gram-positive bacteria. The results indicated that using a dose of 100 mg/ml extract led to a significant decrease in the growth of pathogenic bacteria and fungi [34]. Based on Hojjati et al. (2021) findings, the methanolic extract of A. jesdianum demonstrated greater antimicrobial activity compared to the aqueous extract. Therefore, the A. jesdianum may potentially serve as an alternative treatment for antibiotic-resistant bacteria [35].

3.2.11. Antifungal Effects:

Recent studies have demonstrated the significant antifungal activity of aqueous extract, hydroalcoholic extract, and A. jesdianum essential oil against Candida Albicans in conditions laboratory [34]. For instance, Shahrokh et al. 2017 investigated the antifungal effects of the water-alcoholic extract of A. jesdianum at doses of 250, 500, 1000, and 2000 ug/ml on fluconazole-resistant and sensitive Candida Glabrata isolates obtained from the human female genital tract. The results indicated that the A. jesdianum extract exhibited substantial antifungal effect on a both fluconazole-sensitive and resistant isolates [49]. In another study, it was found that the alcoholic extract of A. jesdianum at a dose of 200 µg/ml exhibited the greatest effect in reducing the keratinase activity of the fungus [50]. Similarly, Khezezlo et al.'s study in 2017 revealed that the ethanolic extract of the *A. jesdianum* plant (640 μ g/ml) displayed strong antifungal effects against *T. Mentagrophytes* isolates [51]. These findings suggest that the hydroalcoholic extract of *A. jesdianum* possesses potent antifungal properties may potentially serve as a rich source of antifungal compounds.

3.4. Non-medical Use

In a 2021 study conducted by Kahkesh, the potential of A. jesdianum extract as a green corrosion inhibitor for mild steel in hydrochloric acid solutions was highlighted. The results of the study revealed that the plant extract not only inhibits corrosion but also has a positive impact on the surface integrity of the metal. Further research and development in this area could lead to the development of more environmentally friendly corrosion protection methods for metal materials [52]. In another study, researchers investigated the effect of phytochemical components of A. jesdianum flower in preventing corrosion of three metals - iron, aluminum, and copper - using the electron density functional method (DFT). The findings showed that utilizing these phytochemical components could be a cost-effective and straightforward method to prevent corrosion in these metals. This research provides valuable scientific insights into the potential use of A. jesdianum extract as a corrosion prevention technique [12]. In Sheikh study, the water Mohseni's extract of A. jesdianum (200 mg/liter) was utilized for the synthesis of a Ni/Montmorillonite Nano catalyst. This method yielded a product with excellent electrochemical properties, suitable for use in the electrochemical oxidation reaction of methane. The results demonstrated that the prepared nanocomposite containing nickel catalyst

exhibited exceptional electrochemical activity in electrochemical oxidation reaction of the methane [53]. In another study, the effect of the alcoholic extract of A. jesdianum plant (0.5 %, 1 %, and 1.5 %) on the shelf life of white fish fillets was investigated [55]. The results revealed that the use of A. jesdianum extract at a dose of 1 % significantly improved the shelf life of white fish fillets. However, doses higher than 1 % had a negative effect on the shelf life of fish fillets [49]. Furthermore, Akrami et al. (2021) indicated that bio nanocomposite films containing nanoliposomes of A. jesdianum extract could be used as food packaging with effective antibacterial properties [54].

Conclusion

Based on the findings of this review, *A. jesdianum* can be regarded as an herbal compound that possesses anticancer, renal protective, radioprotective, hepatoprotective, neuroprotective, analgesic, anti-inflammatory, antioxidant, immune protective, antifungal, and antibacterial effects. This suggests that the extract may serve as a valuable protective agent against drug and chemical toxicity, potentially

References

1. Shirani K, Vahdati Hassani F, Razavi Azar-Khiavi K, Samie Moghaddam Z and Karimi GR. Determination of methanol in Iranian herbal distillates. *J. Complement. Integr. Med.* 2016; 13(2): 123-7. doi: 10.1515/jcim-2015-0041.

2. Yousefsani BS, Bahrami B, Qobadi A, Aghazadeh E, Shirani K, Pourahmad J. The selective cytotoxicity of the hydroalcoholic extract of *Santalum album* Linn wood on A375 and SK-MEL-3 human malignant Melanoma cells. *Iran. J. Pharm. Res.* 2022, 18(2): 128-137. doi: 10.22037/ijps.v18.42139.

3. Štajner D, Milić N, Čanadanović-Brunet J, Kapor A, Štajner M and Popović BM. Exploring

offering significant benefits for human health. However, additional research is essential to understand its efficacy and safety fully. Such studies could explore the extract's mechanisms of action, optimal dosages, and long-term effects, ultimately paving the way for its clinical application and integration into therapeutic strategies.

Author contributions

Research conception and design: KSH and ZSH. Data collection: KSH and ZSH. Review and editing: KSH. All authors contributed to the article and approved the submitted version.

Conflicts of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Acknowledgment

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Allium species as a source of potential medicinal agents. *Phytother. Res.* 2006; 20(7): 581-584. doi: 10.1002/ptr.1917.

4. Lanzotti V, Scala F and Bonanomi G. Compounds from *Allium* species with cytotoxic and antimicrobial activity. *Phytochem. Rev.* 2014; 13(4): 769-791. doi: 10.1007/s11101-014-9366-0.

5. Ramak P and Asri Y. Effect of growth degree days and soil properties on phenology and morphological characters of *Allium jesdianum* Boiss & Buhse in Lorestan province, *Iran. J. Plant. Biol.* 2019; 10(4): 35-52.

Journal of Medicinal Plants

Z. Shamsipour Nehzomi & K. Shirani

Z. Shamsipour Nehzomi & K. Shirani

6. Vahdani R, Mehrabi S, Malekzadeh J, Jannesar R, Sadeghi H and Shafaeifar A. Effect of hydrophilic extract of *Allium jesdianum* on ethylene glycol-induced renal stone in male wistar rats. *Armaghane Danesh*. 2012; 16(6): 566-577.

7. Gholami A, Arabestani MR and Ahmadi M. Evaluation of antibacterial activity of aqueous and methanol extracts of *Allium jesdianum* plant on a number of pathogenic bacteria resistant to antibiotics. *Pajouhan. Sci. J.* 2016; 14(4): 18-26. doi: 10.21859/psj-140418.

8. Amiri H. Chemical composition and antibacterial activity of the essential oil of *Allium jesdianum* Boiss. & Buhse from Iran. *J. Med. Plants*. 2007; 6(Supp. 3): 39-44.

9. Naserabad SS, Zarei S, Rahimi J, Ghafouri Z, Mouludi-Saleh A and Banaee M. Protective effects of *Allium jesdianum* essential oil on rainbow trout (*Oncorhynchus mykiss*) exposed to sub-lethal toxicity of cypermethrin. *Aquat. Toxicol.* 2024; 274: 107051. doi: 10.1016/j.aquatox.2024.107051.

10. Askari Y. Composition of essential oil of *Dorema aucheri* Boiss. and *Allium. jesdianum* Boiss. medicinal plants. *Int. J. Adv. Biol. Biomed. Res.* 2022; 10(1): 72-83. doi: 10.22034/IJABBR.2022.545885.1374.

11. Mimaki Y, Kuroda M, Fukasawa T and Sashida Y. Steroidal glycosides from the bulbs of *Allium jesdianum. J. Nat. Prod.* 1999; 62(1): 194-197. doi: 10.1021/np980346b.

12. Wazzan N. Phytochemical components of *Allium jesdianum* flower as effective corrosion-resistant materials for Fe (1 1 0), Al (1 1 1), and Cu (1 1 1): DFT study. *Arab. J. Chem.* 2023; 16(4): 104625. doi: 10.1016/ j.arabjc. 2023.104625.

13. Keshavarzipoor K, Soodi M and Shirani K. Investigating the effects of hydroalcoholic extract of the *Allium jesdianum* on humoral and cellular indicators of the immune system in a mouse model stimulated with sheep erythrocyte Immunogen compared to Levamisole. *Research in Medicine*. 2024; 47(4): 9-21.

14. Falahi E, Delshadian Z, Ahmadvand H and Jokar SS. Head space volatile constituents and antioxidant properties of five traditional Iranian wild edible plants grown in west of Iran. *AIMS Agri. Food.* 2019; 4(4): 1034-1053. doi: 10.3934/agrfood.2019.4.1034.

15. Dorosti N, Zarabi S, Ahmadi S, Rostami R and Rashidipour M. Anticancer activity evaluation of methanolic extract of *Allium jesdianum* and Nectaroscordeum Coelzi against HeLa and K562 cell lines. *Yafte* 2017; 19(1): 31-41.

16. Rashidi I, Hajalikhani P, Jalili C and Zhaleh M. The cytotoxic and migrastatic potentials of *Allium jesdianum* hydroalcoholic extract on glioblastoma multiforme cell line model. *WCRJ*. 2022; 9: e2151. doi: 10.32113/wcrj_20221_2151.

17. Alidadi H, Shirani M, Samimi A, Salimi A, Ashtari A and Khorsandi L. *Allium jesdianum* extract induces oxidative stress and necroptosis in human colorectal cancer (HT-29) cell line. *BABT*. 2021; 64: e21200491. doi: 10.1590/1678-4324-202120049.

18. Naseri R, Habibi Shams MJ, Gholami MR, Rashidi I and Jalili C. Anti-cancer and apoptosis induction effects of *Allium jesdianum* hydroalcoholic extract on thyroid cancer cell lines (b-cpap and thr. C1-pi 33). *WCRJ*. 2021; 8: e2104. doi: 10.32113/wcrj_20219_2104.

19. Shirani M, Aghazadeh E, Yousefsani B.S and Shirani K. Anticancer activity of *Allium jesdianum* extract: A potential therapeutic approach for Melanoma. *Jundishapur. J Nat. Pharm. Prod.* 2023; 19(1): e139022. doi: 10.5812/jjnpp-139022.

20. Kalantari H, Danesh Pajou M, Kheradmand P, Goodarzian M and Zeidooni L.

Journal of Medicinal Plants

Nephroprotective effect of hydroalcoholic extract *Allium jesdianum* Boiss against carbon tetrachloride induced nephrotoxicity via stress oxidative in mice. *Pharm. Sci.* 2018; 24(2): 89-96. doi: 10.15171/PS.2018.14.

21. Alaee M, Amri J, Karami H, Latifi SA, Salemi Z and Salehi M. *Allium jesdianum* hydro alcoholic extract ameliorates diabetic nephropathy by suppressing connective tissue growth factor (CTGF) and receptor for advanced glycation endproducts (RAGE) gene expression in diabetic rats with streptozotocin. *Horm. Mol. Biol. Clin. Investig.* 2021; 42(2): 167-174. doi: 10.1515/hmbci-2020-0072.

22. Mehrabi A, Mehrabi S, Sadeghi H, Dousti Motlagh AH, Ghadimi Moghadam AK and Ghadimi Moghadam A. Comparison efficacy of combined Tamsulosin and *Allium jesdianum* versus Tamsulosin in treatment of renal colic and passage of urinary stones after Extracorporeal Shock Wave Lithotripsy (ESWL): a randomized clinical trial. *PJMHS*. 2020; 14(2): 1133-1136.

23. Roshankhah S, Zahabi S.S, Gholami M.R, Abdolmaleki А and Salahshoor M.R. Radioprotective effects of Allium jesdianum extract on reduction of Pancreas damages following γ -radiation through down-regulation of Apoptotic genes, antioxidants regulation, and suppression of inflammatory markers. Jundishapur J. Nat. Pharm. Prod. 2021; 16(3): e103997. doi: 10.1016/j.jphotobiol.2017.12.004. 24. Shirani K, Riahi-Zanjani B, Omidkhoda SN, Barangi S and Karimi GR. The hematopoietic potential of methanolic and aqueous extracts of Portulaca oleracea in a phenylhydrazine model of anemia. AJP. 2023; 13(1): 85-96. doi: 10.22038/AJP.2022.20965.

25. Sohrabinezhad Z, Dastan D, Soleimani Asl S and Nili-Ahmadabadi A. *Allium jesdianum* extract improves acetaminophen-induced hepatic failure through inhibition of oxidative/nitrosative

stress. *J. Pharmacopuncture*. 2019; 22(4): 239-247. doi: 10.3831/KPI.2019.22.032.

26. Kalantari H, Shamsi Ehsan T, Samimi A, Kheradmand P, Shirani M and Zeidooni L. Histopathological and biomedical parameters determination in the protective effect of hydroalcoholic extract of *Allium jesdianum* on hepatotoxicity induced by bromobenzene in mice. *Iran. J. Pharm. Sci.* 2018; 14(2): 15-24.

27. Kalantari H, Keshavarzi E Kheradmand P, Goodarzian M and Zeidooni L. Protective effect hydroalcoholic extract *Allium jesdianum* Boiss against CCl4-induced hepatotoxicity in male mice. *J. Rep. Pharm. Sci.* 2018; 7(3): 389-399. doi: 10.4103/2322-1232.254814.

28. Jalili C, Golshani N, Bahrami Y, Roshankhah S, Salahshoor M.R. Protective efficacy of *Allium jesdianum* on liver parameters, inflammatory cytokines, oxidative injury and apoptotic changes in mercuric chloride-induced hepatotoxicity in rats. *Eur. J. Anat.* 2020; 24(5): 381-389.

29. Razavi-Azarkhiavi K, Behravan J, Mosaffa F, Sehatbakhsh S, Shirani K and Karimi GR. Protective effects of aqueous and ethanol extracts of rosemary on H2O2-induced oxidative DNA damage in human lymphocytes by comet assay. *J. Complement. Integr. Med.* 2014; 11(1): 27-33. doi: 10.1515/jcim-2013-0063.

30. Yousefsani BS, Boozari M, Shirani K, Jamshidi AH and Dadmehr M. A review on phytochemical and therapeutic potential of *Iris germanica*. *J. Pharm. Pharmacol.* 2021; 73(5): 611-625. doi: 10.1093/jpp/rgab008.

31. Ghasemi Pirbalouti A. Phytochemical and bioactivity diversity in the extracts from bulbs and leaves of different populations of *Allium jesdianum*, a valuable underutilized vegetable. *Acta Sci. Pol. Hortorum Cultus.* 2019; 18(2): 115-122. doi: 10.24326/asphc.2019.2.11.

Journal of Medicinal Plants

Downloaded from jmp.ir on 2025-07-05

32. Ahmadzadeh CY, Ghasemi PA and Karimia L. Antioxidant activities and total phenolic content of ethanol extracts from *Allium hirtifolium* (Boiss) bulbs and *Allium jesdianum* (Boiss) leaves. In: 3rd National Congress on Medicinal Plants (NCMP 2014). Mashhad, Iran. 2014; 3: 368.

33. Mousavi R, Haghighi Fashi Z, Hadipour Jahromy M and Rasooli R. Potential of the *Allium jesdianum* extract in suppression of anxiety and depression in mice. *British. Biomed. Bull.* 2017; 5(2): 302-307. doi: 10.21767/2471-9897.1000302.

34. Esmaeili M, Ebrahimi Kahrizsangi A, Shahrokh Shahraki S and Mokhtari A. Evaluation of the effects of *Allium jesdianum* bacterial endophytes on some bacterial and fungal pathogens. *Iran. J. Med. Microbiol.* 2021; 15(3): 281-301. doi: 10.30699/ijmm.15.3.281.

35. Hojjati M and Alizadeh Behbahani B. Evaluation of the effect of aqueous and methanolic extraction methods on the antioxidant and antimicrobial characteristics of Allium jesdianum extract: in vitro study. *Iran. Food. Sci. Technol. Res. J.* 2021; 17(1): 83-91.

36. Yousefsani B S, Ghobadi A and Shirani K. Uncovering the neuroprotective powers of *Allium sativum*: exploring its potential to alleviate malathion- associated Parkinson's-like behavioral symptoms in a rat model. *J. Med. Plants* 2024; 23(89): 68-82.

37. Kamranfar F, Pourahmad Jaktaji R, Shirani K, Jamshidi AH, Samiei F, Arjmand AH, Khoramjouy M, Faizi M and Pourahmad J. Protective effect of a standardized *Allium jesdianum* extract in an Alzheimer's disease induced rat model. *Neurosci. Lett.* 2023; 815: 137491. doi: 10.1016/j.neulet.2023.137491.

38. Vahdati Hassani F, Rezaee R, Sazegara H, Hashemzaei M, Shirani K and Karimi GR. Effects of *Silymarin* on neuropathic pain and

formalin-induced nociception in mice. *Iran. J. Basic. Med. Sci.* 2015; 18(7): 715-720.

39. Khaksarian M, Gholami E, Alipour M, Sabooteh T and Asadi-Samani M. Investigation of the effects of the essence and extract of *Allium jesdianum* on the activity of COX-1 and COX-2 enzymes. *Int. J. Adv. Biotechnol. Res.* 2017; 8(2): 1095-1101.

40. Khaksarian M, Meshkatosadat MH, Farazi R, Safarpour F. A study of chemistry and Antinociceptive properties of medicinal plant *Allium jesdianum* leaves and the probable role of Opioidergic system. *Yafte*. 2008; 9(4): 21-26.

41. Lorigooini Z, Ayatollahi SA, Amidi S and Kobarfard F. Evaluation of anti-platelet aggregation effect of some *Allium* species. *Iran. J. Pharm. Res.* 2014; 14(4): 1225-1231. doi: 10.22037/JJPR.2015.1769.

42. Shirani K, Vahdati Hassani F, Razavi-Azarkhiavi K, Heidari S, Riahi Zanjani B and Karimi GR. Phytotherapy of cyclophosphamideinduced immunosuppression. *Environ. Toxicol. Pharmacol.* 2015; 39(3): 1262-1275. doi: 10.1016/j.etap.2015.04.012.

43. Abedi F, Yousefsani BS and Shirani K. The protective effects of *Allium ampeloprasum* Subsp Iranicum on cyclophosphamide-induced immunosuppression in NMRI Mice: A promising natural immunomodulator. *Avicenna*. *J. Phytomed.* 2023; 1-12. doi: 10.22038/ajp.2024.24595.

44. Shiehzadeh F, Vakili P and Shirani K. Evaluation of immune responses induced by hydroalcoholic extract of *Allium ampeloprasum*. L *subsp iranicum* in mice: *Allium iranicum* as a powerful immune system booster. *Iranian. J. Pharmaceutical. Sci.* 2022; 18(3): 176-182.

45. Rahimi S and Madani M. Evaluation of different concentrations of ethanolic extract of *Allium jesdianum* on Humoral immunity in infected mice with *Candida albicans*. Presented

Journal of Medicinal Plants

at the International and Iranian Congress of Microbiology. 2015, 16. [Online]. Available: https://sid.ir/paper/940999/en.

46. Radjabian T, Hosseinpur Yektaei Z, Naghizadeh MM and Ghazanfari T. Immunomodulatory impacts of bulbs extracts from five *Allium* species on IFN-γ, IL-4, and IL-17 Cytokines. *Immunoregulation*. 2021; 4(2): 91-100. doi:

10.32598/IMMUNOREGULATION.4.2.4.

47. Naeini AR, Yaraee R and Shokri H. Antifungal and immunomodulatory activity of *Allium jesdianum* Boiss extracts. *J. Herbmed. Pharmacol.* 2020; 9(1): 75-80. doi: 10.15171/jhp.2020.11.

48. Rezaie AR, Yousefsani BS, Omidi A, Shirani K. Exploring the Immune-Boosting and Hepatoprotective Potential of *Allium jesdianum* against Cyclophosphamide-Induced Toxicity in Mice: A Promising Approach for Immunomodulation. *Avicenna. J. Phytomed* [A ccepted].

49. Shahrokh S, Vahedi Gh, Khosravi AR, Mahzounieh MR, Ebrahimi A, Sharifzadeh A and Balal A. In vitro antifungal activity of aqueous-ethanolic extract of *Allium jesdianum* against fluconazole-susceptible and -resistant human vaginal *Candida glabrata* isolates. *J. Herbmed. Pharmacol.* 2017; 6(3): 136-140.

50. Sarlak M, Nasiri Z, Madani M, Shahanipour K, Tognoni E and Shakib P. Investigating the effects of aqueous and alcoholic extracts of *Allium hirtifolium* and *Allium jesdianum* on Keratinase activity of Trichophyton Mentagrophytes. *Curr. Drug Ther.* 2022; 17(1):

39-44. doi: 10.2174/ 1574885517666211216160301.

51. Razzaghi Khezezlo S, Sharifzadeh A, Soltani M, Shokri H and Khosravi A. Determination of genetic diversity and susceptibility of Trichophyton mentagrophytes isolates against antifungal effects of ethanolic extract of *Allium jesdianum. Iran. Vet. J.* 2017; 13(3): 56-66. doi: 10.22055/IVJ.2017.45921.1669.

52. Kahkesh H and Zargar B. Corrosion protection evaluation of *Allium jesdianum* as a novel and green source inhibitor for mild steel in 1M HCl solution. *J. Mol. Liq.* 2021; 344: 117768. doi: 10.1016/j.molliq.2021.117768.

53. Kogani S and Roomiani L. The effect of alcoholic extract of *Allium jesdanum*, *Tragopon carcifolus* and *Fumaria parviflora* on the shelf life of Rutilus kutum. *Iran. Sci. Fish. J.* 2020; 29(3): 93-108.

54. Ekrami M, Ekrami A, Hosseini MA and Emam-Djomeh Z. Characterization and optimization of Salep Mucilage Bionanocomposite films containing Allium jesdianum Boiss. Nanoliposomes for antibacterial food packaging utilization, Molecules. 2022; 27(20): 7032. doi: 10.3390/molecules27207032.

How to cite this article: Shamsipour Nehzomi Z, Shirani K. A comprehensive review of the pharmacological properties and medicinal uses of *Allium jesdianum* Boiss. & Buhse. *Journal of Medicinal Plants* 2024; 23(90): 1-16. doi: 10.61186/jmp.23.90.1



فصلنامه گياهان دارويي



Journal homepage: <u>www.jmp.ir</u>

مقاله مرورى

مروری بر خواص دارویی و کاربردهای درمانی گیاه پباز یزدی (Allium jesdianum Boiss. & Buhse) زهرا شمسی پور نهزمی، کبری شیرانی*

گروه سم شناسی، دانشکاده علوم پزشکی، دانشگاه تربیت مدرس، تهران، ایران

چکیدہ	اطلاعات مقاله
مقدمه : Allium jesdianum Boiss. & Buhse یک گیاه بومی از خانواده Amaryllidaceae است که در مناطق	گلواژگان:
خاورمیانه یافت میشود. به دلیل خواص دارویی و بیولوژیکی متعدد آن، از این گیاه برای درمان بسیاری از	Allium jesdianum
بیماریها، از جمله سرماخوردگی، عفونتها و سنگ کلیه استفاده میشود. هدف : در این مرور، ما به بررسی	فعاليت بيولوژيكي
کاربردهای دارویی، خواص فیتوشیمیایی و خواص دارویی A. jesdianum پرداختیم. روش بررسی : کاربردهای	
دارویی و خواص فیتوشیمیایی و دارویی A. jesdianum با استفاده از پایگاههای دادهای چون Google Scholar،	ىر كيبات قيبوسيميايي
Scopus ،PubMed و Web of Science با استفاده از کلیدواژههایی نظیر A. jesdianum، فعالیت بیولوژیکی،	اثرات دارویی
تركيبات فيتوشيميايي، اثرات دارويي، گياهان دارويي مورد جستجو قرار گرفت. نتايج : مطالعات نشان دادهاند كه	گیاهان دارویی
A. jesdianum حاوی ترکیبات شیمیایی مختلفی از جمله فلاونوئیدها، پلیفنولها و ترکیبات گوگردی است.	0
علاوه بر این، اثرات آنتیاکسیدانی، ضدسرطانی، مسکن، ضدافسردگی، ضدقارچی، ضدباکتری، محافظ کلیه،	
محافظ کبد، تقویتکننده سیستم ایمنی، محافظت در برابر اشعه و اثرات هماتولوژیکی این گیاه نیز مورد تأکید	
قرار گرفته است. همچنین، A. jesdianum می تواند به عنوان یک نگهدارنده طبیعی برای بهبود پایداری اکسیداتیو	
غذاهای غنی از اسیدهای چرب غیراشباع، کنترل رشد میکروارگانیسمها و تقویت ایمنی غذاها استفاده شود.	
نتیجهگیری : بر اساس نتایج بهدستآمده، A. jesdianum را می توان به عنوان یک ترکیب گیاهی با خواص ضد	
التهابی و آنتیاکسیدانی در نظر گرفت که میتواند به عنوان یک عامل محافظ در برابر سموم شیمیایی و استفاده	
مفرط از داروها مفید باشد. با این حال، مطالعات بیشتری برای بررسی اثرات بالینی این عصاره بر سلامت انسان	
مورد نیاز است.	

B; Bcl-2 T بیماری لنفومای Bcl-2 ۲- بیماری لنفومای Bcl-2 ۲- بیماری لنفومای Bcl-2 ۲- بیماری لنفومای B; Bcl-2 T بیماری لنفومای ALT، مینوترانسفراز؛ Bcl-2 ۲- Bcl-2 بیماری لنفومای ALT، مینوترانسفراز؛ Bcl-2 ۲- بیماری لنفومای COX-2 ۲- (COX-4 بی و تفای Bcl-2 ۲۰ مینوتران Bcl-2 مینوتران Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میکروگرم Back ۲۰ میلی Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میکروگرم در میلی Bcl-4 ۱۰ پروتئین شبیه کیناز خط میکس-2 میکس-2 میلی می متالوپروتئین Top ۲۰ میکنوتران Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میکروگرم در میلی Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میکروگرم در میلی Bcl-2 ۲۰ میکروگرم در میلی Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میکروگره Bcl-2 ۲۰ میکروگرم در میلی Bcl-2 ۲۰ میکرو تولی Bcl-2 ۲۰ میکرولی Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میکروگرم در میلی Bcl-2 ۲۰ میکروگرم در میلی Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میلی Bcl-2 ۲۰ میکروگرم در میلی Bcl-2 ۲۰ میکروگرم در میلی Bcl-2 ۲۰ میکروگرم در میلی Bcl-2 ۲۰ میکروگرم در

تاریخ دریافت: ۹ خرداد ۱۴۰۳؛ تاریخ دریافت اصلاحات: ۸ مهر ۱۴۰۳؛ تاریخ پذیرش: ۱۰ مهر ۱۴۰۳

doi: 10.61186/jmp.23.90.1

© 2023. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<u>https://creativecommons.org/licenses/by-nc/4.0/</u>)