

Research Article

Phytochemical constituents of the fruits of *Kelussia odoratissima* Mozaff., an aromatic plant endemic to Iran

Mahnaz Khanavi^{1,2}, Shima Ghadami¹, Ghazal Sadaghiani-Tabrizi¹, Mohammad-Reza Delnavazi^{1,*}

¹ Department of Pharmacognosy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

² Faculty of Land and Food Systems, University of British Columbia, Vancouver, BC, Canada

ARTICLE INFO

Keywords:

Kelussia odoratissima
Apiaceae
Flavonoid
Essential oil
(Z)-Ligustilide

ABSTRACT

Background: *Kelussia odoratissima* Mozaff. from Apiaceae family is a perennial herbaceous plant endemic to the west of Iran. The aromatic aerial parts of this species are traditionally used by indigenous people to flavor some local foods, as well as for various therapeutic purposes. **Objective:** The present study was designed to analyze phenolic compounds and essential oil constituents of *K. odoratissima* fruits. **Methods:** The *n*-butanol fraction obtained from hydroalcoholic extract of *K. odoratissima* fruits was investigated by chromatography on normal phase and Sephadex LH-20 columns. Chemical structures of the isolated compounds were clarified by ¹H-NMR and ¹³C-NMR spectral analyses. Essential oil constituents of the fruits were also analyzed using GC-MS. **Results:** Phytochemical investigation of the *K. odoratissima* fruits resulted in the isolation of five flavonol glycosides; isorhamnetin 3-O-glucoside (**1**), quercetin 3-O-glucoside (isoquercetin) (**2**), isorhamnetin 3-O-rutinoside (narcissin) (**3**), isorhamnetin 3-O-glucuronide (**4**) and quercetin 3-O-glucuronide (mequilianin) (**5**). GC-MS analysis of the fruits essential oil led to the identification of the thirty six compounds, of which (Z)-ligustilide (15.93 %), δ -cadinene (12.26 %) and germacrene D (12.18 %) were the main compounds. **Conclusion:** The results of this study introduce *K. odoratissima* fruits as a source of flavonoid glycosides and phthalate derivatives. The presence of these compounds with known biological properties and health beneficial effects provides more medicinal potentials for the fruits of *K. odoratissima* and suggest it an appropriate option for further studies.

1. Introduction

The genus *Kelussia* from Apiaceae family is represented by only one member, *Kelussia odoratissima* Mozaff., which is found in central Zagros mountains, west of Iran [1]. The aerial

parts of this aromatic species are used by indigenous people under the local names of "Kellos" and "Karafse-Bakhtiari" as vegetable, as flavoring agent in yogurt, for the treatment of indigestion, rheumatism, gastric ulcer, cough,

Abbreviations: NMR, Nuclear magnetic resonance; GC-MS, Gas Chromatography-Mass Spectrometry; TLC, Thin Layer Chromatography

* Corresponding author: delnavazi@tums.ac.ir

doi: 10.52547/jmp.20.79.1

Received 18 February 2021; Received in revised form 19 June 2021; Accepted 3 July 2021

© 2020. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>)

pain, diabetes and as a sedative agent [2, 3]. A review on literature shows that extracts and essential oils obtained from different parts of *K. odoratissima* have been reported for their biological and pharmacological activities such as antioxidant [4, 5], antibacterial [6], larvicidal [7], antileishmanial [8], cytotoxic [9], spasmolytic [10], anti-inflammatory [11], antihypertensive [12], sedative and anxiolytic [13] effects. One study reported the isolation of two steroids, stigmasterol and β -sitosterol and one phthalide, 3-butylyden-4,5-dihydrophthalide from the *n*-hexane extract of *K. odoratissima* fruits [14]. Furthermore, they identified thirty eight compounds in the fruits essential oil, of which (*Z*)-ligustilide (29.2 %), germacrene B (15.9 %) and germacrene D (15.5 %) were the major compounds [14]. Another study reported the analysis of fatty acids in the fruits of *K. odoratissima* and showed that among the five fatty acids identified in the fruits oil, petroselinic acid (72.35 %) and linoleic acid (19.14 %) were the main acids [15]. In the mentioned study, *K. odoratissima* fruits were also reported as a source of phenolic compounds by a total phenolic content of 218.15 milligrams gallic acid equivalents (GAE) per gram of dry fruit weight [15]. Beside the high phenolic content, there is not any report on phenolic principles of *K. odoratissima* fruits. Therefore, the present study was designed to isolation and structural elucidation of the phenolic compounds present in the fruits of this valuable medicinal plant. The essential oil composition of the fruits was also identified and compared by related data previously reported for *K. odoratissima*.

2. Materials and Methods

2.1. Plant material

The fruits of *Kelussia odoratissima* Mozaff. were purchased from Pakan-Bazr Co., Isfahan,

Iran (Plant source: Fereydunshahr region, Isfahan, Iran.; Collection date: July 2017). The identity of the fruits was authenticated by botanist Dr. Yousef Ajani (Research Institute of Forest and Rangelands, Tehran, Iran) and the code of PMP-2694 was assigned for it at the herbarium of Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran.

2.2. Extraction of Phenolics

The dried and comminuted fruits of *K. odoratissima* (1.4 kg) were subjected to extraction using maceration method with dichloromethane and methanol-water (70:30), successively (4 × 5 L, each). The obtained extracts were concentrated using a rotary evaporator at 45 °C under the reduced pressure and then dried in vacuum oven. Hydroalcoholic extract (250 g) was suspended in water (0.5 L) and fractionated by equal volume of *n*-butanol (×3) to extract its phenolic compounds.

2.3. Isolation and purification of the compounds

A portion of the *n*-butanol fraction (20 g) was chromatographed on a normal phase silica gel column (Mesh 230-400, Merck) using a gradient solvent mixture of CHCl₃-MeOH (100:0 to 50:50) to get eighteen subfractions (B1-B18). Thin layer chromatography (Pre-coated silica gel GF₂₅₄ plates, Merck) was applied to monitor column chromatography and fractions giving similar spots under UV (254 and 366 nm) followed by the exposure to ammonia vapor were combined. Based on the TLC analysis, subfractions B6, B10, B11, B14 and B16 were chosen for further isolation. Column chromatography of these subfractions (B6, B10, B11, B14 and B16) on a Sephadex LH-20 column using methanol as mobile phase resulted in the isolation of compounds **1** (19.0 mg), **2**

(13.5 mg), **3** (35.4 mg), **4** (23.8 mg) and **5** (18.1), respectively. The structures of the compounds were elucidated by $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ (Varian- INOVA, 500 MHz for $^1\text{H-NMR}$ and 125 MHz for $^{13}\text{C-NMR}$) spectral analysis.

2.4. Essential oils extraction

The comminuted fruits (100 g) were subjected to hydrodistillation for 3 hours, using a Clevenger apparatus. The obtained oils were dried over anhydrous sodium sulfate and kept at 4 °C until analysis.

2.5. GC-MS analysis

The essential oil extracted from the fruits of *K. odoratissima* were analyzed on a HP-6890 gas chromatograph with a BPX5 column (30 m \times 0.25 mm id, 0.25 μm film thickness), equipped with HP-5973 mass detector (Ionization energy: 70 eV) under the following conditions; 5 min after injection, oven temperature was increased from 50 °C to 240 °C at a rate of 3 °C/min and then reached to 300 °C at the rate of 15 °C/min and hold 3 min in this temperature. Injector temperature: 250 °C, detector temperature: 220 °C, injection volume: 1.0 μl , split ratio: 1:35, carrier gas: helium (99.999 %, Flow rate: 0.5 ml/min). The retention indices (RIs) were calculated for all identified compounds using a homologous series of *n*-alkanes (C₈-C₂₄) injected under the same conditions described for the analyzed essential oil. Identification of the compounds were carried out based on computer matching with the Wiley 275.L library, as well as by comparison of RIs and mass fragmentation patterns with those published for standard compounds [16].

3. Results

3.1. Isolation and structural elucidation

Phytochemical analysis of the fruits of *K. odoratissima* using chromatography on normal phase silica gel and Sephadex LH-20 columns resulted in the isolation of five compounds (**1-5**) from *n*-butanol fraction of hydroalcoholic extract. The structures of the isolated compounds were characterized as isorhamnetin 3-O- β -D-glucopyranoside (**1**), quercetin 3-O- β -D-glucopyranoside (isoquercetin) (**2**), isorhamnetin 3-O- β -D-glucopyranosyl-(6 \rightarrow 1)- α -L-rhamnopyranoside (isorhamnetin 3-O- β -D-rutinoside, narcissin) (**3**), isorhamnetin 3-O- β -D-glucuronide (**4**) and quercetin 3-O- β -D-glucuronide (mequilianin) (**5**) (Fig. 1) using $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectral analyses, as well as by comparison with published data [17-21].

3.1.1. Spectroscopic data of the isolated compounds

Compound **1**; *Isorhamnetin 3-O- β -D-glucopyranoside* (C₂₂H₂₂O₁₂); Yellow amorphous solid; R_f = 0.6 (CHCl₃-MeOH, 8:2); $^1\text{H-NMR}$ (DMSO-*d*₆, 500 MHz): δ_{H} 7.93 (1H, *d*, *J* = 2.0 Hz, H2'), 7.50 (1H, *dd*, *J* = 8.5 and 2.0 Hz, H6'), 6.93 (1H, *d*, *J* = 8.5 Hz, H5'), 6.46 (1H, *d*, *J* = 2.0 Hz, H8), 6.22 (1H, *d*, *J* = 2.0 Hz, H6), 5.56 (1H, *d*, *J* = 7.4 Hz, H1''), 3.83 (3H, *s*, OCH₃), 3.1-3.7 (6H, *overlapped signals*, H2''-H6''). $^{13}\text{C-NMR}$ (DMSO-*d*₆, 125 MHz): δ_{C} 177.78 (C4), 164.55 (C7), 161.35 (C5), 156.87 (C9), 156.73 (C2), 149.63 (C3'), 147.28 (C4'), 133.46 (C3), 122.51 (C6'), 121.58 (C1'), 115.57 (C5'), 113.87 (C2'), 104.43 (C10), 101.18 (C1''), 99.10 (C6), 94.22 (C8), 77.76 (C5''), 76.62 (C3''), 74.63 (C2''), 70.12 (C4''), 60.93 (C6''), 56.15 (OCH₃) [17].




Fig. 1. The structures of isolated compounds (1-5) from *K. odoratissima* fruits

Compound **2**; *Quercetin 3-O-β-D-glucopyranoside (isoquercetin)* (C₂₁H₂₀O₁₂); Yellow amorphous solid; R_f = 0.5 (CHCl₃-MeOH, 8:2); ¹H-NMR (DMSO-*d*₆, 500 MHz): δ_H 7.58 (1H, *d*, *J* = 2.0 Hz, H2'), 7.56 (1H, *dd*, *J* = 9.0 and 2.0 Hz, H6'), 6.86 (1H, *d*, *J* = 9.0 Hz, H5'), 6.42 (1H, *d*, *J* = 2.0 Hz, H8), 6.21 (1H, *d*, *J* = 2.0 Hz, H6), 5.45 (1H, *d*, *J* = 7.5 Hz, H1''), 3.1-3.7 (6H, *overlapped signals*, H2''-H6''). ¹³C-NMR (DMSO-*d*₆, 125 MHz): δ_C 177.82 (C4), 164.52 (C7), 161.37 (C5), 156.90 (C9), 156.60 (C2), 148.69 (C4'), 145.07 (C3'), 133.80 (C3), 122.09 (C6'), 121.63 (C1'), 116.55 (C5'), 115.60 (C2'), 104.36 (C10), 101.29 (C1''), 99.05 (C6), 94.02 (C8), 77.88 (C5''), 76.73 (C3''), 74.42 (C2''), 70.21 (C4''), 61.26 (C6'') [18].

J = 1.8 Hz, H2'), 7.51 (1H, *dd*, *J* = 8.4 and 1.8 Hz, H6'), 6.91 (1H, *d*, *J* = 8.4 Hz, H5'), 6.44 (1H, *d*, *J* = 2.0 Hz, H8), 6.21 (1H, *d*, *J* = 2.0 Hz, H6), 5.40 (1H, *d*, *J* = 7.4 Hz, H1''), 4.40 (1H, *br s*, H1'''), 3.83 (3H, *s*, OCH₃), 3.0-3.7 (10H, *overlapped signals*, H2''-H6'' and H2'''-H-5'''), 0.94 (3H, *d*, *J* = 6.2 Hz, H6'''); ¹³C-NMR (DMSO-*d*₆, 125 MHz): δ_C 177.71 (C4), 164.34 (C7), 161.26 (C5), 157.01 (C9), 156.94 (C2), 149.57 (C4'), 147.26 (C3'), 133.48 (C3), 122.77 (C6'), 121.54 (C1'), 115.57 (C5'), 113.63 (C2'), 104.44 (C10), 101.54 (C1''), 101.25 (C1'''), 99.08 (C6), 94.31 (C8), 76.54 (C5''), 76.20 (C3''), 74.52 (C2''), 72.03 (C4''), 70.82 (C3'''), 70.57 (C2'''), 70.39 (C4''), 68.68 (C5'''), 67.37 (C6''), 56.12 (OCH₃), 17.99 (C6''') [19].

Compound **3**; *Isorhamnetin 3-O-β-D-rutinoside (narcissin)* (C₂₈H₃₂O₁₆); Yellow amorphous solid; R_f = 0.3 (CHCl₃-MeOH, 8:2); ¹H-NMR (DMSO-*d*₆, 500 MHz): δ_H 7.83 (1H, *d*,

Compound **4**; *Isorhamnetin 3-O-β-D-glucuronide* (C₂₂H₂₀O₁₃); Yellow amorphous solid; R_f = 0.5 (CHCl₃-MeOH, 8:2); ¹H-NMR (DMSO-*d*₆, 500 MHz): δ_H 8.06 (1H, *br s*, H2'),

7.36 (1H, *br d*, $J = 8.4$, H6'), 6.87 (1H, *d*, $J = 8.4$ Hz, H5'), 6.16 (1H, *br s*, H8), 5.99 (1H, *br s*, H6), 5.57 (1H, *d*, $J = 7.7$ Hz, H1''), 3.80 (3H, *s*, OCH₃), 3.21-3.39 (4H, *overlapped signals*, H2''-H5''). ¹³C-NMR (DMSO-*d*₆, 125 MHz): δ_c 177.55 (C4), 173.13 (C6''), 165.44 (C7), 161.21 (C5), 156.81 (C9), 156.81 (C2), 149.38 (C4'), 147.14 (C3'), 133.23 (C3), 122.24 (C6'), 121.68 (C1'), 115.31 (C5'), 114.48 (C2'), 103.68 (C10), 101.28 (C1''), 99.02 (C6), 94.28 (C8), 76.42 (C3''), 76.61 (C5''), 74.35 (C2''), 72.44 (C4''), 56.13 (OCH₃) [20].

Compound **5**; *Quercetin 3-O- β -D-glucuronide (mequilianin)* (C₂₁H₁₈O₁₃); Yellow amorphous solid; $R_f = 0.4$ (CHCl₃-MeOH, 8:2); ¹H-NMR (DMSO-*d*₆, 500 MHz): δ_H 8.23 (1H, *br s*, H2'), 7.33 (1H, *dd*, $J = 8.4$ and 1.2 Hz, H6'), 6.83 (1H, *d*, $J = 8.4$ Hz, H5'), 6.24 (1H, *br s*, H8), 6.06 (1H, *br s*, H6), 5.15 (1H, *d*, $J = 7.3$ Hz, H1''), 3.1-3.5 (4H, *overlapped signals*, H2''-H6''). ¹³C-NMR (DMSO-*d*₆, 125 MHz): δ_c 177.27 (C-4), 172.95 (C6''), 165.06 (C7), 160.99 (C5), 157.25

(C9), 157.25 (C2), 148.62 (C4'), 145.04 (C3'), 134.35 (C3), 121.05 (C6), 121.05 (C1'), 118.10 (C-5'), 115.81 (C-2'), 103.52 (C10), 103.52 (C1''), 100.27 (C-6), 94.79 (C-8), 76.92 (C-3''), 74.54 (C-5''), 74.41 (C-2''), 72.05 (C-4'') [21].

3.2. Essential oil composition

Hydrodistillation of the fruits of *K. odoratissima* led to the extraction of a pale yellowish oil with the yield of 1.5 % (v/w). GC-MS analysis of the obtained essential oil resulted in the identification of the thirty six compounds, representing 90.74 % of the oil (Table 1). Among the identified compounds, (*Z*)-ligustilide (15.93 %), δ -cadinene (12.26 %) and germacrene D (12.18 %) were the main compounds (Fig. 2) and oxygenated sesquiterpenes with relative percentage of 53.68 % were the main group of constituents identified in essential oil of *K. odoratissima* fruits.


Table 1. Chemical composition of the essential oil of *K. odoratissima* fruits obtained by hydrodistillation method

No.	Compound name ^a	%	RI ^b
1	α -Pinene	0.09	939
2	β -Pinene	0.35	978
3	2-Ethyl-2-hexenal	0.71	981
4	β -Phellandrene	3.23	1027
5	5-Pentylcyclohexa-1,3-diene	1.43	1161
6	Lavandulol	0.14	1170
7	Acetic acid octyl ester	0.24	1215
8	Citronellol	0.27	1227
9	Lavandulyl acetate	0.75	1292
10	Ylangene	0.62	1374
11	α -Copaene	2.92	1378
12	β -Cubebene	0.48	1388
13	β -Elemene	2.04	1391
14	α -Barbatene	0.42	1415
15	Lavandulyl isobutyrate	3.80	1417
16	γ -Elemene	4.34	1435
17	Sativene	1.36	1437
18	α -Humulene	1.09	1453
19	γ -Muurolene	0.49	1481
20	Germacrene D	12.18	1483

Table 1. Chemical composition of the essential oil of *K. odoratissima* fruits obtained by hydrodistillation method (Continued)

No.	Compound name ^a	%	RI ^b
21	Alloaromadendrene	1.81	1487
22	γ -Amorphene	1.07	1492
23	α -Selinene	1.84	1498
24	α -Muurolene	1.78	1504
25	γ -Cadinene	2.80	1512
26	δ -Cadinene	12.26	1521
27	Selina-3,7(11)-diene	2.16	1532
28	Neryl 2-methyl butanoate	4.81	1571
29	Germacrene B	4.02	1556
30	Spatulenol	0.56	1582
31	Salvial-4(14)-en-1-one	0.27	1599
32	α -epi-Muurolol	0.47	1648
33	3-Butylphthalide	0.61	1658
34	(Z)-Butylidene phthalide	3.05	1678
35	(Z)-Ligustilide	15.93	1797
36	(E)-Ligustilide	0.35	1741
	Monoterpene hydrocarbons	3.67	
	Oxygenated monoterpenes	9.77	
	Sesquiterpene hydrocarbons	53.68	
	Oxygenated sesquiterpenes	1.30	
	Non-terpenes	22.32	
	Total identified	90.74	

^a Compounds listed in order of elution from BPX5 column; ^b Retention indices to C₈-C₂₄ n-alkanes on BPX5 column.

**Fig. 2.** The structures of main compounds identified in essential oil of *K. odoratissima* fruits

4. Discussion

Five flavonol 3-O-glycosides, namely, isorhamnetin 3-O- β -D-glucopyranoside (**1**), quercetin 3-O- β -D-glucopyranoside (isoquercetin) (**2**), isorhamnetin 3-O- β -D-rutinoside (narcissin) (**3**), isorhamnetin 3-O- β -D-glucuronide (**4**) and quercetin 3-O- β -D-glucuronide (mequilianin) (**5**) were isolated from the fruits of *K. odoratissima* in present study for the first time.

Previously, two steroid derivatives, stigmasterol and β -sitosterol and one phthalide, 3-butyliden-4,5-dihydrophthalide reported from *n*-hexane extract of *K. odoratissima* fruits [14]. Flavonoids are phenolic compounds with well-known free radical scavenging activity which makes them appropriate food supplements to prevent oxidative stress related diseases such as cancers, diabetes, cardiovascular, inflammatory

and neurodegenerative diseases (such as Alzheimer and Parkinson) [22].

The potent antioxidant activity of isorhamnetin 3-O- β -D-rutinoside, a major flavonol 3-O-glycoside isolated in present study, has been reported in DPPH (IC₅₀; 9.01 μ M, IC₅₀ value of ascorbic acid as positive control; 11.93 μ M) and ONOO⁻ (IC₅₀; 2.56 μ M, IC₅₀ value of DL-penicillamin as positive control; 5.1 μ M) methods [23]. This compound (isorhamnetin 3-O- β -D-rutinoside) has also been reported as one of antimicrobial principles of *Atriplex halimus* aerial parts with considerable antibacterial activity against *Streptococcus pyogenes* (inhibition zone; 17 \pm 0.09 mm), *Escherichia coli* (inhibition zone; 16 \pm 0.09 mm) and *Acinetobacter baumannii* (inhibition zone; 16 \pm 0.13 mm), having a low cytotoxicity on PMNCs (IC₅₀; 450 μ g ml⁻¹) [24].

Isorhamnetin 3-O- β -D-glucopyranoside and quercetin 3-O- β -D-glucopyranoside (isoquercetin), two other flavonol glycosides isolated from *K. odoratissima* fruits in present study, have been documented for its antioxidant [25], antidiabetic [26] and hepatoprotective [27] activities. Furthermore, in a bioassay-guided fractionation study quercetin 3-O-glucoside was isolated as active compounds of *Sambucus ebulus* L. leaves with remarkable wound healing activity [28].

Isorhamnetin and quercetin were also isolated as their 3-O- β -D-glucuronide derivatives in the present study. A bioactivity-guided isolation study reported the isolation of isorhamnetin 3-O- β -D-glucuronide as one of active compounds of *Sanguisorba officinalis* L. (Rosaceae) with considerable lipid accumulation inhibition on 3T3-L1 cells (IC₃₀; 18.43 μ M) [29]. *Chuquiraga spinosa* (Asteraceae), *Cichorium spinosum* L. (Asteraceae), *Foeniculum vulgare* L. (Apiaceae) and some cultivars of *Vitis vinifera* L. (Vitaceae) are examples of plants reported to contain isorhamnetin 3-O- β -D-glucuronide [30-34].

Quercetin 3-O- β -D-glucuronide has been found in some plant species such as *Calligonum comosum* L. (Polygonaceae), *Rubus ulmifolius* Schott. (Rosaceae) and *Phaseolus vulgaris* L. (Fabaceae), as well as one of major human metabolites of quercetin [35-38]. One study suggested that supplementation of quercetin 3-O- β -D-glucuronide can reduce the relative risk for developing Alzheimer's disease (AD) dementia [39]. They showed that quercetin 3-O- β -D-glucuronide can significantly reduce the generation of β -amyloid (A β) peptides and improves AD-type deficits in hippocampal formation basal synaptic transmission and long-term potentiation, possibly through mechanisms involving the activation of the c-Jun N-terminal kinases and the mitogen-activated protein kinase signaling pathways [39]. In another study, quercetin 3-O- β -D-glucuronide showed to possess anti-neuroinflammatory activity in lipopolysaccharide (LPS)-induced inflammation in BV2 cells by inhibition of the production of NO and PGE2 and reducing the levels of some pro-inflammatory cytokines such as TNF- α and interleukin-1 β [40]. Moreover, beneficial effects of quercetin 3-O- β -D-glucuronide in arteriosclerosis prevention has been reported via elevating plasma HDLC followed by induction of ABCA1 (ATP-binding cassette, subfamily A, member 1) expression, a crucial cholesterol transporter involved in reverse cholesterol transport [41].

In the present study, GC-MS analysis of the essential oil extracted from the fruits of *K. odoratissima* resulted in the identification of thirty six compounds, of which (*Z*)-ligustilide (15.93 %), δ -cadinene (12.26 %) and germacrene D (12.18 %) were the main compounds (Fig. 2). One study reported α -caryophyllene (22.60 %), α -humulene (20.0 %) and cyclopropane (11.54 %) as the main compounds of the essential oil of *K. odoratissima* fruits, gathered from Zardkooh

Mountain, (Charmahal-Bakhtiari, Iran) [42]. In another study on the fruits of this plant collected from Samsami region (Chaharmahal-Bakhtiari, Iran) (*Z*)-Ligustilide (86.0 %), (*2E*)-decen-1-ol (8.0 %), pentyl cyclohexa-1,3-diene (4.4 %) were characterized as main compounds of its essential oil [43]. The results of previous reports on

essential oil composition of different parts of *K. odoratissima* have been summarized in Table 2 [5, 7, 8, 13, 15, 43-47]. Beside possible genetic diversity, variation observed in essential oil constituents may be raised from some geographical differences between the populations of *K. odoratissima* [48].

Table 2. The results of essential oil analysis of *K. odoratissima* from the previous and present studies.

Location of collection	Date	Method	Part	Main compounds (%)	Ref.
Fereydunshahr, Isfahan, Iran	Dec 2018	HD ^a	Fruit	(<i>Z</i>)-Ligustilide (15.9 %), δ -Cadinene (12.3 %), Germacrene D (12.2 %)	Present study
Dishmook, Kohgiluyeh-Buyer Ahmad, Iran	May 2014	HD	Leaf	(<i>Z</i>)-Ligustilide (58.7%), Carvacrol (7.8%), <i>trans</i> -3-Butylidene phthalide (4.9%), Thymol (4.5%),	5
Keloseh region, Isfahan, Iran				(<i>Z</i>)-Ligustilide (53.5%), Thymol (7.9%), <i>trans</i> -3-Butylidene phthalide (3.3%)	
Fereydunshahr, Isfahan, Iran				(<i>Z</i>)-Ligustilide (51.3%), Thymol (8.7%), Carvacrol (3.2%)	
Central Zagros Mountain, Chaharmahal-Bakhtiari, Iran	Apr 2009	HD	Leaf	(<i>Z</i>)-Ligustilide (77.7%), 2-Octen-1-ol acetate (6.3%), (<i>E</i>)-Ligustilide (2.3%)	7
Kohgiluyeh-Buyer Ahmad, Iran	Apr-May 2012	HD	Aerial part	(<i>Z</i>)-Ligustilide (34.5%), (<i>E</i>)-Ligustilide (11.8%), 3-(<i>Z</i>)-Butylidene phthalide (8.8%), Dec-9-en-1-ol (5.9%)	8
Central Zagros mountain, Iran	Mar 2006	HD	Aerial part	(<i>Z</i>)-Ligustilide (85.9%), α -Copaene (1.4%), δ -Cadinene (0.7%)	13
Fereydunshahr, Isfahan, Iran	Aug 2009	HD	Fruit	<i>Z</i> -ligustilide (51.0%), δ -Terpinen-7-al (10.3%), δ -terpinene (5.3%), Cumin aldehyde (5.2%)	15
Zardkooh Mountain, Charmahal-Bakhtiari, Iran	Jul-Aug 2011	HD	Fruits	α -Caryophyllene (22.6%), α -Humulene (20.0%), Cyclopropane (11.5%)	42
Samsami region, Chaharmahal-Bakhtiari, Iran	Jun-Jul 2014	HD	Root	(<i>Z</i>)-Ligustilide (54.0%), (<i>2E</i>)-Decen-1-ol (10.7%), Pentyl cyclohexa-1,3-diene (6.4%), (<i>3Z</i>)-Butylidene phthalide (5.8%)	43
			Stem	(<i>Z</i>)-Ligustilide (58.7%), (<i>2E</i>)-Decen-1-ol (11.6%), Pentyl cyclohexa-1,3-diene (4.4%)	
			Leaf	(<i>Z</i>)-Ligustilide (66.8%), (<i>2E</i>)-Decen-1-ol (12.3%), Pentyl cyclohexa-1,3-diene (3.8%)	
			Flower	(<i>Z</i>)-Ligustilide (62.4%), Geranyl butyrate (9.0%), <i>trans</i> -Muurolo-4(14) 5-diene (5.5%)	
			Fruit	(<i>Z</i>)-Ligustilide (86.0%), (<i>2E</i>)-Decen-1-ol (8.0%), Pentyl cyclohexa-1,3-diene (4.4%)	

Table 2. The results of essential oil analysis of *K. odoratissima* from the previous and present studies (Continued)

Location of collection	Date	Method	Part	Main compounds (%)	Ref.
Keloseh region, Fereydunshahr, Isfahan, Iran	Aug 2007	HD	Stem	Borneol (36.9 %), Bornyl acetate (14.0 %), 1,8-Cineol (13.6 %), Camphor (9.5 %)	44
			Flower	1,8-Cineol (22.0%), Camphor (20.1%), α -Pinene (19.0%), Camphene (12.0%), Bornyl acetate (5.8 %)	
			Leaf	β -Terpinene (23.0%), Sabinene (9.0%), α -Thujene (8.4%)	
Fereydunshahr, Isfahan, Iran	Jul 2007	HD	Aerial part	(<i>Z</i>)-Ligustilide (87.6%), (<i>E</i>)-Ligustilide (3.2 %), Piperitone epoxide (3.1%)	45
Bazoft region, Chaharmahal- Bakhtiari, Iran	April 2008	HD	Aerial part	(<i>Z</i>)-Ligustilide (47.3%), (<i>3E</i>)-Butyldiene phthalide (17.38%), (<i>E</i>)-Ligustilide (6.3%), 2-Octen-1-ol acetate (5.4%)	46
Koohrang region, Chaharmahal- Bakhtiari, Iran				(<i>Z</i>)-Ligustilide (33.7%), (<i>3E</i>)-Butyldiene phthalide (20.1%), (<i>E</i>)-Ligustilide (6.6%), 2-Octen-1-ol acetate (5.2%)	
Samsami region, Chaharmahal- Bakhtiari, Iran				(<i>Z</i>)-Ligustilide (37.6%), (<i>3E</i>)-Butyldiene phthalide (19.9%), (<i>E</i>)-Ligustilide (7.0%), Kessane (5.3%)	
Zagros mountain, Iran	May-Jun 2012	HD	Aerial part	α -Pinene (20.1%), 1,8-Cineole (18.2%), (<i>Z</i>)- Ligustilide (15.5%)	47

(*Z*)-Ligustilide, a phthalide derivative identified as major constituent in most of previously studied *K. odoratissima* essential oils, has received attention for its interesting pharmacological and biological effects such as neuroprotective, anti-oxidation, anti-inflammatory, analgesic and anticancer effects [49].

5. Conclusion

The presence of flavonoid glycosides (1-5) and phthalide derivatives with known health beneficial effects make the fruits of *K. odoratissima* as a natural remedy with valuable therapeutic potentials and suggest it an appropriate option for further studies. Meanwhile, restricted distribution of *K. odoratissima* underline the importance of an appropriate conservation approach for the uses of this species for medicinal and food purposes.

Author contributions

MK: Study supervision and data interpretation; SG: Experimental analysis and preparation of manuscript draft; GS: Experimental analysis; MD: Original idea presentation, study design, study supervision, data interpretation and revision of the manuscript.

Conflict of interest

The authors declare that there is no conflict of interest.

Acknowledgment

This research was supported by Tehran University of Medical Sciences and Health Services grant (No. 48353).

References

1. Mozaffarian V. Flora of Iran, No.54: Umbelliferae. Tehran: Publication of Research Institute of Forests and Rangelands; 2007.
2. Amiri MS and Joharchi MR. Ethnobotanical knowledge of Apiaceae family in Iran: a review. *Avicenna J. Phytomed.* 2016; 6(6): 621-35. doi: 10.22038/AJP.2016.6696.
3. Ahmadi K, Omidi H, Amini Dehaghi M and Naghdi Badi H. A review on the botanical, phytochemical and pharmacological characteristics of *Kelussia odoratissima* Mozaff. *J. Med. Plants* 2020; 18(72): 30-45.
4. Ahmadi F, Kadivar M and Shahedi M. Antioxidant activity of *Kelussia odoratissima* Mozaff. in model and food systems. *Food Chem.* 2007; 105(1): 57-64. doi: 10.1016/j.foodchem.2007.03.056.
5. Akbarian A, Rahimmalek M, Sabzalian MR and Sarfaraz D. Variation in essential oil composition, phenolic, flavonoid and antioxidant activity of *Kelussia odoratissima* Mozaff based on three model systems. *J. Appl. Res. Med. Aromat. Plants* 2019; 13: 100208. doi: 10.1016/j.jarmap.2019.100208.
6. Sureshjani MH, Yazdi FT, Mortazavi SA, Behbahani BA and Shahidi F. Antimicrobial effects of *Kelussia odoratissima* extracts against food borne and food spoilage bacteria "in vitro". *Arch. Adv. Biosci.* 2014; 5(2): 115-20. doi:10.22037/JPS.V5I2.5943.
7. Vatandoost H, Sanei Dehkordi A, Sadeghi SMT, Davari B, Karimian F, Abai MR and Sedaghat MM. Identification of chemical constituents and larvicidal activity of *Kelussia odoratissima* Mozaffarian essential oil against two mosquito vectors *Anopheles stephensi* and *Culex pipiens* (Diptera: Culicidae). *Exp. Parasitol.* 2012; 132(4): 470-574. doi: 10.1016/j.exppara.2012.09.010.
8. Kheirabadi KP, Dehkordi SS and Kheibari P. Effect of *Kelussia odoratissima* Mozaff. essential oil on promastigot form of *Leishmania major* (in vitro). *J. HerbMed Pharmacol.* 2015; 4(1): 10-14.
9. Karimian H, Arya A, Fadaeinasab M, Razavi M, Hajrezaei M, Khan AK, Ali HM, Abdulla MA and Noordin MI. *Kelussia odoratissima* Mozaff. activates intrinsic pathway of apoptosis in breast cancer cells associated with S phase cell cycle arrest via involvement of p21/p27 in vitro and in vivo. *Drug Des. Devel. Ther.* 2017; 11: 337-50. doi: 10.2147/dddt.s121518.
10. Sedighi M, Rafieian-Kopaei M, Noori-Ahmadabadi M. *Kelussia odoratissima* Mozaffarian inhibits ileum contractions through voltage dependent and beta adrenergic receptors. *Life Sci. J.* 2012; 9(4): 1033-8.
11. Minaiyan M, Sajjadi SE, Naderi N and Taheri D. Anti-inflammatory effect of *Kelussia odoratissima* Mozaff. hydroalcoholic extract on acetic acid-induced acute colitis in rats. *J. Rep. Pharm. Sci.* 2014; 3(1): 28-35.
12. Safaeian L, Sajjadi SE, Javanmard SH and Gholamzadeh H. Antihypertensive and antioxidant effects of hydroalcoholic extract from the aerial parts of *Kelussia odoratissima* Mozaff. in dexamethasone-induced hypertensive rats. *Adv. Biomed. Res.* 2016; 5: 25. doi: 10.4103/2277-9175.176342.
13. Rabbani M, Sajjadi SE and Sadeghi M. Chemical composition of the essential oil from *Kelussia odoratissima* Mozaff. and the evaluation of its sedative and anxiolytic effects in mice. *Clinics.* 2011; 66(5): 843-8. doi: 10.1590/s1807-59322011000500022.
14. Sajjadi S, Shokoohinia Y and Mehrmiri P. Isolation and characterization of steroids, phthalide and essential oil of the fruits of

- Kelussia odoratissima* Mozaff., an endemic mountain celery. *Res. Pharm. Sci.* 2013; 8(1): 35-41.
15. Saeedi KA and Omidbaigi R. Chemical characteristics of the seed of Iranian endemic plant *Kelussia odoratissima*. *Chem. Nat. Compd.* 2010; 46(5): 813-5. doi: 10.1007/s10600-010-9754-7
16. Adams RP. Identification of essential oil components by gas chromatography/mass spectrometry. Carol Stream: Allured Publishing Corporation; 2007.
17. Aliotta G, Della Greca M, Monaco P, Pinto G, Pollio A and Previtera L. In vitro algal growth inhibition by phytotoxins of *Typha latifolia* L. *J. Chem. Ecol.* 1990; 16(9): 2637-46. doi: 10.1007/bf00988075.
18. Eldahshan OA. Isolation and structure elucidation of phenolic compounds of carob leaves grown in Egypt. *Curr. Res. J. Biol. Sci.* 2011; 3(1): 52-5.
19. Güvenalp Z, Özbek H, Ünsalar T, Kazaz C, Demirezer LÖ. Iridoid, flavonoid, and phenylethanoid glycosides from *Wiedemannia orientalis*. *Turk. J. Chem.* 2006; 30(3): 391-400.
20. Im SH, Wang Z, Lim SS, Lee OH and Kang IJ. Bioactivity-guided isolation and identification of anti-adipogenic compounds from *Sanguisorba officinalis*. *Pharm. Biol.* 2017; 55(1): 2057-64. doi: 10.1080/13880209.2017.1357736.
21. Nugroho A, Song BM, Lee KT, Park HJ. Quantification of antidepressant miquelianin in mature and immature fruits of Korean *Rubus species*. *Nat. Prod. Sci.* 2014; 20(4): 258-61. doi: 10.1080/13880209.2017.1357736.
22. Shahidi F and Ambigaipalan P. Phenolics and polyphenolics in foods, beverages and spices: Antioxidant activity and health effects- a review. *J. Funct. Food.* 2015; 18: 820-97. doi: 10.1016/j.jff.2015.06.018.
23. Hyun SK, Jung YJ, Chung HY, Jung HA, Choi JS. Isorhamnetin glycosides with free radical and ONOO⁻ scavenging activities from the stamens of *Nelumbo nucifera*. *Arch. Pharm. Res.* 2006; 29(4): 287-92. doi: 10.1007/bf02968572.
24. El-Aasr M, Kabbash A, El-Seoud KAA, Al-Madboly LA and Ikeda T. Antimicrobial and Immunomodulatory activities of flavonol glycosides isolated from *Atriplex halimus* L. herb. *J. Pharm. Sci. Res.* 2016; 8(10): 1159-68.
25. Zheng Y-Z, Deng G, Liang Q, Chen D-F, Guo R and Lai R-C. Antioxidant activity of quercetin and its glucosides from propolis: A theoretical study. *Sci. Rep.* 2017; 7(1): 1-11. doi: 10.1038/s41598-017-08024-8.
26. Panda S and Kar A. Antidiabetic and antioxidative effects of *Annona squamosa* leaves are possibly mediated through quercetin-3-O-glucoside. *Biofactors* 2007; 31(3-4): 201-10. doi: 10.1002/biof.5520310307.
27. Lee S, Lee J, Lee H and Sung J. Relative protective activities of quercetin, quercetin-3-glucoside, and rutin in alcohol-induced liver injury. *J. Food Biochem.* 2019; 43(11): e13002. doi: 10.1111/jfbc.13002.
28. Süntar IP, Akkol EK, Yalçın FN, Koca U, Keleş H and Yesilada E. Wound healing potential of *Sambucus ebulus* L. leaves and isolation of an active component, quercetin 3-O-glucoside. *J. Ethnopharmacol.* 2010; 129(1): 106-14. doi: 10.1016/j.jep.2010.01.051.
29. Im SH, Wang Z, Lim SS, Lee OH and Kang IJ. Bioactivity-guided isolation and identification of anti-adipogenic compounds from *Sanguisorba officinalis*. *Pharm. Biol.* 2017; 55(1): 2057-64. doi: 10.1080/13880209.2017.1357736.
30. Landa A, Casado R and Calvo MI. Identification and quantification of flavonoids from *Chuquiraga spinosa* (Asteraceae). *Nat.*

- Prod. Commun.* 2009; 4(10): 1353-1355. doi: 10.1177/1934578X0900401008.
- 31.** Petropoulos SA, Fernandes Â, Vasileios A, Ntatsi G, Barros L and Ferreira IC. Chemical composition and antioxidant activity of *Cichorium spinosum* L. leaves in relation to developmental stage. *Food Chem.* 2018; 239: 946-52. doi: 10.1016/j.foodchem.2017.07.043.
- 32.** Parejo I, Jauregui O, Sánchez-Rabaneda F, Viladomat F, Bastida J and Codina C. Separation and characterization of phenolic compounds in fennel (*Foeniculum vulgare*) using liquid chromatography-negative electrospray ionization tandem mass spectrometry. *J. Agric. Food Chem.* 2004; 52(12): 3679-87. doi: 10.1021/jf030813h.
- 33.** Castillo-Muñoz N, Gómez-Alonso S, García-Romero E and Hermosín-Gutiérrez I. Flavonol profiles of *Vitis vinifera* white grape cultivars. *J. Food Compos. Anal.* 2010; 23(7): 699-705. doi: 10.1016/j.jfca.2010.03.017.
- 34.** Del-Castillo-Alonso MÁ, Monforte L, Tomás-Las-Heras R, Martínez-Abaigar J and Núñez-Olivera E. Phenolic characteristics acquired by berry skins of *Vitis vinifera* cv. Tempranillo in response to close-to-ambient solar ultraviolet radiation are mostly reflected in the resulting wines. *J. Sci. Food Agric.* 2020; 100(1): 401-9. doi: 10.1002/jsfa.10068.
- 35.** Badria FA, Ameen M and Akl MR. Evaluation of cytotoxic compounds from *Calligonum comosum* L. growing in Egypt. *Z. Naturforsch. C.* 2007; 62(9-10): 656-60. doi: 10.1515/znc-2007-9-1005.
- 36.** Dall'Acqua S, Cervellati R, Loi MC and Innocenti G. Evaluation of *in vitro* antioxidant properties of some traditional Sardinian medicinal plants: Investigation of the high antioxidant capacity of *Rubus ulmifolius*. *Food Chem.* 2008; 106(2): 745-9. doi: 10.1016/j.foodchem.2007.06.055.
- 37.** Plumb GW, Price KR and Williamson G. Antioxidant properties of flavonol glycosides from green beans. *Redox Rep.* 1999; 4(3): 123-7. doi: 10.1179/135100099101534800.
- 38.** Moon J-H, Tsushida T, Nakahara K and Terao J. Identification of quercetin 3-O- β -D-glucuronide as an antioxidative metabolite in rat plasma after oral administration of quercetin. *Free Radical Bio. Med.* 2001; 30(11): 1274-85. doi: 10.1016/s0891-5849(01)00522-6.
- 39.** Ho L, Ferruzzi MG, Janle EM, Wang J, Gong B, Chen TY, Lobo J, Cooper B, Wu QL, Talcott ST, Percival SS, Simon JE and Pasinetti GM. Identification of brain-targeted bioactive dietary quercetin-3-O-glucuronide as a novel intervention for Alzheimer's disease. *FASEB J.* 2013; 27(2): 769-81. doi: 10.1096/fj.12-212118.
- 40.** Yoon CS, Kim DC, Ko WM, Kim KS, Lee DS, Kim DS, Cho HK, Seo J, Kim SY, Oh H and Kim YC. Anti-neuroinflammatory effects of quercetin-3-O-glucuronide isolated from the leaf of *Vitis labruscana* on LPS-induced neuroinflammation in BV2 cells. *Korean J. Pharmacog.* 2014; 45(1): 17-22.
- 41.** Ohara K, Wakabayashi H, Taniguchi Y, Shindo K, Yajima H and Yoshida A. Quercetin-3-O-glucuronide induces ABCA1 expression by LXR α activation in murine macrophages. *Biochem. Bioph. Res. Co.* 2013; 441(4): 929-34. doi: 10.1016/j.bbrc.2013.10.168.
- 42.** Mahmoudi R, Kosari M, Zare P and Barati S. *Kelussia odoratissima* essential oil: biochemical analysis and antibacterial properties against pathogenic and probiotic bacteria. *J. Agroaliment. Processes Technol.* 2014; 20(1): 109-15. doi: 10.26226/morressier.5f4ccc57648d3d8b3f362be9.
- 43.** Raeisi S, Mirjalili MH, Nadjafi F and Hadian J. Variability in the essential oil content and composition in different plant organs of *Kelussia odoratissima* Mozaff. (Apiaceae) growing wild

in Iran. *J. Essent. Oil Res.* 2015; 27(4): 283-8. doi: 10.1080/10412905.2015.1025917.

44. Esmaeili A, Sharafian S, Mirian S and Larijani K. Chemical composition of essential oil from leaves, stems and flowers of *Kelussia odoratissima* Mozaff. grown in Iran. *J. Essent. Oil-Bear. Plants.* 2011; 14(5): 643-6. doi: 10.1080/0972060x.2011.10643984.

45. Omidbaigi R, Sefidkon F and Saeedi K. Essential oil content and composition of *Kelussia odoratissima* Mozaff. as an Iranian endemic plant. *J. Essent. Oil-Bear. Plants.* 2008; 11(6): 594-7. doi: 10.1080/0972060X.2008.10643672.

46. Shojaei ZA, Ebrahimi A and Salimi M. Chemical composition of three ecotypes of wild celery (*Kelussia odoratissima*). *J. Herbs Spices Med. Plants.* 2011; 17(1): 62-8. doi: 10.1080/10496475.2011.560089.


47. Pirbalouti AG, Sedaghat L, Hamed B and Tirgir F. Chemical composition and antioxidant activity of essential oils of three endemic

medicinal plants of Iran. *Bangladesh J. Bot.* 2013; 42(2): 327-32. doi: 10.3329/bjb.v42i2.18038.

48. Baser KHC and Buchbauer G. Handbook of essential oils: science, technology, and applications: CRC press; 2015.

49. Xie Q, Zhang L, Xie L, Zheng Y, Liu K, Tang H, Liao Y and Li X. Z-Ligustilide: a review of its pharmacokinetics and pharmacology. *Phytotherapy Res.* 2020; 34(8): 1966-1991. doi: 10.1002/ptr.6662.

How to cite this article: Khanavi M, Ghadami Sh, Sadaghiani-Tabrizi Gh, Delnavazi MR. Phytochemical constituents of the fruits of *Kelussia odoratissima* Mozaff., an aromatic plant endemic to Iran. *Journal of Medicinal Plants* 2021; 20(79): 1-13. doi: 10.52547/jmp.20.79.1



ترکیبات فیتوشیمیایی میوه کلوس (*Kelussia odoratissima* Mozaff.)، گیاهی معطر و انحصاری ایران

مهناز خانوی^۱، شیما قدمی^۱، غزل صدقیانی تبریزی^۱، محمدرضا دلنوازی^{۱*}

^۱ گروه فارماکوتکنولوژی، دانشکده داروسازی، دانشگاه علوم پزشکی تهران، تهران، ایران

^۲ دانشکده سیستم‌های زمین و غذا، دانشگاه بریتیش کلمبیا، ونکوور، BC، کانادا

چکیده

اطلاعات مقاله

مقدمه: گیاه کلوس (*Kelussia odoratissima* Mozaff.) از تیره چتریان، گیاهی است علفی و پایا که به صورت انحصاری در غرب ایران می‌روید. مردم بومی از بخش‌های هوایی معطر این گیاه به صورت سنتی به عنوان عطر و طعم‌دهنده در تهیه بعضی غذاهای محلی و همچنین در درمان بیماری‌های مختلف استفاده می‌کنند. **هدف:** مطالعه حاضر با هدف جداسازی و شناسایی ترکیبات فنولی و همچنین آنالیز ترکیبات اسانس میوه‌های گیاه کلوس طراحی شده است. **روش بررسی:** ترکیبات فراکسیون بوتانولی نرمال بدست آمده از عصاره هیدروالکلی میوه گیاه با استفاده از کروماتوگرافی روی ستون‌های فاز نرمال و سفادکس ال اچ-۲۰ مورد جداسازی و خالص‌سازی قرار گرفت. ترکیبات اسانس بدست آمده از نمونه نیز با استفاده از تکنیک کروماتوگرافی گازی متصل به طیف‌سنج جرمی آنالیز شد. **نتایج:** مطالعه فیتوشیمیایی روی میوه‌های گیاه کلوس به جداسازی و شناسایی پنج گلیکوزید فلاونولی با نام‌های ایزورامنتین ۳-ا^۱، گلوکوزید (۱)، کوئرستین ۳-ا^۱، گلوکوزید (ایزوکوئرستین) (۲)، ایزورامنتین ۳-ا^۱، روتینوزید (نارسیسین) (۳)، ایزورامنتین ۳-ا^۱، گلوکورونید (۴) و کوئرستین ۳-ا^۱، گلوکورونید (مکوئیلینانین) (۵) انجامید. تعداد سی و شش ترکیب نیز در نتیجه آنالیز اسانس حاصل از میوه‌های گیاه شناسایی شد که از این میان ترکیبات (Z) - لیگوستیلید (۱۵/۹۳ درصد)، دلتا- کادینن (۱۲/۲۶ درصد) و جرماکرن دی (۱۲/۱۸ درصد) ترکیبات عمده بودند. **نتیجه‌گیری:** نتایج این مطالعه، میوه‌های گیاه کلوس را به عنوان منبعی از گلیکوزیدهای فلاونوئیدی و مشتقات فتالات معرفی می‌کند. حضور این ترکیبات که خواص بیولوژیک و اثرات سودمند آنها بر سلامتی در مطالعات پیشین نشان داده شده است، میوه‌های این گیاه را به عنوان گزینه مناسبی برای مطالعات مرتبط بیشتر مطرح می‌کند.

گل‌واژگان:

کلوس

تیره چتریان

فلاونوئید

اسانس

(Z) - لیگوستیلید

مخفف‌ها: NMR، رزونانس مغناطیسی هسته‌ای؛ GC-MS، کروماتوگرافی گازی متصل به طیف‌سنج جرمی؛ TLC، کروماتوگرافی لایه نازک

* نویسنده مسؤول: delnavazi@tums.ac.ir

تاریخ دریافت: ۳۰ بهمن ۱۳۹۹؛ تاریخ دریافت اصلاحات: ۲۹ خرداد ۱۴۰۰؛ تاریخ پذیرش: ۱۲ تیر ۱۴۰۰

doi: 10.52547/jmp.20.79.1

© 2020. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>)