

Journal of Medicinal Plants

Journal homepage: www.jmp.ir



Research Article

Design, reformulation, and standardization of a traditional-based memory enhancer herbal preparation originated from Persian medicine

Farid Dabaghian^{1,2}, Amir Azadi³, Mohammad Mehdi Zarshenas^{1,4,*}

¹ Medicinal Plants Processing Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

² Department of Pharmacognosy, School of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

³ Department of Pharmaceutics, School of Pharmacy, Shiraz University of Medical Sciences, Shiraz, Iran

⁴ Department of Phytopharmaceuticals (Traditional Pharmacy), School of Pharmacy, Shiraz University of Medical Sciences, Shiraz, Iran

ARTICLE INFO	ABSTRACT
Keywords:	Background: Alzheimer's disease (AD) is the most common type of dementia that has
Alzheimer's disease	been cited in traditional Persian medicine (TPM) manuscripts under the heading
Tablet	"Farāmūshkārī", "Fesād-e-Zekr" and "Nesiān". Since AD is a significant challenge for
Bunium persicum	health professionals, searching for natural and effective compounds is crucial.
Piper nigrum	Objective: Given the importance of AD, this study has provided a standardized plant-
Standardization	based tablet based on TPM. Methods: Related preparations were extracted from leading
I raditional medicine	pharmaceutical manuscripts of TPM. Out of 127 gathered compound formulations, a
	preparation containing Bunium persicum (Boiss.) B.Fedtsch. (Persian cumin) and Piper
	nigrum L. (black pepper) was selected from the Canon of Medicine of Avicenna. After
	evaluating various excipients for tablet preparation, the proportions of Persian cumin (40
	%), black pepper (40 %), hydroxypropyl methylcellulose (HPMC), and
	polyvinylpyrrolidone (PVP) as solution binder (20 %) were chosen. The final
	formulation was undergone through ethanol and dichloromethane and essential oil
	extractions to prepare for HPTLC fingerprints and GC assessments, respectively. The
	total alkaloid in the tablet's ethanol extract was also determined via spectrophotometry.
	Results: The final tablet was standardized via GC based on trans-caryophyllene (29.22)
	%) and γ -terpinene (17.72 %). Each tablet contains an average of 0.041 mg of trans-
	caryophyllene and 0.018 mg of 7-terpinene. The amount of the total alkaloid in each
	tablet was determined 0.06 mg, equivalent to atropine. Conclusion: This prepared
	simple natural formulation can be considered a traditional-based product to be evaluated
	clinically and presented as a memory enhancer supplement.

Abbreviations: AD, Alzheimer's disease; TPM, Traditional Persian medicine; HPMC, Hydroxypropyl Methylcellulose; PVP, Polyvinylpyrrolidone; GC, Gas chromatography; MS, Mass spectrometry; FID, Flame ionization detector; HPTLC, High-performance thin-layer chromatography

^{*}Corresponding author: <u>zarm@sums.ac.ir</u>

doi: 10.52547/jmp.21.82.93

Received 28 March 2022; Received in revised form 27 May 2022; Accepted 27 May 2022

^{© 2020.} Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<u>https://creativecommons.org/licenses/by-nc/4.0/</u>)

1. Introduction

According to the International Working Group (IWG), Alzheimer's disease (AD) is now recognized in vivo as a dual clinico-biological entity that can be identified prior to the onset of the dementia syndrome, on the basis of: i) a distinct core clinical phenotype characterized by hippocampal amnestic syndrome, and ii) changing of biomarkers that reflect the location or nature of Alzheimer's-type [1]. AD, the most common form of dementia, begins gradually and progressively disrupts the brain's cognitive functions [2]. About 35.6 million people worldwide have dementia, which is expected to double every 20 years, with an estimated 115.4 million people worldwide by 2050 [3]. The disease, which mainly affects the elderly [4], is flagrantly becoming a widespread problem as the population ages increase worldwide. AD is currently imposing a vast economic and social burden on societies. Economically, it is estimated that dementia accounted for approximately 604 billion dollars worldwide in 2010, indicating that the disease can cause many economic problems for communities, especially low-income groups [5].

There are still many unknowns about the pathophysiology of AD. Several investigations have been carried out into influential factors including acetylcholine deficiency, Tau protein hyperphosphorylation, and oxidative stress- used to develop new medicines to treat this disorder [6]. Loss of memory and severity of AD is closely linked to cholinergic activity decline in certain parts of the brain [7]. According to the cholinergic hypothesis, medicines with acetylcholinesterase enzyme inhibitory activity might effectively manage AD [8]. Recently, known medicines such Donepezil, as Rivastigmine, Ladostigil, and Galantamine have been used as acetylcholinesterase inhibitors [9].

Much evidence also implies the role of oxidative stress and free radicals in the pathogenesis and possible cause of this disease [10]. Correspondingly, the use of antioxidant therapies in the pre-clinical stages has been successful [11].

However, no drug that can completely prevent AD has been developed. As a result, many pharmaceutical companies are developing and producing novel medications to help slow the progression of AD [12, 13].

Since AD is a significant challenge for health professionals, searching for natural and effective compounds is crucial. Different traditional medical systems may offer good opportunities for treating AD patients [14]. Traditional Persian medicine (TPM) offers various therapies to manage such chronic diseases [15]. Persian practitioners were well aware of amnesia, dementia, and Alzheimer's disease (Nesiān and Fesād-e-Zekr) [16]. They provided a variety of medicinal plants to help with memory loss, including Piper nigrum, Cinnamomum verum, Zingiber officinale, Boswellia spp., Terminalia chebula, Bunium persicum, Acorus calamus, Allium sativum and etc [17]. Animal studies and clinical trials have confirmed the biological activity of some of these; for example, when an aqueous extract of Boswellia serrata was administered orally to AD-induced rats, there was an increase in Ach levels and a decrease in AchE activity in brain homogenates [18] or the combination of Piper nigrum and Cinnamomum significantly zeylanicum increased the recognition index in mice with scopolamineinduced memory impairment [19]. In another study the ginger was given orally to volunteers to evaluate cognitive task and working memory of middle-aged women, and its use significantly improved digit vigilance, word recognition, choice reaction, spatial working memory, and

Journal of Medicinal Plants

94

numeric working memory scores [20]. As a result, these medicinal plants can assist us in developing effective medications to prevent or slow the progression of AD; but, in order to improve patient compliance, modern medicine dosage forms should be used instead of traditional medicine dosage forms such as *habb*, *majoon*, *yaqooti*, *itrifal* and etc. Some of these are produced in new dosage forms, such as a biocompatible active film of nanochitosan containing free and nanoliposome caraway seed extract or gastroretentive matrix tablets of Boswellia Oleogum Resin [21, 22].

The purpose of this study is to design and reformulate one of Aviccena's traditional compounding formulations (containing Persian cumin and black pepper) as a memory enhancer from powder to tablet dosage form in order to improve patient compliance and satisfaction, and to standardize the new dosage form based on the major component of essential oil and total alkaloid in order to improve the accuracy of the amount of medicine used.

2. Materials and Methods

2.1 Materials

The required materials used in this study are listed as follows: ethanol. methanol, chloroform. dichloromethane toluene, Bromocresol Green (BCG) sodium and hydroxide (NaOH) were purchased from Merck Co. (Germany); all these analytical grade compounds. The following materials manufactured by Sigma-Aldrich Co. (Germany) have also been used: HPMC, PVP, starch, acacia and avicel.

2.2. Manuscripts review

A line of the traditional pharmaceutical encyclopedias (*Qarābādins*) in TPM, including

Qarābādin-e-Kabir [23], Qarābādin-e-Salehi [24], Qarābādins-e-Ghāderi [25], Qarābādin-e-Azam [26], Tohfah al-Momenin (Rarity of the Faithful) [27] and the third volume of Avicenna's Canon of Medicine [28], were studied with the exact keywords; "Nesiān", "Farāmūshkārī", "Fesād-e-Zekr". Memory enhancer formulations were collected and cited [29]. From the third volume of the Canon of Medicine, a twoingredient formulation with repeated citations in those encyclopedias was chosen to enter the experimental phases. This preparation contained two main components: Piper nigrum L. (Black pepper) fruits and Bunium persicum (Boiss.) B.Fedtsch. (Persian Cumin) seeds. The daily dose to prevent memory loss was recommended at 1.7 g.

2.3. Plants authentication and microscopic characterization

The purchased samples of *Piper nigrum* L. and *Bunium persicum* (Boiss.) B.Fedtsch. were authenticated and deposited with a specific voucher number in the herbarium of the School of Pharmacy of Shiraz University of medical sciences. The vouchers were PM 1008 for Persian cumin and PM 862 for black pepper. Pepper and cumin samples were ground to determine botanical organs using an electron microscope and a digital camera.

2.4. Tablet preparation and Pharmaceutical tests

Using several different excipients, those herbal components (milled, and sieved previously) were mixed (1:1 ratio) to prepare the initial mixture of the tablet. Also, solution binders were used to improve the pharmaceutical properties of the dosage form. For this purpose, Avicel, Starch, Acacia, HPMC, and PVP were used as solution binders in a concentration of 24 % W/V in different proportions. Various pharmaceutical tests, including the angle of repose [30] and Hausner ratio for assessment of final powder characteristics and also various tablet tests such as hardness, weight variation, friability, and disintegration time, were performed according to British Pharmacopoeia [31].

2.5. Thin Layer Chromatography (TLC) and High-Performance Thin Layer Chromatography (HPTLC) fingerprinting

TLC technique was employed to seek for the optimized solvent system for HPTLC. Two types of extracts including dichloromethane and ethanol, were prepared by using ultrasonic bath technique from each component and the final formulation to determine the TLC profile. In this method, 10.0 g of each powder was poured into dichloromethane and then subjected to an ultrasonic bath for 15 minutes three times; the plant residues were then used for ethanol extraction, resulting in the extraction of polar and non-polar compounds separately. The extracts were then dissolved in their respective solvents (1:10) before being subjected to the TLC test. To determine the best solvent system for the TLC assessment based on the ability to distinguish different components of extracts, various solvents such as chloroform, ethanol, methanol, toluene, dichloromethane, and water were investigated. Chromatographic plates were examined in visible light first, followed by UV light at 254 nm and 366 nm, respectively. Anisaldehyde reagent was then applied to the plates. The stains reappeared after drying, and the plate was examined under visible and UV light again. Also, HPTLC was used to determine the final profile as a fingerprint, using an appropriate solvent system for each extract.

2.6. Essential oil preparation and Gas chromatography

Approximately, 50.0 g of Persian cumin, black pepper, and the final tablet were ground into powder and soaked in 500.0 ml of distilled water (1:10 w/v) separately to extract the essential oil. Hydrodistillation was carried out for 4 hours using Clevenger-type apparatus, yielding essential oils that were stored in the dark at 4 °C until analysis.

Essential oil components were analyzed using technologies model 7890A Agilent gas chromatograph (column; HP-5MS, 25 mm, 30 mm) attached to a US Agilent technology mass spectrometer (MS). Helium was selected as the carrier gas with a 1 ml/min flow rate with a split ratio of 1:100. The injector temperature was 250.0 °C, while column temperature was linearly programmed from 60.0 to 220.0 °C (rate of 5 °C/min) and then held for 10 min at 220.0 °C. The applied voltage was 70 EV, and its interface temperature was set at 280.0 °C. The mass range was set at 30-600 m/z. Then, each sample was injected into the device, and the final compound profile was analyzed.

Identification of essential oil components was accomplished based on the comparison of their retention times with those of authentic standards and by comparison of their mass spectral fragmentation patterns (NIST Chemistry WebBook and the Adams' textbook [32]).

2.7. Gas chromatography coupled to flame ionization detector (GC/FID)

Following GC/MS analysis of the compounds in Persian cumin, black pepper, and final formulation, Agilent technology US 7890A GC/FID was used to standardize the product

Journal of Medicinal Plants

96

Downloaded from jmp.ir on 2025-07-12

based on major components (transcaryophyllene and *y*-terpinene). For this purpose, nitrogen gas with 1 ml/min flow was selected as the carrier gas. Since the device had an FID detector, it needed two hydrogen and oxygen gases that gave it the best flame at the ratio of 1 to 10. We used split mode $(1/_{50})$, and 1 µl of the sample was injected into the device. The injection chamber was set to 270 °C, and the column used was HP-5 (30 m \times 320 mm). The stationary phase was phenylmethyl siloxane with a diameter of 0.25 µm. The column temperature was set at 60 °C. The sample was kept at this temperature for 2 minutes, then increased at a rate of 6 °C/min until it reached 250 °C, where it was held for 10 minutes. The temperature of the detector was also set at 300 °C.

As standards, trans-caryophyllene and y-terpinene (Analytical Standard grade, Sigma Aldrich) were used. By sequential dilution with n-hexane, different concentrations of transcaryophyllene (0.18, 0.36, 0.9, 1.8, and 3.6 mg/ml) and *y*-terpinene (0.85, 1.7, 8.5, 17, and 42.5 mg/ml) were prepared. Three injections were performed for each standard to check the reproducibility, and the standard curve was plotted by the mean of three injections for each standard. The mean of three injections was used as the sample concentration for the essential oil in the final formulation. In addition, to validate the method, a concentration of 8.5 mg/ml was injected into the device three times daily for three different days to calculate intra- and inter-day differences as well as relative standard deviation (RSD). The limit of detection and the limit of quantification (LOQ) were measured separately for each marker.

2.8. Determination of total alkaloid content in the final tablet

F. Dabaghian, et al

Dried ethanol extract of the final formulation was dissolved in hydrochloric acid (2 N) and filtered subsequently. Afterward, 1 ml of the solution was transferred to a funnel and washed three times with chloroform (10 ml every time). Enough sodium hydroxide was added to the residual solution to neutralize the solution. Finally, 5 ml of BCG, 5 ml of phosphate buffer, and separately, 1, 2, 3 and 4 ml of chloroform were added to the neutralized solution. respectively. The chloroform phases were collected, and the volume was increased to 10 ml with chloroform [33]. This procedure was also used for atropine as a standard alkaloid, and absorption at various concentrations (40, 80, 100, and 120 mg/ml) was measured using a spectrophotometer at 470 nm, and a calibration curve was established. Finally, the absorbance of the resulting solution of the sample was measured at 470 nm, and the total alkaloid content of the formulation was calculated.

3. Result

3.1. Traditional literature review

In total, 127 prescriptions were extracted from the aforementioned traditional encyclopedias. During this search, 3442 pages of traditional texts were studied. Finally, a medicinal prescription mentioned in Canon of Medicine, consisting of Persian cumin and black pepper, was chosen for subsequent processes.

3.2. Microscopic Characterization

An electron microscope was used to examine the fruit of Persian cumin and black pepper. Specific characteristics of each fruit limb were determined in Fig. 1.

DOI: 10.52547/jmp.21.82.93]





в





Fig. 1. A: Microscopic characterization of Persian cumin (in chloral hydrate): a) Endocarp b) Epicarp c) Sclereid of mesocarp d) Cells of inner fruit with duct. B: Microscopic characterization of black pepper (in chloral hydrate): a) Exocarp b) Sclereid of endocarp c) Stone cells d) Pitted cells of mesocarp

3.3. Assessment of different tablet formulations

Following the use of various excipients and methods to create the tablet from the selected prescription, each formulation was tested for hardness, weight, and disintegration time. Among 54 different formulations, a tablet containing 40 % Persian cumin, 40 % black pepper, 10 % HPMC, and 10 % PVP as solution binder had good characteristics and was selected for more specific pharmaceutical controls (Fig. 2). The evaluation of powder flow by measuring the angle of repose revealed that the

Journal of Medicinal Plants

flow improved after using HPMC and PVP solution binders from 42.56° to 33.55°.

The mean weight of the 20 tablets was 0.401 g, and the percentage of weight changes was calculated as 1.412 %. Furthermore, the measured thickness of 10 tablets was 3.02 ± 0.042 mm, the friability was less than 1 %, the

hardness was 4.87 ± 0.24 kilogram-force and the final formulation tablets were disintegrated in 24 minutes. The final tablet contained an average of 0.041 mg *trans*-caryophyllene, 0.018 mg of γ -terpinene, and 0.06 mg total alkaloid (equivalent to atropine).



Fig. 2. Final prepared tablets



Fig. 3. High-performance thin-layer chromatogram (A: Dichloromethane extract, B: Ethanol extract) under UV light at 366 nm before adding reagent (right), at 254 nm before adding reagent (middle), under visible light after adding reagent (Left side). Sample arrangement from left: a: Black pepper, b: Final formulation, c: Persian cumin, d: Piperine

Journal of Medicinal Plants

3.4. Results of pharmacognostic evaluations

TLC, HPTLC, GC/MS, GC/FID, and total alkaloid content were all determined through various analyses.

3.4.1. HPTLC Analysis of Extract

As previously stated, a concentration of 3.0 mg/ml was prepared for various extracts of Persian cumin, black pepper, and the final tablets. Finally, toluene and ethyl acetate ($\underline{80}$:20) was used for ($\underline{70}$:15:5) was selected for ethanol extracts. HPTLC chromatogram for dichloromethane and ethanol extracts is shown in Fig. 3.

3.4.2. GC/MS analysis of the essential oil samples

The yields of the essential oils of Persian cumin, black pepper, and final formulations were 2.8 %, 1 %, and 3.4 %, respectively. *trans*-Caryophyllene and γ -terpinene were identified as major compounds after determining more than 91 % of the essential oil components of the finalformulation. The chemical constituents of the essential oil samples are shown in Table 1.

 Table 1. Essential oil analysis of compounds in Persian cumin (PC), Black Pepper (BP) and Final formulation (F) by

 GC/MS

PC	BP	F	KI _{CAL} *	KI _{ST} **
0.63	0.7	2.01	935	939
0.66	-	1.55	975	975
1.34	1.95	3.69	980	979
1.04	0.83	1.67	991	991
-	1.68	2.11	1008	1003
-	5.98	7.17	1015	1015
6.93	-	-	1030	1026
22.73	7.15	-	1033	1029
0.43	-	0.34	1048	1050
26.8	-	17.72	1064	1060
0.3	0.74	0.76	1091	1089
-	0.75	0.72	1102	1097
0.66	-	0.41	1180	1177
1.23	-	0.81	1197	1207
2.08	-	1.04	1201	1193
1.15	-	0.56	1208	1217
2.65	-	1.04	1244	1244
13.11	-	-	1258	1243
0.39	-	-	1271	1268
	PC 0.63 0.66 1.34 1.04 - - 6.93 22.73 0.43 26.8 0.3 - 0.66 1.23 2.08 1.15 2.65 13.11 0.39	PCBP 0.63 0.7 0.66 - 1.34 1.95 1.04 0.83 - 1.68 - 5.98 6.93 - 22.73 7.15 0.43 - 26.8 - 0.3 0.74 - 0.75 0.66 - 1.23 - 2.08 - 1.15 - 2.65 - 13.11 - 0.39 -	PCBPF 0.63 0.7 2.01 0.66 - 1.55 1.34 1.95 3.69 1.04 0.83 1.67 - 1.68 2.11 - 5.98 7.17 6.93 22.73 7.15 - 0.43 - 0.34 26.8 - 17.72 0.3 0.74 0.76 - 0.75 0.72 0.66 - 0.41 1.23 - 0.81 2.08 - 1.04 1.15 - 0.56 2.65 - 1.04 13.11 0.39	PCBPFKIcat* 0.63 0.7 2.01 935 0.66 - 1.55 975 1.34 1.95 3.69 980 1.04 0.83 1.67 991 - 1.68 2.11 1008 - 5.98 7.17 1015 6.93 1030 22.73 7.15 - 1033 0.43 - 0.34 1048 26.8 - 17.721064 0.3 0.74 0.76 1091 - 0.75 0.72 1102 0.66 - 0.41 1180 1.23 - 0.81 1197 2.08 - 1.04 1201 1.15 - 0.56 1208 2.65 - 1.04 1244 13.11 1258 0.39 1271

Journal of Medicinal Plants

DOR: 20.1001.1.2717204.2022.21.82.5.3]

[Downloaded from jmp.ir on 2025-07-12]

Table 1. Essential oil analysis of compounds in Persian cumin (PC), Black Pepper (BP) and Final formulation (F) by
GC/MS (Continued)

Components	РС	BP	F	KI _{CAL} *	KIst**
Perilla aldehyde	0.43	-	-	1280	1272
Unknown	5.48	-	2.45	-	-
Unknown	-	-	5.9	-	-
Unknown	7.95	-	-	-	-
1,4-p-Menthadien-7-ol	0.65	-	-	1332	1332
δ -Elemene	-	7.48	4.68	1344	1338
α-Copaene	-	6.26	2.81	1383	1377
β -Elemene	-	1.86	0.39	1397	1391
Cuminyl acetate	0.88	-	-	1425	1432
(E)-Caryophyllene	-	51	29.22	1439	1419
α-Humulene	-	4.22	2.46	1464	1455
Germacrene D	0.29	0.54	0.63	1486	1485
β -Selinene	-	1.24	0.95	1494	1490
α-Selinene	-	1.21	0.91	1502	1498
β -Bisabolene	-	0.59	0.62	1512	1506
δ -Cadinene	-	2.86	1.52	1530	1523
Caryophyllene oxide	-	1.03	0.85	1591	1583
Dill apiole	1.89	-	2.65	1630	1621
(-)-Spathulenol	-	1.89	1.51	1637	1619
Palmitic acid	-	-	0.86	1966	1963
Identification	86.63	99.96	91.66	-	

* **KI**_{CAL}: Calculated kovats index.

**** KI**ST: Standard kovats index.

3.4.3. GC/FID-based determination of transcaryophyllene and γ -terpinene in the final formulation

According to GC/MS data analysis, the main volatile compound in the essential oil of the final formulation was *trans*-caryophyllene, followed by γ -terpinene. A stock solution and subsequent serial dilutions of *trans*-caryophyllene (0.18,

0.36, 0.9, 1.8, and 3.6 mg/ml) and γ -terpinene (0.85, 1.7, 5 8, 17, 42.5 mg/ml) were prepared to draw calibration curves with correlation coefficients of 0.9996 and 0.9962, respectively. All data related to the determination of markers are presented in tables 2 to 4. Also, respective calibration curves are shown in Fig. 4 and Fig. 5.

Journal of Medicinal Plants

101

[DOR: 20.1001.1.2717204.2022.21.82.5.3]

3.4.4. Determination of total alkaloid

The final formulation was standardized based on the amount of total alkaloid equivalent to atropine in each tablet, in addition to the major compounds in the essential oil. The absorption of atropine at different concentrations (40.0, 80.0, 100.0 and 120.0 mg/ml) were 0.031, 0.048, 0.056, and 0.063, respectively. The calibration curve was drawn by this data and the total alkaloid equivalent to atropine for ethanol extract of final formulation was calculated. The extract concentration of 12.5 mg/ml resulted in absorbances of 0.049, 0.049, and 0.048 in the first, second, and third replicates, respectively. The absorbances were entered into the following equation:

 $Abs = 0.0004 \times Concentration + 0.01512$

The alkaloid content was calculated to be 664.906 ± 11.44 mg in 100.0 g of extract.

Table 2. <i>trans</i> -Caryophynelle and <i>p</i> -terpinelle calibration data

Compound	Concentration(mg/ml)	Sample number	Area	Mean± SD	RSD
trans-Caryophyllene	0.18	1 2 3	3190.5 2798.7 3473.5	3272.7 ± 174.8	5.3
	0.36	1 2 3	6512.7 6826.7 6446.6	6595.3 ± 203.7	3
	0.9	1 2 3	13046.8 13181.4 13219.8	13149.3 ± 90.84	0.69
	1.8	1 2 3	27561.1 23985 24762	25436.03 ± 1880.92	7.3
	3.6	1 2 3	53811.9 51453.2 46655.7	50460.2 ± 3646.7	7.2
	0.85	1 2 3	12564 10920 14345	12609 ± 1712.95	13.58
	1.7	1 2 3	34645 30241 30584	31823.3 ± 2449.64	7.7
	8.5	1 2 3	147534 145819 138750	144034 ± 4656.01	3.23
	17	1 2 3	280561 268687 237145	262131 ± 22438.2	8.6
	42.5	1 2 3	494772 532713 502671	510085 ± 20018.47	3.9

C1	Days -	Area			Mean ± SD	RSD %	RSD %
Compound		A_1	A_2	A 3	(intra-day)	(intra-day)	(inter-day)
<i>trans-</i> Caryophyllene	1	3154.2	3297.66	2969.06	3140.3 ± 164.7	5.2	
	2	3195.6	3181.7	2865.72	3081 ± 186.5	6.02	2.29
	3	3203.3	3198	3273.04	3224.7 ± 41.8	1.29	-
≁Terpinene -	1	154413	154880	147534	152275.7 ± 4113.04	2.7	
	2	159115.5	162315	145819	155749.83 ± 8747.9	5.62	5.6
	3	144034	140119	138750	140967.7 ± 2742.32	1.94	-

Table 3. Intra- and inter-day differences of *trans*-caryophyllene and γ -terpinene

Table 4. *trans*-Caryophyllene and γ -terpinene content in the essential oil of the final formulation

Compound	Injection No.	Area	Concentration (mg/ml)	Mean ± SD	RSD %	LOD (mg/ml)	LOQ (mg/ml)
trans- Caryophyllene	1	41442.1	2.94				
	2	42692.4	3.03	3.04±0.12	3	0.06	0.18
	3	44615.8	3.17	-			
<i>γ</i> -Terpinene	1	23439.6	1.429				
	2	22937.3	1.339	1.359±0.06	4.57	2.6* 10-4	8*10-4
	3	24955	1.309	-			



Fig. 4. Standard curve of trans-caryophyllene

[DOI: 10.52547/jmp.21.82.93]

Journal of Medicinal Plants

103



Fig. 5. Standard curve of γ -terpinene

4. Discussion

The ancient sages are thought to have established the foundation for modern medicine, and their instructions can be used to discover and design new pharmaceutical dosage forms and preparations. This valuable information could be used to either treat diseases directly or to provide good sources of various natural medicines [34, 35].

Natural and traditional medicines are now used to prevent and treat a wide range of diseases, including cancers diabetes [36], [37]. gastrointestinal and neurological disorders [38], inflammatory issues [39], and bacterial and viral infections [40, 41]. Furthermore, the potential of traditional medicine in treating diseases like Alzheimer's, for which there is no complete cure, can be significant. However, due to a lack of standardization or quality controls, there are numerous concerns and pitfalls associated with the use of traditional medicines [42, 43]. As a result, these medicines must be designed and developed based on new dosage forms with standardized doses in order to improve drug quality and patient satisfaction.

The present study selected a simple traditional formulation reported as a memory enhancer from the list of memory enhancer preparations cited in TPM manuscripts. This formulation includes Persian cumin and black pepper.

Persian cumin has been used in folk medicine to treat digestive issues such as bloating and stomach pain [44]. There have also been reports of anticonvulsant [45], anti-anxiety, antidepressant [46], and anti-inflammatory effects [47]. Furthermore, this herbal remedy may inhibit acetylcholinesterase enzyme, the potentially aiding in the treatment of AD [48]. The effects of caraway extract on nootropic activities in normal and stress-induced rats were studied, and it was discovered that it has significant anti-stress activity and reduces memory loss due to its central cholinomimetic effect as well as its free radical scavenging mechanism [49].

In addition, black pepper is used as an antiflatulent, appetizer, and anti-inflammatory in medicine. It has nutritional value in addition to being a spicy condiment [50]. Various studies on the pharmacological effects of black pepper

Journal of Medicinal Plants

104

Design, reformulation ...

show that it can control and manage Alzheimer's disease through a variety of underlying mechanisms [51, 52]. Piperine, one of the bioactive compounds found in black pepper, has been shown in adult male Wistar rats at all doses to reduce memory impairment and nerve damage to the hippocampus. This role could be explained by a decrease in lipid peroxidation and an inhibitory effect on the acetylcholinesterase enzyme [53]. Furthermore, in numerous studies, this bioactive compound has been shown to act as an antioxidant and to eliminate the effects of acetyridine mustard acetylcholine ion brain destruction in an animal model of AD [54, 55]. Moreover, the use of methanol extract of black pepper in rat Alzheimer's disease models was shown to improve memory function and performance via antioxidant activities [56]. As a result, black pepper may play an important role in reducing brain neuron damage in the treatment of Alzheimer's disease [57].

Another study showed that oral consumption of β -caryophyllene reduces neuroinflammation by activating cannabinoid two receptors as well as the peroxisome proliferator-activated receptor- γ (PPAR γ) pathway in mice with Alzheimer's disease [58]. Also, this compound can play a role as one of the effective compounds in the treatment of Alzheimer's patients by reducing the load of β-amyloid, Nitric Oxide Prostaglandin (NO), and E2 (PGE2), Cyclooxygenase-2 (COX-2) and by acting on pro-inflammatory factors [59].

Standardization of herbal products is a critical step after formulation. In Traditional Chinese Medicine (TCM), for example, standardization is carried out in accordance with the principles of "unification, simplification, coordination, and optimization" to ensure TCM quality and safety, to promote TCM modernization and international communication in order to achieve good economic and social outcomes [60]. Standardization of herbal products ensures their quality, purity, safety, and efficacy [61].

According to the traditional use of Persian cumin and black pepper, as well as the mentioned related activities in modern medicine, these two components may have a synergistic effect, and act as a good supplement to prevent AD. Following formulation design, control, and standardization of the traditional prescription, a new tablet formulation containing 40 % Persian cumin, 40 % black pepper, 10 % HPMC, and 10 % PVP was established.

To investigate the phytochemical information of herbal medicines, the chromatographic fingerprint is used as a comprehensive identification method [62]. HPTLC was used to determine the presence of piperine in the final formulation. Because the components in each plant can vary depending on external conditions, reproducing the results without conforming to the analytical data is impossible. As a result, measurements quantitative on the final formulation were performed. Overall, the current study has created a new simple tablet formulation based on traditional Persian pharmaceutical manuscripts with memory enhancing effects. Analytical outcomes of this investigation include constituent and final product fingerprints, including GC/MS and HPTLC data, content determination of the final formulation via GC/FID, and spectrophotometry. In this study, each tablet contained an average of 0.041 mg trans-caryophyllene, 0.018 mg of y-terpinene, and 0.06 mg of the total alkaloid (equivalent to atropine). The produced medicine can be clinically evaluated and considered a simple memory enhancer natural formulation.

Journal of Medicinal Plants

105

5. Conclusion

This work has designed, developed, and standardized a herbal formulation containing Persian cumin and black pepper based on traditional Persian medicine reports. This research can be used not only for the production and development of a new supplement, but also as a process for the preparation of herbal products in the form of solid dosage form while controlling and standardizing methods are taken into account. This new tablet has been standardized based on trans-caryophyllene, γ terpinene and total alkaloid content. This new dosage form can ensure the quality and safety of this herbal product while also increasing patient compliance. The final formulation has introduced a memory enhancer preparation that can be evaluated clinically in various related groups of participants. With an adjusted low dose of administration and safe side for unwanted

References

1. Dubois B, Epelbaum S, Santos A, Di Stefano F, Julian A, Michon A, Sarazin M and Hampel H. Alzheimer disease: from biomarkers to diagnosis. *Rev. Neurol.* 2013; 169(10): 744-751. doi: 10.1016/j.neurol.2013.07.016.

2. Alzheimer's Association, Thies W, Bleiler L. 2013 Alzheimer's disease facts and figures. *Alzheimers Dement*. 2013; 9(2): 208-245. doi: 10.1016/j.jalz.2013.02.003.

3. Prince M, Bryce R, Albanese E, Wimo A, Ribeiro W and Ferri CP. The global prevalence of dementia: a systematic review and metaanalysis. *Alzheimers Dement.* 2013; 9(1): 63-75. e62. doi: 10.1016/j.jalz.2012.11.007.

4. Mangialasche F, Solomon A, Winblad B, Mecocci P and Kivipelto M. Alzheimer's disease: clinical trials and drug development. *Lancet Neurol.* 2010; 9(7): 702-716. doi: 10.1016/S1474-4422(10)70119-8.

5. Wimo A, Jönsson L, Bond J, Prince M, Winblad B and International AD. The worldwide

effects, this formulation may be applied for an extended period, mainly for preliminary steps of dementia or preferably as prevention.

Author contribution

F.D: experimental parts, writing the manuscript, A.A and MM.Z: Supervision, Experimental validation, and developing the draft of the paper.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgment

Shiraz University of Medical Sciences supported this work as Thesis of Pharm-D degree of Farid Dabaghian [Grant No.: 8864].

economic impact of dementia 2010. *Alzheimers Dement*. 2013; 9(1): 1-11. doi: 10.1016/j.jalz. 2012.11.006.

6. Francis PT. The interplay of neurotransmitters in Alzheimer's disease. *CNS Spectr.* 2005; 10(S18): 6-9. doi: 10.1017/ S1092852900014164.

7. Garcia-Alloza M, Zaldua N, Diez-Ariza M, Marcos B, Lasheras B, Javier Gil-Bea F and Ramirez MJ. Effect of selective cholinergic denervation on the serotonergic system: implications for learning and memory. *J. Neuropathol. Exp. Neurol.* 2006; 65(11): 1074-1081. doi: 10.1097/01.jnen. 0000240469.20167. 89.

8. Heinrich M and Teoh HL. Galanthamine from snowdrop-the development of a modern drug against Alzheimer's disease from local Caucasian knowledge. *J. Ethnopharmacol.* 2004; 92(2-3): 147-162. doi: 10.1016/ j.jep.2004.02. 012.

9. Galimberti D and Scarpini E. Diseasemodifying treatments for Alzheimer's disease. *Ther. Adv. Neurol. Disord.* 2011; 4(4): 203-216. doi: 10.1177/1756285611404470.

10. Perry G, Cash AD and Smith MA. Alzheimer disease and oxidative stress. *J. Biomed. Biotechnol.* 2002; 2(3): 120-123. doi: 10.1155/S1110724302203010.

11. Aliev G, Obrenovich ME, Reddy VP, Shenk JC, Moreira PI, Nunomura A, Zhu X, Smith MA and Perry G. Antioxidant therapy in Alzheimer's disease: theory and practice. *Mini Rev. Med. Chem.* 2008; 8(13): 1395-1406. doi: 10.2174/138955708786369582.

12. Costa N, Derumeaux H, Rapp T, Garnault V, Ferlicoq L, Gillette S, Andrieu S, Vellas B, Lamure M and Grand A. Methodological considerations in cost of illness studies on Alzheimer disease. *Health Econ. Rev.* 2012; 2(1): 18. doi: 10.1186/2191-1991-2-18.

13. Margolin RA, Araujo JA, Sheinerman K and Van Epps D. P4-280: Maximizing the potential of small companies in Alzheimer's disease research: Challenges and opportunities. *Alzheimers Dement.* 2015; 11(7S_Part_19): P893-P893. doi: 10.1016/j.jalz.2015.08.110.

14. Yang M, Xu DD, Zhang Y, Liu X, Hoeven R and Cho WCS. A systematic review on natural medicines for the prevention and treatment of Alzheimer's disease with meta-analyses of intervention effect of ginkgo. *Am. J. Chin. Med.* 2014; 42(03): 505-521. doi: 10.1142/S0192415X14500335.

15. Adhami HR, Farsam H and Krenn L. Screening of medicinal plants from Iranian traditional medicine for acetylcholinesterase inhibition. *Phytother. Res.* 2011; 25(8): 1148-1152. doi: 10.1002/ptr.3409.

16. Hosseinkhani A, Sahragard A, Namdari A and Zarshenas M. Botanical sources for Alzheimer's: A review on reports from traditional Persian medicine. Am. J. Alzheimers Dis. Other Demen. 2017; 32(7): 429-437. doi: 10.1177/ 1533317517717013.

17. Jamshidi AH, Eghbalian F, Mahroozade S, Mohammadi Kenari H, Ghobadi A and Yousefsani BS. Recommended natural products in Alzheimer's disease based on traditional Persian medicine. *J. Med. Plant.* 2020; 19(75): 17-29. doi: 10.29252/jmp.1975.17.

18. Yassin N, El-Shenawy S, Mahdy KA, Gouda N, Marrie A, Farrag A and Ibrahim B. Effect of Boswellia serrata on Alzheimer's disease induced in rats. *J. Arab Soc. Med. Res.* 2013; 8(1): 1-11. doi: 10.7123/ 01.JASMR. 0000429323.25743.cc.

19. Teymuori M, Yegdaneh A and Rabbani M. Effects of Piper nigrum fruit and *Cinnamum zeylanicum* bark alcoholic extracts, alone and in combination, on scopolamine-induced memory impairment in mice. *Res. Pharm. Sci.* 2021; 16(5): 474. doi: 10.4103/1735-5362.323914.

20. Saenghong N, Wattanathorn J, Muchimapura S, Tongun T, Piyavhatkul N, Banchonglikitkul C and Kajsongkram T. Zingiber officinale improves cognitive function of the middle-aged healthy women. *Evid. Based Complement. Alternat. Med.* 2012; 2012. doi: 10.1155/2012/383062.

21. Homayounpour P, Shariatifar N and Alizadeh-Sani M. Development of nanochitosanbased active packaging films containing free and nanoliposome caraway (*Carum carvi* L) seed extract. *Food Sci. Nutr.* 2021; 9(1): 553-563. doi: 10.1002/fsn3.2025.

22. Yusif RM, Abu Hashim II, Mohamed EA and Badria FA-E. Gastroretentive matrix tablets of boswellia oleogum resin: preparation, optimization, in vitro evaluation, and cytoprotective effect on indomethacin-induced gastric ulcer in rabbits. *AAPS Pharm. Sci. Tech.* 2016; 17(2): 328-338. doi: 10.1208/s12249-015-0351-8.

23. Shirazi M.H, Qarabadin-e-Kabir. Tehran: Institute of Medical History, Islamic and Complementary Medicine Studies, Iran University of Medical Sciences Publications; 2007. [In Persian].

24. Heravi MS. Qarabadin-e-salehi. Tehran: Institute of Medical History, Islamic and Complementary Medicine Studies, Iran University of Medical Sciences Publications; 2004. [In Persian].

25. Arzani MA. Qarabadin-e-Ghaderi. Tehran: Institute of Medical History, Islamic and Complementary Medicine Studies, Iran University of Medical Sciences Publications; 2007. [In Persian].

26. Azam Khan M. Qarabadin-e-Azam. Tehran: Institute of Medical History, Islamic and Complementary Medicine Studies, Iran University of Medical Sciences Publications; 2004. [In Persian].

27. Tonekaboni MM. Tohfat-ol-Moeminin. Qom: Nour Publications; 2011. [In Persian].

28. Avicenna. The Canon of Medicine. Vol. 3. Tehran: Soroush Publications; 2013. [In Persian].
29. Dabaghian F, Azadi A, Setooni M and Zarshenas MM. An overview on multi-ingredient memory enhancers and anti-Alzheimer's formulations from traditional Persian pharmacy. *Trends in Pharmaceutical Sci.* 2017; 3(4): 215-220. doi: 10.1111/TIPS.V3I4.152.

30. Geldart D, Abdullah EC, Hassanpour A, Nwoke LC and Wouters IJ. Characterization of powder flowability using measurement of angle of repose. *China Particuology*. 2006; 4(3-4): 104-107. doi: 10.1016/S1672-2515(07)60247-4.
31. British Pharmacopoeia Commission. *British Pharmacopoeia* 2017. London: TSO; 2017.

32. Adams RP. Identification of essential oil components by gas chromatography/mass spectroscopy. Fourth ed., Carol Stream: Allured publishing corporation; 2007: 544-545.

33. Shamsa F, Monsef H, Ghamooghi R and Verdian R. Spectrophotometric determination of total alkaloids in Peganum harmala L. using bromocresol green. *Res. J. Phytochem.* 2007; 1(2): 79-82. doi: 10.3923/rjphyto.2007.79.82.

34. Beutler JA. Natural products as a foundation for drug discovery. *Curr. Protoc. Pharmacol.* 2009; 46(1): 9-11. doi: 10.1002/0471141755. ph0911s46.

35. Süntar I. Importance of ethnopharmacological studies in drug discovery: role of medicinal plants. *Phytochem. Rev.* 2019; 19(5): 1199-1209. doi: 10.1007/s11101-019-09629-9.

36. Rutebemberwa E, Lubega M, Katureebe SK, Oundo A, Kiweewa F and Mukanga D. Use of traditional medicine for the treatment of diabetes in Eastern Uganda: a qualitative exploration of reasons for choice. *BMC Int. Health Hum. Rights.* 2013; 13(1): 1. doi: 10.1186/1472-698X-13-1.

37. Jalali A, Dabaghian F and Zarshenas MM. Alkaloids of *Peganum harmala*: Anticancer Biomarkers with Promising Outcomes. *Curr. Pharm. Des.* 2021; 27(2): 185-196. doi: 10.2174/1381612826666201125103941.

38. Alanis A, Calzada F, Cervantes J, Torres J and Ceballos G. Antibacterial properties of some plants used in Mexican traditional medicine for the treatment of gastrointestinal disorders. *J. Ethnopharmacol.* 2005; 100(1-2): 153-157. doi: 10.1016/j.jep.2005.02.022.

39. Dawid-Pać R. Medicinal plants used in treatment of inflammatory skin diseases. *Postepy. Dermatol. Alergol.* 2013; 30(3): 170. doi: 10.5114/pdia.2013.35620.

40. Rojas G, Lévaro J, Tortoriello J and Navarro V. Antimicrobial evaluation of certain plants used in Mexican traditional medicine for the treatment of respiratory diseases. *J. Ethnopharmacol.* 2001; 74(1): 97-101. doi: 10.1016/S0378-8741(00)00349-4.

41. Yang Y, Islam MS, Wang J, Li Y and Chen X. Traditional Chinese medicine in the treatment of patients infected with 2019-new coronavirus (SARS-CoV-2): a review and perspective. *Int. J. Biol. Sci.* 2020; 16(10): 1708. doi: 10.7150/ijbs.45538.

42. Kunle OF, Egharevba HO and Ahmadu PO. Standardization of herbal medicines-A review. *Int. J. Biodivers. Conserv.* 2012; 4(3): 101-112. doi: 10.5897/IJBC11.163.

43. Vandana G, Dhar VJ, Anupam S and Rohit D. Facts about standardization of herbal medicine: a review. *Chin. J. Integr. Med.* 2012; 10(10): 1077-1083. doi: 10.3736/jcim20121002.
44. Nasab FK and Khosravi AR. Ethnobotanical study of medicinal plants of Sirjan in Kerman Province, Iran. *J. Ethnopharmacol.* 2014; 154(1): 190-197. doi: 10.1016/j.jep.2014.04.003.

45. Mandegary A, Arab-Nozari M, Ramiar H and Sharififar F. Anticonvulsant activity of the essential oil and methanolic extract of Bunium persicum (Boiss). B. Fedtsch. *J. Ethnopharmacol.* 2012; 140(2): 447-451. doi: 10.1016/j.jep.2012.01.024.

46. Shah Z, Ali T and Shafi S. Phytopharmacological review of *Bunium persicum* (Boiss) B. fedtsch. *Journal of Drug Delivery and Therapeutics*. 2019; 9(2): 458-460. doi: 10.22270/jddt.v9i2.2509.

47. Bukvicki D, Gottardi D, Prasad S and Tyagi AK. The healing effects of spices in chronic diseases. *Curr. Med. Chem.* 2020; 27(26): 4401-4420. doi: 10.2174/0929867325666180831145800.
48. Gholamhoseinian A, Moradi M and Sharifi-Far F. Screening the methanol extracts of some Iranian plants for acetylcholinesterase inhibitory activity. *Res. Pharm. Sci.* 2009; 4(2): 105-112.

49. Koppula S, Kopalli SR and Sreemantula S. Adaptogenic and nootropic activities of aqueous extracts of Carum Carvi Linn (Caraway) fruit:

An Experimental Study in Wistar Rats. *Aust. J. Medical Herbal.* 2009; 21(3): 72-78.

50. Nasrollah GD. Indicative Medicinal Plants in Medicine and Industry: Macroscopic and Microscopic Controls. Isfahan: Chaharbagh; 2013: 159-161. [In Persian].

51. Ahmed HH, Salem AM, Sabry GM, Husein AA and Kotob SE. Possible Therapeutic Uses of Salvia triloba and Piper nigrum in Alzheimer's Disease–Induced Rats. *J. Med. Food.* 2013; 16(5): 437-446. doi: 10.1089/jmf.2012.0165.

52. Subedee L, Suresh R, Jayanthi M, Kalabharathi H, Satish A and Pushpa V. Preventive role of Indian black pepper in animal models of Alzheimer's disease. *J. Clin. Diagn Res.* 2015; 9(4): FF01-FF04. doi: 10.7860/JCDR/2015/8953.5767.

53. Chonpathompikunlert P, Wattanathorn J and Muchimapura S. Piperine, the main alkaloid of Thai black pepper, protects against neurodegeneration and cognitive impairment in animal model of cognitive deficit like condition of Alzheimer's disease. *Food Chem. Toxicol.* 2010; 48(3): 798-802. doi: 10.1016/ j.fct.2009. 12.009.

54. Essa MM, Vijayan RK, Castellano-Gonzalez G, Memon MA, Braidy N and Guillemin GJ. Neuroprotective effect of natural products against Alzheimer's disease. *Neurochem. Res.* 2012; 37(9): 1829-1842. doi: 10.1007/s11064-012-0799-9.

55. Vurmaz A, Bozkurt F and Akbulut MD. The effect of betaine and piperine on rats with created alzheimer-like dementia by AF64A. *Kocatepe Tip. Dergisi.* 2020; 21(2): 183-190. doi: 10.18229/kocatepetip.579496.

56. Hritcu L, Noumedem JA, Cioanca O, Hancianu M, Kuete V and Mihasan M. Methanolic extract of Piper nigrum fruits improves memory impairment by decreasing brain oxidative stress in amyloid beta (1–42) rat

model of Alzheimer's disease. *Cell Mol. Neurobiol.* 2014; 34(3): 437-449. doi: 10.1007/ s10571-014-0028-y.

57. Mostafa NM, Mostafa AM, Ashour ML, Elhady SS. Neuroprotective effects of black pepper cold-pressed oil on scopolamine-induced oxidative stress and memory impairment in rats. *Antioxidants*. 2021; 10(12): 1993-2008. doi: 10.3390/antiox10121993.

58. Cheng Y, Dong Z and Liu S. β -Caryophyllene ameliorates the Alzheimer-like phenotype in APP/PS1 Mice through CB2 receptor activation and the PPAR γ pathway. *Pharmacol.* 2014; 94(1-2): 1-12. doi: 10.1159/000362689.

59. Hu Y, Zeng Z, Wang B and Guo S. Transcaryophyllene inhibits amyloid β (A β) oligomerinduced neuroinflammation in BV-2 microglial cells. *Int. Immunopharmacol.* 2017; 51: 91-98. doi: 10.1016/j.intimp.2017.07.009.

60. Wang J, Guo Y and Li GL. Current status of standardization of traditional Chinese medicine

in China. *Evid. Based Complement. Alternat. Med.* 2016; 2016. doi: 10.1155/2016/9123103.

61. Govindaraghavan S and Sucher NJ. Quality assessment of medicinal herbs and their extracts: Criteria and prerequisites for consistent safety and efficacy of herbal medicines. *Epilepsy Behav.* 2015; 52: 363-371. doi: 10.1016/j.yebeh. 2015.03.004.

62. Liang Y, Xie P and Chau F. Chromatographic fingerprinting and related chemometric techniques for quality control of traditional Chinese medicines. *J. Sep. Sci.* 2010; 33(3): 410-421. doi: 10.1002/jssc.200900653.

How to cite this article: Dabaghian F, Azadi A, Zarshenas MM. Design, reformulation, and standardization of a traditional-based memory enhancer herbal preparation originated from Persian medicine. *Journal of Medicinal Plants* 2022; 21(82): 93-110. doi: 10.52547/jmp.21.82.93

Downloaded from jmp.ir on 2025-07-12



فصلنامه گیاهان دارویی

Journal homepage: <u>www.jmp.ir</u>



مقاله تحقيقاتي

طراحی، ساخت و استانداردسازی داروی گیاهی تقویت کننده حافظه مبتنی بر طب سنتی ایران
فرید دباغیان ^{۲۰۱} ، امیر آزادی ^۳ ، محمدمهدی زرشناس ^{۲۰۱.} *
ا مرکز تحقیقات فرآوری گیاهان دارویی، دانشگاه علوم پزشکی شیراز، شیراز، ایران
۲ گروه فارماکوگنوزی، دانشکده داروسازی، دانشگاه علوم پزشکی تهران، تهران، ایران
۳گروه فارماسیوتیکس، دانشکده داروسازی شیراز، دانشگاه علوم پزشکی شیراز، شیراز، ایران
*گروه فیتوفارماسیوتیکس (داروسازی سنتی)، دانشکاره داروسازی، دانشگاه علوم پزشکی شیراز، شیراز، ایران

چکیدہ	اطلاعات مقاله
مقدمه : آلزایمر، شایعترین نوع دمانس میباشد که در طب سنتی تحت عناوینی همچون "فراموشکاری"، "فسادِ	گلواژگان:
ذکر" و "نِسیان" ذکر شده است. با توجه به چالشهای این بیماری، تحقیقات جهت دستیابی به ترکیبات طبیعی	بيمارى آلزايمر
موثر در آن حائز اهمیت میباشد. هدف : هدف، ارائهی قرص گیاهی استاندارد موثر در پیشگیری و درمان آلزایمر	قرص
مبتنی بر طب سنتی ایرانی میباشد. روش بررسی : فرآوردههای مرکبه مرتبط از کتب مرجع طب سنتی ایران	زيره ايراني
گردآوری شد. از میان ۱۲۷ فرمولاسیون، فرآورده حاوی زیره ایرانی و فلفل سیاه از کتاب قانون ابن سینا انتخاب	فلفل سياه
و پس از ارزیابی اکسپیانهای مختلف، فرمولاسیون حاوی ۴۰ درصد زیره ایرانی، ۴۰ درصد فلفل سیاه و ۲۰ درصد	استانداردسازى
HPMC و PVP به عنوان محلول چسباننده، تهیه گردید. فرمولاسیون نهایی و هریک از اجزای آن جهت ارزیابی	طب سنتی
GC و استانداردسازی فرآورده، اسانسگیری شدند؛ همچنین عصارهگیری با استفاده از اتانول و دیکلرومتان به	
منظور تهیه اثر انگشت HPTLC و تعیین میزان آلکالوئید تام موجود در قرص با استفاده از روش طیفسنجی	
انجام شد. نتایج : آنالیز اسانس قرص موید حضور دو ترکیب ترانس کاریوفیلن (۲۹/۲۲ درصد) و گاماترپینن	
(۱۷/۷۲ درصد) بود و استانداردسازی قرص براساس این دو ترکیب انجام شد؛ به گونهای که بصورت میانگین در	
هر قرص ۰/۰۴۱ میلیگرم ترانس کاریوفیلن و ۰/۰۱۸ میلیگرم گاماترپینن وجود داشت. میزان آلکالوئید تام موجود	
در هر قرص، ۰/۰۶ میلیگرم معادل آتروپین محاسبه گردید. نتیجهگیری : این فرمولاسیون طبیعی میتواند به عنوان	
یک محصول طب سنتی ایرانی به صورت بالینی مورد ارزیابی قرار گیرد و به عنوان یک مکمل تقویتکنندهی	
حافظه ارائه شود.	

مخففها: AD، بیماری آلزایمر؛ TPM، طب سنتی ایران؛ HPMC، هیدروکسی پروپیل متیل سلولز؛ PVP، پلی وینیل پیرولیدون، GC، کروماتوگرافی گازی؛ MS، طیفسنجی جرمی؛ FID، آشکارساز یونش شعلهای؛ HPTLC، کروماتوگرافی لایه نازک با کارایی بالا * نویسنده مسؤول: <u>zarm@sums.ac.ir</u>

doi: 10.52547/jmp.21.82.93

© 2020. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<u>https://creativecommons.org/licenses/by-nc/4.0/</u>)