




Institute of
Medicinal Plants

Journal of Medicinal Plants

Journal homepage: www.jmp.ir



Research Article

The essential oil of *Alpinia officinarum* Hance: chemical compounds, antioxidant, antimicrobial, and anti-inflammatory activities

Nguyen Thi Bich Thuyen^{1,*}, Tran Thanh Men², Huynh Lien Huong¹, Cao Luu Ngoc Hanh¹

¹Faculty of Chemical Engineering, Can Tho University, Campus II, 3/2 Street, Ninh Kieu District, Can Tho City, Vietnam

²Can Tho University, Campus II, 3/2 Street, Ninh Kieu District, Can Tho City, Vietnam

ARTICLE INFO

Keywords:

Alpinia officinarum

Hance

Biological

Essential oil

Galangal

Steam distillation

ABSTRACT

Background: *Alpinia officinarum* Hance (*A. officinarum*) roots are used as a treatment for various illnesses, including colds, stomachaches, inflammation, and discomfort. **Objective:** This work aimed to evaluate the anatomy, chemical compounds, antioxidant, antimicrobial, and anti-inflammatory activities of *A. officinarum* roots essential oil (EO). **Methods:** First, the plant's anatomy was studied by the alum-green iodine staining method, followed by steam distillation to extract EOs. The determination of chemical compositions by Gas chromatography-mass spectroscopy (GC-MS). The antioxidant activities were conducted by free radical resistance by 1,1-diphenyl-2-picrylhydrazyl (DPPH), ABTS (2,2-Azino-bis-3-ethylbenzothiazoline-6-sulphonic acid), and FRAP (Ferric Reducing Antioxidant Power). The paper plate method was used to determine antimicrobial activity. Finally, the anti-denaturation of the Bovine serum albumin assay was measured for the anti-inflammatory properties of EO. **Results:** Anatomical results show that the EO of *A. officinarum* is distributed in the form of secretory sacs. By GC-MS, twenty-eight components were identified; the main compounds are Eucalyptol (24.57 %), β -Bisabolene (9.72 %), and β -Sesquiphellandrene (5.95 %). The IC₅₀ of FRAP, ABTS, and DPPH assays are 810.00 μ g/mL, 1343.71 μ g/mL, and 26953.21 μ g/mL, respectively. In addition, the EO exhibited anti-inflammatory activity with an IC₅₀ of 2.55 mg/mL, compared to the standard Diclofenac (1.03 mg/mL). Additionally, the antimicrobial activity test demonstrated that the EO exhibited antimicrobial activity against various bacterial strains, including *Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonella typhimurium*, *Staphylococcus aureus*, *Bacillus cereus*, and *Listeria innocua*. **Conclusions:** These findings demonstrate that EOs can be utilized in the production of anti-inflammatory compounds and ferric-reducing additives.

1. Introduction

With 53 genera and 1300 species, the Zingiberaceae family, the Zingiberaceae family is considered the largest family in the

Zingiberales family, including *Zingiber*, *Alpinia*, *Curcuma*, *Globba*, *Etlingera*, *Renealmia*, *Riedelia*, *Amomum*, *Aframomum*, *Boesenbergia*, *Meisteria*, *Hedychium*, and

Abbreviations: DPPH, 1,1-diphenyl-2-picrylhydrazyl; ABTS, 2,2-Azino-bis-3-ethylbenzothiazoline-6-sulphonic acid; BSA, Bovine serum albumin; D, diameter; EO, Essential oil; FRAP, Ferric Reducing Antioxidant Power; GC-MS, Gas chromatography-mass spectroscopy; IC₅₀, Half maximal inhibitory concentration

*Corresponding author: ntbthuyen@ctu.edu.vn

doi:

Received 15 July 2025; Received in revised form 14 December 2025; Accepted 28 December 2025

© 2023. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>)

Hornstedtia [1]. The family is commonly found in the tropical regions of Africa, the Americas, and Asia, particularly in countries such as Egypt, Sri Lanka, Malaysia, India, Indonesia, and Thailand, due to their favorable agroecological conditions [2]. Most of the Zingiberaceae species, such as *Zingiber* and *Alpinia* species (*Z. officinale* Roscoe, *Z. zerumbet* (L.) Roscoe ex Sm., *Z. cassumunar* Roxb., *Alpinia galanga* Linn., *A. conchigera* Griff., *A. katsumadai* Hayata, *A. oxyphylla* Miq., *A. zerumbet* (Pers.) Burt. et Smith, *A. purpurata* (Viell.) K. Schum., and *A. officinarum* Hance) have been employed in conventional medicine [1].

Alpinia officinarum Hance (*A. officinarum*), called galangal, belongs to the Zingiberaceae family, is utilized throughout Southeast Asia, particularly in Thailand and Indonesia, as a traditional culinary herb, and since it has antiviral, antibacterial, and antioxidant properties, it can be used as a treatment for a number of illnesses, such as colds, stomachaches, inflammation, and discomfort [2-4]. *A. officinarum* has long been utilized in Chinese, Ayurvedic, Thai, and Greek medicine to increase blood flow and decrease swelling, diarrhea, headaches, renal issues, ulcers, joint pain, and respiratory issues [5]. In addition to its reputation as a medicinal herb, *A. officinarum* is also utilized as a preservative and taste enhancer in cooking. In Asian cooking, it is a common pickling spice due to its distinct sweet and spicy scent, which enhances flavor and prolongs the expiry date [6]. The historic use of *A. officinarum* has been supported by recent investigations that have demonstrated its varied pharmacological properties. For example, Laksmiawati et al. (2022) revealed that *Alpinia* is an antioxidant agent due to its polyphenol and total flavonoid components [7]. There is

evidence that *Alpinia* can be used to treat inflammatory pain because it inhibits p38 mitogen-activated protein kinases, nuclear factor-kappa B, nod-like receptor protein 3 signals, TRPV1 activation, regulates the expression of COX-2, and NF- κ B/TNF- α [8, 9]. In addition, extracts from *A. officinarum* have demonstrated significant sun protection factor (SPF) and free radical scavenging properties, which enhance their efficacy as natural sunscreens [10]. Methanol extracts of *A. officinarum* also showed significant antibacterial activity against Gram-negative *Enterobacter sakazakii* and Gram-positive *Staphylococcus aureus* pathogens [11]. Besides, the antidiabetic effect is also shown by *Alpinia officinarum*, mainly through inhibiting the α -glucosidase enzyme to decrease the conversion of carbohydrates to glucose [12]. Regarding anticancer activity, Yakufu revealed that flavonoids from *A. officinarum* may be used for stomach cancer treatment, probably with minimal toxicity. It might be linked to the suppression of cancer-associated inflammation, tumor cell proliferation, and tumor angiogenesis [13]. The study of Panigrahi et al. (2023) showed that ethyl-p-methoxycinnamate in *A. officinarum* has antiproliferative and cytotoxic effects on human breast and cervical cancer cells, suggesting that it may have anticancer characteristics [14]. This study enriches the scientific data on anatomy, chemical compounds, antioxidant, antimicrobial, and anti-inflammatory activities of *A. officinarum* roots essential oil.

2. Materials and methods

2.1. Studies on Anatomy

A. officinarum roots were harvested in Can Tho City, cleaned, and the damaged roots were further removed. According to morphological

figures [15], plant identification was conducted at the Biological Laboratory of the School of Education, Can Tho University, Vietnam, and a sample code of RIENG-01 was encoded. The sample was then observed to contain vesicles of essential oil. Firstly, use a 5 % sodium hypochlorite solution to immerse the sliced roots for twenty minutes, and then remove the solution with deionized water. The samples were cleaned once again using distilled water after steeping in a 0.5 % acetic acid solution for fifteen minutes. Following cleaning, the samples were dyed using iodine green and alum carmine. It was eventually determined by microscopic inspection that the samples included essential oil (EO) vesicles [16-18].

2.2. Essential oils distillation

By employing the Clevenger apparatus and about a 1 mm sample size, a 1:2 (g/ml) ratio between the material and water, and 240 minutes of distillation, the essential oil (EO) of *A. officinarum* roots was extracted by steam distillation. Physicochemical indexes, chemical composition, and biological activity were assessed after the EO was removed from water with Na₂SO₄, and stored at 4 °C until use.

2.3. Analysis of essential oil

The EO chemical compositions were analyzed by Gas chromatography-mass spectrometry (GC-MS, Thermo Fisher Scientific), which ran at 70 eV in electron impact mode. Operating in SCAN mode (range 35–350 m/z), the mass spectrometer had a scan rate of 0.2 scans per second. The DB-5MS column was used to produce the reported results of the EOs analysis (30 m × 0.25 mm × 0.25

µm, Agilent Technologies, USA), using helium as the carrier gas, and 1 mL/min was employed. The injection system ran at 220 °C in split mode (40:1). The following was the program for the GC oven temperature: 60 °C as the starting temperature, maintained for five minutes, then a 3 °C/min ramp up to 250 °C, which was maintained for 10 minutes. 1 µl of the EO solution (diluted with CH₂Cl₂ in a ratio of 1:100 v/v) was injected for each analysis. The EO ingredients were identified by comparing their retention indices (RI), retention times (RT), and mass spectra with those of authentic samples and data already available in the Institute of Standards and Technology version 2.4 (2020).

2.4. Antioxidant activities

2.4.1. DPPH assay

Referred to Anh et al. [19], the DPPH (1,1-diphenyl-2-picrylhydrazine) free radical scavenging activity was performed as follows: 960 µL of various EO concentrations in methanol were mixed with 40 µl of a methanolic solution containing 1000 µg/mL of DPPH radicals. Additionally, a blank control was applied by combining 960 µl of methanol with 40 µl of DPPH solution. A popular antioxidant, ascorbic acid, was compared. The mixtures were measured spectrophotometrically at 517 nm following 30 minutes in the dark. The following formula (I %) was used to express the DPPH scavenging activity of the extract samples as the percentage of free radical inhibition. The experiment was repeated three times for each concentration of extract. The IC₅₀ value was determined by calculating the effective concentrations of the EOs required to scavenge 50 % of the DPPH in the test solution.

$$I(\%) = \frac{A_0 - A_c}{A_0} \times 100 \quad (1)$$

Ac: Samples absorbance; A₀: DPPH Absorbance (control)

2.4.2. ABTS assay

The experiment of ABTS + was carried out according to Sun et al. [20] with small adjustments. The reaction was performed at room temperature in the dark for 12 to 16 hours after a 2.45 mM potassium persulfate and 7 mM ABTS solution were combined. This solution was then diluted with ethanol, yielding an absorbance of 0.70 ± 0.05 at 734 nm. A specified quantity of 100 μ l of the standard and samples in different concentrations dissolved in 10 % DMSO was added to 900 μ l of the diluted ABTS + solution. After six minutes of mixing, the solution was then measured at 734 nm using spectrophotometry. The ABTS + inhibition percentage was calculated in the same way as the DPPH inhibition percentage.

2.4.3 FRAP assay

The FRAP of EO was referenced to Berrezig et al. [21]. 0.5 ml of the sample, 0.5 ml of 0.8 M phosphate buffer at pH 7, and 0.5 ml of 1 % potassium ferricyanide were combined and incubated at 50 °C for 20 minutes. Then, 0.5 ml of 10 % trichloroacetic acid was added to the mixture, stirred, and centrifuged for 10 minutes at 3000 rpm. Then, 0.1 ml of 0.1 % FeCl₃ and 0.5 ml of distilled water were combined with 0.5 ml of the supernatant. At a wavelength of 700 nm, the concentration at which the Abs value = 0.5 is used to calculate the EC₅₀ value [21-22].

2.5. Antimicrobial activity

The antimicrobial activity was estimated using the paper plate method, as described by Maleki and Bakhsh (2023) with slight adjustments [23]. A clean tube was filled with four milliliters of Luria-Bertani liquid medium and six of the microorganisms, then mixed at 200 rpm overnight. Next, 200 μ l of the mixture

($4-5 \times 10^8$ CFU/ml) was evenly distributed on a petri dish with Luria-Bertani solid medium. A sterile paper plate (6 mm, Whatman No.1) was then set on top of the Petri plates, and 10 μ l of diluted EO (10^0-10^{-1}) was added to it. The control for the samples was the negative control (DMSO 1 % and Tween 0.1 %) and the positive control (tetracycline 320 μ g/ml). After that, the disks were incubated for 18 hours at 37 °C. The diameter of the sterile zone is measured as D (mm), which includes the diameter of the paper disk.

2.6. Anti-denaturation of Bovine serum albumin assay

To evaluate the possible anti-arthritis effectiveness of *A. officinarum* EO, the in vitro protein denaturation assay was utilized, with slight modifications, as described in Ameena et al. (2023). A mixture of 0.5 mL of 0.5 % bovine serum albumin (BSA) and 0.5 mL of EO (0.16 mg/mL to 5.00 mg/mL). The mixture was allowed to stand at room temperature for 10 minutes before being heated for 25 minutes at 70 °C. Once the reaction mixture had cooled to room temperature, the absorbance at 660 nm was measured. As a positive control, diclofenac sodium was utilized. The following method was used to determine the % inhibition of protein denaturation:

$$\text{Inhibition (\%)} = \frac{\text{blank} - \text{sample}}{\text{blank}} \times 100 \quad (2)$$

Where blank is the buffer solution and the control, and the sample is a diclofenac Sodium or tested sample [24, 25].

2.7. Statistical analysis

Using the Microsoft Excel application (Microsoft 365), the study's findings were expressed as the average of three separate tests.

3. Results

3.1. Studies on anatomy

The morphology of *A. officinarum* root was displayed in Figure 1. at magnifications of 4X (Fig. 1.a), and 10X (Fig. 1.b). As can be seen from the pictures, the EOs of *A. officinarum* are

disseminated as secretory vesicles (circled in red), in large numbers and scattered throughout the root. This observation is similar to that of Trimanto et al., who revealed that the oils of *Alpinia galanga* roots scattered in the parenchyma cells [26].




Fig. 1. Magnification of white *Alpinia officinarum* Hance root at 4X (a) and 10X (b)

3.2. Chemical compositions

The chemical compositions of *A. officinarum* EO are listed in Table 1 and Figure 2, including 27 components, mainly sesquiterpene (50.0 %), followed by monoterpene (39.29 %) and other components (10.71 %). The highest contents are

Eucalyptol (24.57 %), β -bisabolene (9.72 %), β -sesquiphellandrene (5.95 %), β -pinene (5.85 %), α -bergamotene (5.54 %), α -pinene (4.12 %), contributing to the characteristic smell of *A. officinarum*.

Table 1. The EO compositions

No	Name	Area (%)	RT (min)	RI	Class
1	α -Pinene	4.12	4.679	936	Monoterpene
2	β -Phellandrene	1.49	5.289	940	Monoterpene
3	β -Pinene	5.85	5.419	943	Monoterpene
4	β -Myrcene	1.30	5.480	988	Monoterpene
5	2-Carene	0.56	6.041	1011	Monoterpene
6	P-Cymene	0.82	6.148	1026	Monoterpene
7	D-Limonene	3.66	6.278	1027	Monoterpene
8	Eucalyptol	24.57	6.388	1046	Monoterpene
9	δ -Terpinene	1.05	6.772	1062	Monoterpene
10	Terpinen-4-ol	3.39	9.136	1182	Monoterpene
11	α -Terpineol	2.04	9.390	1189	Monoterpene

Table 1. The EO compositions (Continued)

No	Name	Area (%)	RT (min)	RI	Class
12	β -Elemene	1.58	12.965	1398	Sesquiterpene
13	Eugenol Methyl Ether	2.33	13.022	1403	Sesquiterpene
14	β -Caryophyllene	2.57	13.586	1418	Sesquiterpene
15	α -Bergamotene	5.54	13.704	1435	Sesquiterpene
16	β -Farnesene	1.21	13.936	1455	Sesquiterpene
17	α -Caryophyllene	1.44	14.201	1457	Sesquiterpene
18	α -Curcumene	0.92	14.497	1478	Sesquiterpene
19	Germacrene D	2.66	14.614	1481	Sesquiterpene
20	Octadecane	2.60	14.719	1800	alkane
21	β -Bisabolene	9.72	14.947	1509	Sesquiterpene
22	β -Sesquiphellandrene	5.95	15.228	1529	Sesquiterpene
23	1,3-Diisopropyl-1,3 Cyclopentadiene	3.40	15.272	1012	Cyclic alkenes
24	Ar-Tumerone	2.59	17.415	1665	Sesquiterpene
25	α -Tumerone	1.77	17.494	1689	Sesquiterpene
26	(9E)-9-Icosene	1.85	17.676	2017	Hydrocarbon
27	α -trans-Bergamotol	1.81	17.907	1700	Sesquiterpene

RT (retention time), RI (retention indices), monoterpene (39.29%), sesquiterpene (50.0%), and others (10.71%).




Fig. 2. GC-MS spectrum of *A. officinarum* EO

3.3. Biological activities

3.3.1. Antimicrobial activity

The paper plate method for measuring the antibacterial activity of the six pathogens investigated is shown in Table 2 and Figure 3. The pure EO showed resistance against six different kinds of bacteria, including

Escherichia coli ATCC 25922TM, *Pseudomonas aeruginosa* ATCC 27853TM, *Salmonella typhimurium* ATCC 13311, *Staphylococcus aureus* ATCC 25923TM, *Bacillus cereus* ATCC 10876TM, and *Listeria innocua* ATCC 33090TM.

Strongest, the EO has good resistance to *Escherichia coli* with a sterile ring diameter of 17.33 ± 0.58 mm, followed by *Salmonella typhimurium* $\approx$ *Listeria innocua* (16.33 ± 1.50 mm) > *Pseudomonas aeruginosa* (16.00 ± 1.00 mm) > *Bacillus cereus* (15.33 ± 0.58 mm) > *Staphylococcus aureus* (13.33 ± 0.58 mm). When the EO concentration was diluted up to 10 times, the EO did not show antibacterial activity ($D = 8.00 \pm 0.00$ mm), as seen in Table 2.

Table 2. Antibacterial efficacy based on sterile ring diameter (D)

Strains	Sterile ring diameter D (mm)			
	Control concentration		EO dilution	
	Negative control	Positive control	10 ⁰	10 ⁻¹
	DMSO (1 %) & Tween (0.1 %)	Tetracycline (320 µg/µl)		
<i>Escherichia coli</i> ATCC 25922TM	8.00 ± 0.00	23.67 ± 1.15	17.33 ± 0.58	8.00 ± 0.00
<i>Pseudomonas aeruginos</i> ATCC 27853TM	8.00 ± 0.00	22.67 ± 1.15	16.00 ± 1.00	8.00 ± 0.00
<i>Salmonella typhimurium</i> ATCC 13311	8.00 ± 0.00	23.33 ± 0.58	16.33 ± 1.50	8.00 ± 0.00
<i>Staphylococcus aureus</i> ATCC 25923TM	8.00 ± 0.00	25.66 ± 0.58	13.33 ± 0.58	8.00 ± 0.00
<i>Bacillus cereus</i> ATCC 10876TM	8.00 ± 0.00	25.33 ± 1.15	15.33 ± 0.58	8.00 ± 0.00
<i>Listeria innocua</i> ATCC 33090TM	8.00 ± 0.00	27.00 ± 1.00	16.33 ± 0.58	8.00 ± 0.00




Fig. 3. The antibacterial activity of *A. officinarum* EO, (+): Tetracycline 320 µg/mL; (-): DMSO (1.0 %) & Tween (0.1 %)

3.3.2. Antioxidant activity

3.3.2.1. DPPH assay

Ascorbic acid was used in this study as a positive control to find the DPPH free radical scavenging activity of the EO. The relationship between ascorbic acid concentration and DPPH free radical inhibition is shown in Fig. 4. When the ascorbic acid concentration increased from 6.0 to 40.0 µg/ml, the inhibition percentage increased from 20.95 % ± 1.1 to 92.99 % ± 0.99. Through the regression equation in Fig. 4,

the IC₅₀ value of ascorbic acid was determined to be 20.41 µg/mL ± 1.30.

In parallel with the experiment of DPPH free radical inhibition of the ascorbic acid, the DPPH free radical inhibition of the EO was performed in Fig. 5. Through the equation $y = 0.0019x - 1.2111$, $R^2 = 0.991$, the IC₅₀ of the EO was found to be 26953.21 µg/mL ± 0.30. This result indicates that the EO has a weak ability to scavenge DPPH free radicals.




Fig. 4. DPPH free radical inhibition of ascorbic acid




Fig. 5. DPPH free radical inhibition of the EO

3.3.2.2. ABTS and FRAP Assay

In addition to the investigation of PDDH free radical activity, the ABTS free radical activity, as well as the ferric-reducing antioxidant power (FRAP) of the EO, were also studied and presented in Table 3, Figure 6, and Figure 7. The EO showed the best ferric-reducing antioxidant power with $IC_{50} = 810.00 \mu\text{g/mL} \pm$

0.01, followed by ABTS free radical scavenging activity ($IC_{50} = 1343.71 \mu\text{g/mL} \pm 0.01$).

3.3.3. Anti-inflammatory

By protein denaturation assay, the possible anti-arthritic effectiveness of *A. officinarum* EO was performed in Figure 8 and compared with the positive control, diclofenac (Figure 9).

Table 3. The antioxidant capacity of *A. officinarum* EO

Assays	DPPH	ABTS•+	FRAP
	(IC ₅₀ , μg/ml)		
Standards (Ascorbic acid)	20.41 ± 1.30 (y = 2.2677x + 3.727 R ² = 0.9919)	3.05 ± 0.10 (y = 16.328x + 0.1301 R ² = 0.9975)	4.30 ± 0.03 (y = 0.0506x + 0.2823 R ² = 0.9953)
EO	26953.21 ± 0.30 (y = 0.0019x - 1.2111 R ² = 0.991)	1343.71 ± 0.01 (y = 0.0291x + 10.898 R ² = 0.9526)	810.00 ± 0.01 (y = 0.0003x + 0.257 R ² = 0.9539)




Fig. 6. The ABTS inhibition of the EO




Fig. 7. The Ferric reducing antioxidant power (FRAP) of EO




Fig. 8. Anti-inflammatory activity of EO




Fig. 9. Anti-inflammatory activity of diclofenac

Figure 8 shows that when the EO concentration increases from 0.16 to 5.00 mg/mL, the capacity to inhibit BSA gradually increases from 3.17 % $\pm$ 0.02 to 94.99 % $\pm$ 0.03. *A. officinarum*'s IC₅₀ value was found to be 2.55 mg/ml $\pm$ 0.02, which is greater than the standard Diclofenac's IC₅₀ value of 1.03 mg/ml $\pm$ 0.04 (Figure 9). This suggests that the EO has anti-inflammatory action that is 2.63 times lower than that of Diclofenac. The results demonstrate

the anti-inflammatory properties of *A. officinarum* EO.

4. Discussion

Chemical profiling of *A. officinarum* EO in three regions is represented in Table 4. As seen in the table, 1,8-Cineole, γ -Cadinene, α -terpineol, α -fenchyl acetate and β -Selinene are the main constituents of the EO of *A. officinarum* grown in China and Imphal (India) [27-28]. Meanwhile, α -fenchyl acetate,

1,8-cineole, borneol, bornyl acetate are the main compounds of *A. officinarum* EO in Southern India [29], and Eucalyptol (24.57%), β -bisabolene are the main constituents of the EO in Vietnam (this study). Thus, with the same

plant, the chemical compositions and contents differ depending on various factors, and the different geographical conditions in which the tree lives are also a contributing factor [30-33].

Table 4. Comparison of *A. officinarum* EO's main chemical constituents in different regions

No	Can Tho, Vietnam (This study)	China [27]	Imphal, India [28]	Southern India [29]
01	Eucalyptol (24.57 %)	1,8-Cineole (9.06 %)	1,8-cineole (28.3 %)	α -Fenchyl acetate (40.9 %)
02	β -Bisabolene (9.72 %)	γ -Cadinene (6.39 %)	α -fenchyl acetate (15.2 %)	1,8-cineole (9.4 %)
03	β -Sesquiphellandrene (5.95 %)	α -Terpineol (3.73 %)	carotol (8.9 %)	borneol (6.3 %)
04	β -Pinene (5.85 %)	β -Selinene (2.70 %)	α -Terpineol (6.7 %)	bornyl acetate (5.4 %)
05	α -Bergamotene (5.54 %)	α -Bergamotene (2.46 %)	β -pinene (3.1 %)	Elemol (3.1 %)
06	α -Pinene (4.12 %)	α -Pinene (0.72 %)	α -pinene (1.2 %)	Camphene (2.43 %)
07	D-Limonene (3.66 %)	D-Limonene (1.24 %)	zingiberenol (1.6 %)	Myrcene (1.8 %)
08	Terpinen-4-ol (3.39 %)	β -Pinene (0.60 %)	terpinen-4-ol (1.2 %)	<i>p</i> -Cymene (1.1 %)

The antimicrobial activity of the EO in this study was similar to that of the EO grown in India [34], as revealed by Indrayan et al. (2007), the zone diameter of inhibition of the EO against the microbial strains ranged from 12.3 – 17.1 mm for the tested strains including *Staphylococcus aureus*, *Bacillus subtilis*, and *Escherichia coli* [34]. Compared with the EOs grown elsewhere, for example, Zhang et al. (2020) studied 10 EO samples harvested in 10 provinces in China and found that the EOs were resistant to *Staphylococcus aureus* and *Bacillus subtilis* with diameters ranging from 2.10 to 10.9 mm. However, eight samples were not resistant to *Escherichia coli*, and nine samples were not resistant to *Pseudomonas aeruginosa* [27]. Thus, the different antibiotic activity of EOs from distinct locations may be due to the influence of their different chemical components.

The DPPH assay showed that the IC₅₀ of the EO was 26953.21 μ g/ml $\pm$ 0.30. This result indicates that the EO has a weak ability to

scavenge DPPH free radicals. The result is similar to the outcomes of Rahmi et al. (2024) and Bellarmine et al. (2024), who reported that the EC₅₀ values of *A. officinarum* rhizome essential oil were 20400 μ g/ml and 37580 μ g/ml, respectively, with no special activity towards DPPH [35, 36]. Thus, there appears to be a relationship between the primary chemical composition of oil and its ability to combat DPPH free radicals. Eucalyptol (24.57 %) is the main ingredient in *A. officinarum* EO in this study, as well as the main compound in *A. officinarum* EO (27.52 %) in Bellarmine's report [36] did not show anti-DPPH free radical activity. However, the activity can also be influenced by other ingredients and the interactions between them in the mixture [37, 38].

Regarding ABTS and FRAP Assay, the EO displayed the best ferric-reducing antioxidant power with IC₅₀ = 810.00 μ g/ml $\pm$ 0.01, followed by ABTS free radical scavenging activity (IC₅₀ =1343.71 μ g/ml $\pm$ 0.01). In the

literature, only two studies about the IC₅₀ index of ABTS and the FRAP experiment of the EO were found. As Belamine et al. reported, the *A. officinarum* EO in Morocco has ABTS free radical scavenging ability with IC₅₀ = 45230 µg/mL [36], which is 33 times higher than the IC₅₀ in this study, or according to Ivanović et al., the EO of the *A. officinarum* showed weak activity toward both ferric reducing power and ABTS [3]. But *A. officinarum* showed strong antioxidant activity in its extracts; for example, Laksmiawati et al. said that the 70 % ethanol extract of *Alpinia galanga* had an IC₅₀ for ABTS of 121.20 µg/ml [39]. In addition, the water, ethanol, and water/ethanol extracts (50:50 v/v) of *Alpinia galanga* for ABTS had IC₅₀ of 33.007, 16.119, 12.377 µg/mL, and for FRAP of 1.332 ± 0.002, 1.976 ± 0.002 and 2.020 ± 0.003 µg/ml, respectively [40].

It is reported that 10 samples of *A. officinarum* EO exhibit a good inhibitory effect on 12-O-tetradecanoylphorbol-13-acetate-induced mice's ear edema with an inhibition percentage of 64.42 % - 91.62 % compared to the anti-inflammatory medication dexamethasone (86.95 %) [26]. From the in vitro results of Rajendiran et al., it is evident that *A. officinarum* EO efficiently reduced HT-29 cell growth and suppressed the production of inflammatory mediators [41].

References

1. Yuandani, Jantan I, Haque A, Rohani AS, Nugraha SE, Salim E, Septama AW, Juwita NA, Khairunnisa NA, Nasution HR, Utami DS and Ibrahim S. Immunomodulatory effects and mechanisms of the extracts and secondary compounds of *Zingiber* and *Alpinia* species: A review. *Front. Pharmacol.* 2023; 14. 1222195. doi: 10.3389/fphar.2023.1222195.
2. Abd Rahman IZ, Adam SH, Hamid AA, Mokhtar MH, Mustafar R, Kashim MIAM ,

5. Conclusion

A. officinarum essential oil is in the form of secretory vesicles throughout the roots. The essential oil consists mainly of sesquiterpene (50.0 %), then monoterpene (39.29 %). In relation to anti-bacterial properties, the essential oil inhibited six different kinds of bacteria. Regarding anti-inflammatory capabilities, *A. officinarum* roots' essential oil displayed a moderate anti-inflammatory effect with the IC₅₀, which is 2.63 times lower than the standard. In terms of antioxidant properties, *A. officinarum* showed good ferric-reducing antioxidant power (FRAP). These biological activity findings demonstrate that essential oils can be utilized in the production of anti-inflammatory agents as well as ferric-reducing antioxidant additives.

Author contributions

NTBT conducted most of the experiments, including those on antioxidant, antimicrobial, and anti-inflammatory activities. TTM performed the antimicrobial experiment, while HLH carried out the anatomy, and chemical compounds were analyzed by CLNH. All authors read and approved the final manuscript.

Conflicts of interest

No potential conflict of interest was reported by the authors.

- Febriza A and Mansor NI. Potential Neuroprotective Effects of *Alpinia officinarum* Hance (Galangal): A review. *Nutrients*. 2024; 16(19): 3378. doi: 10.3390/nu16193378.
3. Ivanović M, Makoter K and Razboršek MI. Comparative study of chemical composition and antioxidant activity of essential oils and Crude extracts of four characteristic *Zingiberaceae* Herbs. *Plants*. 2021; 10(3): 501. doi: 10.3390/plants10030501.

4. Lei X, Wang J, Zuo K, Xia T, Zhang J, Xu X, Liu Q and Li X. *Alpinia officinarum* Hance: a comprehensive review of traditional uses, phytochemistry, pharmacokinetic and pharmacology. *Front. Pharmacol.* 2024; 15: 1414635. doi: 10.3389/fphar.2024.1414635.
5. Akbar S (Ed.). *Alpinia officinarum* Hance. (Zingiberaceae). In Handbook of 200 Medicinal Plants: A Comprehensive Review of Their Traditional Medical Uses and Scientific Justifications. Springer International Publishing: Cham, Switzerland. 2020, pp: 217-224. doi: 10.1007/978-3-030-16807-0.
6. Hamad A, Djalil AD and Hartanti D. Galangal and ginger essential oils exerted microbial growth inhibitory activity and preservation potential on tofu. *Food Res.* 2023; 7(Supp. 1): 27-34. doi: 10.26656/fr.2017.7(S1).7.
7. Laksmiawati DR, Pratami DK, Widowati W, Kusuma HSW, Wijayanti CR and Rizal R. The Potency of *Alpinia galanga* as natural antioxidant. *Maj. Obat Tradis.* 2022; 27(3): 165-171. doi: 10.22146/mot.72450.
8. Lin K, Fu D, Wang Z, Zhang X and Zhu C. Analgesic and anti-inflammatory effects of galangin: A potential pathway to inhibit transient receptor potential vanilloid 1 receptor activation. *Korean. J. Pain.* 2024; 37(2): 151-163. doi: 10.3344/kjp.23363.
9. Hassanein EHM, El-Maksoud MSA, Ibrahim IM, Abd-alhameed EK, Althagafy HS, Mohamed NM and Ross SA. The molecular mechanisms underlying anti-inflammatory effects of galangin in different diseases. *Phytother. Res.* 2023; 37(7): 3161-3181. doi: 10.1002/ptr.7874.
10. Khan MS, Adnan Q and Akhtar N. Profiling of phytochemicals using LC-ESI-MS2, in vitro, in vivo characterization and cosmeceutical effects of *Alpinia galanga* (wild) extract loaded emulgel. *J. Cosmet. Dermatol.* 2023; 22(5): 1628-1641. doi: 10.1111/jocd.15623.
11. Fu J, Wang Y, Sun M, Xu Y and Chen L. Antibacterial activity and components of the Methanol-phase extract from rhizomes of pharmacophagous plant *Alpinia officinarum* Hance. *Molecules.* 2022; 27(13): 4308. doi: 10.3390/molecules27134308.
12. Tian Y, Jia X, Wang Q, Lu T, Deng G, Tian M and Zhou Y. Antioxidant, antibacterial, enzyme inhibitory, and anticancer activities and chemical composition of *Alpinia galanga* flower essential oil. *Pharmaceuticals.* 2022; 15(9): 106. doi: 10.3390/ph15091069.
13. Yakufu M, Hailiwu R, Cong Y, Habaike A, Wang X and Abulizi P. Antitumor activity of flavonoids from *Alpinia officinarum* hance on gastric cancer. *Eur. J. Inflamm.* 2021; 19: 1-11. doi: 10.1177/20587392211051119.
14. Panigrahi R and Parida R. Preliminary screening of in vitro *Kaempferia galanga* oil for anti-proliferative activities against HeLa and MCF-7 cell. *Pharmacogn. Mag.* 2023; 19(3): 660-666. doi: 10.1177/09731296231172544.
15. Ho PH. Plants in Vietnam: Vietnamese. Youth Publishing House; 1999: 440-441
16. Bercu R and Popoviciu DR. Comparative anatomical study on leaves of three *Euphorbia* L. species. *Wulfenia.* 2015; 22: 271-6.
17. Plugatar YV, Sakhno TM, Bulavin I, Shevchuk OM and Feskov S. Morphology, anatomy and essential oil characterization of *Pinus radiata* needles in the conditions of the Southern Coast of the Crimea. In: IOP Conf. Ser.: Earth Environ. Sci. Russian; 2021; 843(1): 012046. doi: 10.1088/1755-1315/843/1/012046.
18. Thuyen NTB, Phong HQ, Hanh CLN, Huong HL and Men TT. Study on morphology, distillation process, chemical compositions, and biological activities of *Mentha arvensis* L. var. *javanica* (Blume) Hook. f essential oil. *J.*

- Taiwan Inst. Chem. Engrs.* 2023; 153: 105209. doi: 10.1016/j.jtice.2023.105209.
19. Anh VTT, Trang DTX, Kamei K, Linh TC, Pham-khanh NH, Tuan NT and Danh LT. Phytochemicals, antioxidant and antidiabetic activities of extracts from *Milusa velutina* flowers. *Horticul.* 2021; 7(12): 555. doi: 10.3390/horticulturae7120555.
20. Sun L, Wang H, Du J, Wang T and Yu D. Ultrasonic-assisted extraction of grape seed procyanidins, preparation of liposomes, and evaluation of their antioxidant capacity. *Ultrason. Sonochem.* 2024; 105: 106856. doi: 10.1016/j.ultsonch.2024.106856.
21. Berrezig AE, Achraf K, Kheira M and Yazid A. Phytochemical analysis and antioxidant activities of different extracts of *Silene vulgaris* (Moench) Garcke roots and leaves grown wild in the Algerian West. *Egypt. J. Chem.* 2024; 67(12): 67-76. doi: 10.21608/EJCHEM.2024.272992.9390.
22. Paiva L, Rego C, Lima E, Marcone M and Baptista J. Comparative analysis of the Polyphenols, Caffeine, and Antioxidant activities of green tea, white tea, and flowers from azorean *Camellia sinensis* varieties affected by different harvested and processing conditions. *Antioxidants* 2021; 10(2): 183. doi: 10.3390/antiox10020183.
23. Hajian-Maleki H and Shams-bakhsh M. Identification of the chemical profile and evaluation of the antimicrobial effect of *Eryngium billardieri* Delar essential oil component against bacterial species of agricultural and food interest. *Front. Microbiol.* 2023; 14: 1249780. doi: 10.3389/fmicb.2023.124978.
24. Aameena M, Arumugham M, Ramalingam K and Rajeshkumar S. Evaluation of the anti-inflammatory, antimicrobial, antioxidant, and cytotoxic effects of Chitosan thiocolchicoside-lauric acid nanogel. *Cureus.* 2023; 15(9): e46003. doi: 10.7759/cureus.46003.
25. Derbel H, Elleuch J, Mahfoudh W, Michaud P, Fendri I and Abdelkafi S. *In vitro* antioxidant and anti-inflammatory activities of bioactive proteins and peptides from *Rhodomonas* sp. *Appl. Sci.* 2023; 13(5): 3202. doi: 10.3390/app13053202.
26. Trimanto T, Hapsari L and Diwiyanti D. *Alpinia galanga* (L.) Willd: Plant morphological characteristics, histochemical analysis and review on pharmacological. *ICLST.* 2021; 2353(1): 030021. doi: 10.1063/5.0052687.
27. Zhang L, Pana C, Oua Z, Liang X, Shia Y, Chia L, Zhang Z, Zheng X, Lia C and Xiang H. Chemical profiling and bioactivity of essential oils from *Alpinia officinarum* Hance from ten localities in China. *Ind. Crops Prod.* 2020; 153: 112583. doi: 10.1016/j.indcrop.2020.112583.
28. Rana VS, Verdeguer M and Blazquez MA. GC and GC/MS Analysis of the Volatile Constituents of the Oils of *Alpinia galanga* (L.) Willd and *A. officinarum* Hance Rhizomes. *J. Essent. Oil Res.* 2010; 22(6). doi: 10.1080/10412905.2010.9700388.
29. Jirovetz L, Buchbauer G, Shafi MP and Leela NK. Analysis of the essential oils of the leaves, stems, rhizomes, and roots of the medicinal plant *Alpinia galanga* from southern India. *Acta Pharm.* 2003; 53(2): 73-81.
30. Santos SM, Cardoso CAL, Junior PLO, Silva ME, Pereira ZV, Silva RMMF and Formagio ASN. Seasonal and geographical variation in the chemical composition of essential oil from *Allophylus edulis* leaves. *S. Afr. J. Bot.* 2023; 154: 41-45. doi: 10.1016/j.sajb.2022.12.013.
31. Khan M, Dar NA, Alie BA, Dar SA, Lone AA, Mir GH, Fayaz U, Ali S, Tyagi A, El-Sheikh MA and Alansi S. Unraveling the variability of essential oil composition in

- different accessions of *Bunium persicum* collected from different temperate microclimates. *Molecules*. 2023; 28(5): 2404. doi: 10.3390/molecules28052404.
32. Ogwu MC, Izah SC and Joshua MT. Ecological and environmental determinants of phytochemical variability in forest trees. *Phytochem. Rev.* 2025. doi: 10.1007/s11101-025-10066-0.
33. Lima A, Arruda F, Medeiros J, Baptista J, Madruga J and de Castro Lima EM. Variations in essential oil chemical composition and biological activities of *Cryptomeria japonica* (Thunb. ex L.f.) D. Don from different geographical origins—A critical review. *Appl. Sci.* 2021; 11: 11097. doi: 10.3390/app112311097.
34. Indrayan AK, Garg SN, Rathi AK and Sharma V. Chemical composition and antimicrobial activity of the essential oil of *Alpinia officinarum* rhizome. *Indian J. Chem.* 2007; 46(B): 2060-2063.
35. Rahmi D, Yunilawati R, Jati BN and Astuti RI. Antioxidant and antiaging activities of cinnamic acid from Indonesian *Alpinia galangal* oil. In: 4th International Conference on Chemical Processing and Engineering (4th IC3PE). Indonesia; 2024, 3027: 020001. doi: 10.1063/5.0204732.
36. Belamine H, El Ghallab Y, Zahid B, Kadiri F, Kasrati A and Said AAH. *Alpinia officinarum* Hance essential oil as potent antipseudomonal agent: Chemical profile, antibacterial activity, and computational study. *Chem. Biodiver.* 2025; 22(2): e202401344. doi: 10.1002/cbdv.202401344.
37. Lakache Z, Hacib H, Aliboudhar H, Toumi M, Mahdid M, Lamrani N, Tounssi H and Kameli A. Chemical composition, antidiabetic, anti-inflammatory, antioxidant, and toxicity activities of the essential oil of *Fortunella margarita* peels. *J. Biol. Res.* 2022; 95: 10641. doi: 10.4081/jbr.2022.10641.
38. Bunse M, Daniels R, Gründemann C, Heilmann J, Kammerer DR, Keusgen M, Lindequist U, Melzig MF, Morlock GE, Schulz H, Schweiggert R, Simon M, Stintzing FC and Wink M. Essential oils as multicomponent mixtures and their potential for human health and well-being. *Front Pharmacol.* 2022; 13: 956541. doi: 10.3389/fphar.2022.956541.
39. Laksmitawati DR, Pratami DK, Widowati W, Kusuma HSW, Wijayanti CR and Rizal R. The potency of *Alpinia galanga* as natural antioxidant. *Trad. Med. J.* 2022; 27(3): 165-171. doi: 10.22146/mot.72450.
40. Kösea LP, Gülc I, Görenc AC, Namiesnik J, Martinez-Ayala AL and Gorinstein Sh. LC-MS/MS analysis, antioxidant and anticholinergic properties of galanga (*Alpinia officinarum* Hance) rhizomes. *Ind. Crops Prod.* 2015; 74: 712-721. doi: 10.1016/j.indcrop.2015.05.034.
41. Rajendiran V, Natarajan V and Devaraj SN. Anti-inflammatory activity of *Alpinia officinarum* hance on rat colon inflammation and tissue damage in DSS induced acute and chronic colitis models. *Food Sci. Hum. Wellness.* 2018; 7(4): 273-281. doi: 10.1016/j.fshw.2018.10.004.

How to cite this article: Nguyen Thi Bich Th, Tran Thanh M, Huynh Lien H, Cao Luru Ngoc H. The essential oil of *Alpinia officinarum* Hance: chemical compounds, antioxidant, antimicrobial, and anti-inflammatory activities. *Journal of Medicinal Plants* 2025; 24(96): 68-82.

doi: