

Antimicrobial Activity of the Leaves of *Pistacia khinjuk*

Taran M (Ph.D.)^{1*}, Sharifi M (Ph.D.)², Azizi E (M.Sc. Student)³, Khanahmadi M (M.Sc.)⁴

1- Microbiology Laboratory, Department of Biology, Razi University, Kermanshah, Iran

2- Department of Biology, Faculty of Science, Razi University, Kermanshah, Iran

3- Department of Biology, Faculty of Science, Razi University, Kermanshah, Iran

4- Department of Chemistry, Kermanshah Branch of ACECR, Kermanshah, Iran

*Corresponding author: Microbiology Laboratory, Department of Biology, Razi University, Kermanshah, Iran, Tel: +98- 831- 4274545

Email: mtaran@razi.ac.ir, mojtabataran@yahoo.com

Receive: 7 Nov. 2009

Acceptance: 17 Feb. 2010

Abstract

Background: *Pistacia khinjuk* is one of the three species *Pistacia* growing in Iran. The essential oils of some *Pistacia* plants contain variety of components with different therapeutical effects.

Objective: The purpose of this study was to provide the examination of antibacterial and antifungal effects of chloroform, ethyl acetate, ethyl alcohol and diethyl ether extracts obtained from the leaves of *P. khinjuk*.

Method: for obtaining different extracts of *P. khinjuk*, 30 gr of leaves powder was macerated with 120 ml of each solvent (chloroform, ethyl acetate, ethyl alcohol, diethyl ether). The antibacterial and antifungal activities of different extracts was individually tested against a panel of microorganisms, including *Bacillus subtilis*, *Enterococcus faecalis*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Candida albicans* and *Saccharomyces cerevisiae*

Results: Some major constituents of essential oil from the aerial parts of *P. khinjuk* are α -pinene, β -pinene, Myrcene, beta-caryophyllene, Germacrene B and Spathulenol. The extracts showed antimicrobial activity against bacteria (MIC = 0.02 - 0.5 mg/ml) and fungi (MIC = 0.06 - 0.4 mg/ml).

Conclusion: In conclusion the different leaves extracts of *P.khinjuk* (chloroform, ethyl acetate, ethyl alcohol, diethyl ether) like other species of *Pistacia* show antibacterial and antifungal activities.

Keywords: *Pistacia khinjuk*, Antibacterial activity, Antifungal activity

Plant

The genus *Pistacia* belongs to the family Anacardiaceae. Among 15 known species of pistachios, only 3 species grow in Iran, including *Pistacia Vera*, *Pistacia khinjuk* and *Pistacia atlantica*. They are the most important species of pistachio and for this reason, Iran is known as the origin of pistachios. *P.khinjuk* is spread in places where the attitude is 700 - 2000 m [1].

Uses in traditional medicine

Some species of *Pistacia* have been used in folk medicine and these plants are used as anti-inflammatory, antipyretic, antibacterial, antiviral, in treatment diarrhea and throat infection [2 - 4].

Previously isolated compounds

Phenolic composition consisting of gallic acid and quinic acid [5], flavonoids (kaempferol, quercetin, myricetin, taxifolin, distylin, naringenin, cyanidin) [6-8], essential oils (pinene, camphene, limonene, myrecene, gurjenene, etc) [9-12] and terpenoids [13, 14].

Plant material

The leaves of *P. khinjuk* were collected from Kerend of Kermanshah (western part of Iran) in May 2008. The voucher specimen (No.2406) is deposited in the Herbarium of Agricultural Faculty of Razi University, Kermanshah, Iran. The leaves were air dried in shadow. The essential oil hydrodistilled in a Clevenger-type apparatus (Clevenger 1928) according to the British method and Gas chromatography (GC) and Gas Chromatography-mass spectrometry (GC-MS) have been reported in the literature [15].

Tested materials

For obtaining different extracts of

P. khinjuk, 30 gr of leaves powder was macerated with 120 ml of each solvent (chloroform, ethyl acetate, ethyl alcohol, diethyl ether). Yields for different extracts were 16.9, 5.1, 12.9, 0.13 percent respectively.

Used microorganism

The antimicrobial activity of different extracts was individually tested against a panel of microorganisms, including *Bacillus subtilis* (ATCC 127111), *Enterococcus faecalis* (ATCC 29737), *Staphylococcus aureus* (ATCC 29737), *Staphylococcus epidermidis* (ATCC 12229), *Escherichia coli* (ATCC 8739), *Klebsiella pneumoniae* (ATCC 10031), *Candida albicans* (ATCC 10231) and *Saccharomyces cerevisiae* (ATCC 9763). Bacterial strains were cultured overnight at 37 °C in Mueller Hinton broth (Merck). Yeasts were cultured at 28 °C for 22h in Sabouraud dextrose broth (Merck). For the determination of antimicrobial activities MIC (minimum inhibitory concentration) method was employed.

Results

The major constituents of essential oil from the aerial parts of *P. khinjuk*, their Kovats indices and their percentage are given in table 1.

Antibacterial and antifungal activity of the different extracts is shown in the table 2. The MIC of the extracts against the test microorganisms were determined by the Micro dilution method [16]. Antibacterial and antifungal differences of the extracts was determined by One way variance analysis (ANOVA), using the SPSS 11.5 software package. Data were considered statistically significant at $p \leq 0.05$.

Discussion

Different extracts of the leaves of



Table 1- Chemical composition (%) of identified compounds in the oil of *P. khinjuk*

| | Component | Percent | ^a KI |
|----|----------------------------------|---------|-----------------|
| 1 | Tricyclene | 0.64 | 926 |
| 2 | α -pinene | 2.11 | 936 |
| 3 | camphene | 1.52 | 953 |
| 4 | Sabinene | 0.67 | 967 |
| 5 | β -pinene | 1.49 | 980 |
| 6 | Myrcene | 2.85 | 1000 |
| 7 | α -3-carene | 0.29 | 1010 |
| 8 | β cymene | 0.58 | 1024 |
| 9 | Limonene | 0.58 | 1029 |
| 10 | γ Terpinene | 1.15 | 1057 |
| 11 | Nonanol | 1.13 | 1104 |
| 12 | Comphenol | 0.63 | 1125 |
| 13 | Trans-pinocarveol | 0.35 | 1138 |
| 14 | Verbenol | 0.41 | 1148 |
| 15 | P Mentha-1,5-dien8-ol | 0.75 | 1169 |
| 16 | α -terpineol | 0.58 | 1195 |
| 17 | Bornyl Acetate | 2.77 | 1286 |
| 18 | δ -Elemene | 1.52 | 1339 |
| 19 | beta-caryophyllene | 3.67 | 1421 |
| 20 | α -Guaiene | 2.24 | 1444 |
| 21 | Aromadendrone<dehydro> | 8.80 | 1464 |
| 22 | Germacrene B | 9.53 | 1500 |
| 23 | Luparene | 1.38 | 1520 |
| 24 | Occidentalol cadinene <deltha> | 3.77 | 1532 |
| 25 | Thusopsanone | 3.84 | 1653 |
| 26 | Eudesmol | 2.29 | 1659 |
| 27 | Spathulenol | 20.87 | 1584 |
| 28 | α -Eudesmol | 6.78 | 1666 |
| 29 | Occidentalol Acetate | 3.68 | 1687 |
| 30 | Zizanol | 6.51 | 1699 |
| 31 | Khusimol | 1.48 | 1722 |
| 32 | Aristolone | 0.55 | 1768 |
| 33 | Muurolene<14-hydroxy- α > | 0.28 | 1842 |
| 34 | Dibutyl phthalate | 0.33 | 1956 |
| 35 | Pentacosune | 0.54 | 2500 |
| 36 | Heptacosane | 0.35 | 2580 |

^aKI: kovats indices on column

Table 2- The minimum inhibitory concentration (MIC, mg/ml) of the leaves of *P.khinjuk* against bacteria and fungi

| Microorganism | Extract | | | |
|-----------------------|------------|---------------|---------------|---------------|
| | Chloroform | Ethyl acetate | Ethyl alcohol | Diethyl ether |
| <i>B. subtilis</i> | 0.09 | 0.05 | 0.34 | 0.16 |
| <i>E. faecalis</i> | 0.21 | 0.38 | 0.19 | 0.36 |
| <i>S. aureus</i> | 0.04 | 0.13 | 0.09 | 0.42 |
| <i>S. epidermidis</i> | 0.25 | 0.06 | 0.25 | 0.13 |
| <i>E. coli</i> | 0.19 | 0.36 | 0.34 | 0.17 |
| <i>K. pneumoniae</i> | 0.02 | 0.13 | 0.04 | 0.16 |
| <i>C. albicans</i> | 0.10 | 0.34 | 0.43 | 0.38 |
| <i>S. cerevisiae</i> | 0.06 | 0.36 | 0.18 | 0.09 |

The results are the conclusion of three replicates.

P. khinjuk inhibited gram positive and gram negative bacteria probably due to its particular chemical pattern mainly the amounts of Germacrene B, α -pinene and β -pinene. Chloroform and ethyl alcohol extracts inhibited *K. pneumoniae* more than others bacteria. Ethylacetate and diethyl ether extracts inhibited *B. subtilis* and *S. epidermidis* respectively more than other bacteria.

Also the extracts of the leaves of *P. khinjuk* inhibited fungi. Among different extracts, chloroform extract inhibited growth of fungi more than others (MIC 0.10 for *C. albicans*

and 0.06 for *S. cerevisiae*). In conclusion the leaves extracts of *P.khinjuk* (especially chloroform extract) like other species of *Pistacia* show antibacterial and antifungal activities.

Acknowledgment

This study financially supported by Razi University of Kermanshah .We wish to express our thanks to Dr. S.M. Maassoumi and Mr. E. Samadian for identification of plant and laboratory help.

References

- Behboodi B.Sh. Ecological distribution study of wild pistachios for selection of rootstock. *Options Mediterran* 2003; 63: 61 - 6.
- Kordali S, Cakir A, Zengin H, Duru ME. Antifungal activities of the leaves of three *Pistacia* species grown in Turkey. *Fitoterapia* 2003; 74: 164 - 7.
- Villar A, Sanz MJ, Payo M. Hypotensive effect of *Pistacia lentiscus* L. *Int. J. Crude Drug Res.* 1987; 25 (1): 1 - 3.
- Benhammou N, Bekkara FA, Panovska T K. Antioxidant and antimicrobial activities of the *Pistacia lentiscus* and *Pistacia atlantica* extracts. *Afr. J. Pharmacol.* 2008; 2 (2): 22 - 8.
- Romani A, Pinelli P, Galardi C, Mulinacci N, Tattini M. Identification and Quantification of Galloyl Derivatives, Flavonoid Glycosides and Anthocyanins in Leaves of *Pistacia lentiscus* L. *Phytochem. Anal.* 2002; 13: 79 - 86.
- Kawashty SA, Mosharrata SAM, El-Gibali M, Saleh NAM. The flavonoids of four *Pistacia* species in Egypt. *Biochem. Syst. Ecol.* 2000; 28 (9): 915 - 7.
- Bouguet J. *Matiere Medicale Indigene de*



- l'Afrique du Nord. *Bull Soc. Pharmacol.* 1921; 28: 22 - 36.
8. Keys JD. Chinese Herbs, their Botany, Chemistry and Pharmacodynamics. Charles E. Tuttle Company, Rutland, 1976.
9. Longo L, Scardino A, Vasapollo G. Identification and quantification of anthocyanins in the berries of *Pistacia lentiscus* L., *Phillyrea latifolia* L. and *Rubia peregrina* L. *Food Innovat. Sci. Emerg. Tech.* 2007; 8: 360 - 4.
10. Ben Douissa F, Hayder N, Chekir-Ghedira L, Hammami M, Ghedira K, Mariotte AM. New study of the essential oil from leaves of *Pistacia lentiscus* L. (Anacardiaceae) from Tunisia. *Flav. Frag. J.* 2005; 20: 410.
11. Anastasia T, Katerina G, Eleni M, Prokopios M, Eugenia T. Composition and natiomeric Analysis of the Essential Oil of the Fruits and the Leaves of *Pistacia vera* from Greece. *Molecules* 2007; 12 (6): 1233 - 9.
12. Duru ME, Cakir A, Kordali S, Zengin H, Harmandar M, Izumi S. Chemical composition and antifungal properties of essential oils of three *Pistacia* species. *Fitoterapia* 2003; 74: 170 – 6.
13. Boar RB, Couchman LA, Jagues AJ, Perkins JM. Isolation from *Pistacia* resins of a bicyclic triterpenoid representing an apparent trapped intermediate of squalene 2, 3-epoxide cyclization. *J. Am. Chem. Soc.* 1984; 106 (8): 2476 - 7.
14. . Ansari SH, Ali M, Quadry JS. New tetracyclic triterpenoids from *Pistacia integerrima* galls. *Pharmazie* 1994; 49 (5): 356 - 7.
15. Ghasempour H, Shirinpour E, Heidari H. Analysis by Gas Chromatography-mass Spectrometry of Essential Oil from Seeds and Aerial Parts of *Ferulago angulata* (Schlecht.) Boiss Gathered in Nevakoh and Shahoo, Zagross Mountain, West of Iran. *Pakistan. J. Biol Sci.* 2007; 10 (5): 814 - 7.
16. NCCLS (National Committee for Clinical Laboratory Standards). Performance standards for antimicrobial susceptibility testing. 12th Informational Supplement 2002, M100.