Effects of *Pinus eldarica* Medw. Nut Extract on Blood Glucose and Cholesterol Levels in Hypercholesterolemic Alloxan-induced Diabetic Rats

Fallah huseini H $(Ph.D.)^1,$ Mehrzadi S $(D.V.M.)^{1,2},$ Ghaznavi H $(M.D.)^{2,3},$ Tajallizadehkhoob Y $(M.D.)^4,$ Fakhrzadeh H $(Ph.D.)^{4*}$

- 1- Pharmacology & Applied Medicine Department of Medicinal Plants Research Center, Institute of Medicinal Plants, ACECR, Karaj, Iran
- 2- Department of Pharmacology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran
- 3- Cellular and Molecular Research Center, Zahedan University of Medical Sciences, Zahedan, Iran
- 4- Endocrinology & Metabolism Research Institute, Tehran University of Medical Sciences, Tehran, Iran
- * Corresponding author: Endocrinology and Metabolism Research Institute, Tehran University of Medical Sciences, 5th floor, Dr Shareeati Hospital, North Karegar Avenue, Tehran, Iran

Tel: +98-21-84902476-7, Fax: +98-21-88220052

Email: Fakhrzad@tums.ac.ir

Received: 1 Dec. 2012 Accepted: 17 Mar. 2013

Abstract

Background: Diabetes mellitus type 2 with dyslipidemia is a common disease. Previous studies suggest that chemical constituent present in *pinus eldarica* (*P. eldarica*) nut posesess antioxidant properties and positively affect glucose metabolism. However blood glucose and cholesterol lowering effects of *P. eldarica* nut have not been studied so far.


Objective: The present study was undertaken to explore the possibility of anti-hyperglycemic and anti-hypercholesterolemic effects of the *P. eldarica* nut extract in hypercholesterolemic diabetic rats.

Methods: Sixty male wistar rats six months of age from central animal house of the institute of medicinal plants were selected. 10 rats were kept as normal group and diabetes was induced in the remaining rats by intraperitonial injection of 120 mg/kg aloxan monohydrate. After one week the diabetic rats with fasting blood glucose between 180 to 250 mg/kg were assigned to 5 groups of 10 rats each and were fed on hypercholesterolemic diet. One group was kept as control group (untreated diabetic rats) and *P. eldarica* nut extract in doses of 50, 100, 200 and 400 mg/kg was gavaged daily to the remaining rats. After one month, the fasting blood glucose, cholesterol and triglyceride levels were determined in all groups.

Results: The results indicate that fasting blood glucose in 200 and 400 mg/kg *P. eldarica* nut extract treated groups significantly decreased (P=0.000 and P=0.000) and fasting blood cholesterol and triglyceride levels did not change significantly compared with control group.

Conclusion: P. eldarica nut extract lowers blood glucose level withought affecting blood cholesterol and triglyceride levels in hypercholesterolemic diabetic rats.

Keywords: Pinus eldarica nut, Blood glucose, Rat, Diabetes, Hypercholesterolemia



Introduction

Diabetes is a chronic disorder of carbohydrate, fat and protein metabolism characterized by increased fasting and postprandial blood glucose levels. Diabetes mellitus type 2 is one of the most prevalent and fastest growing diseases in almost all countries [1]. Apart from conventional antidiabetic therapy, several studies have shown that medicinal plants have beneficial effects and improve glucose homeostasis in diabetic patients [2, 3]. A number of potential antidiabetic and anti-hyperlipidemic plants were used by herbalists in the Iranian folk medicine [2]. It has been reported that medicinal plants with antioxidant properties or dietary antioxidant intake have beneficial effects on diabetes and hyperlipidemia [4-6]. P. eldarica nut is one of the dietary antioxidants which is used as food and medicine in several countries and contains several phenols and essential fatty acids with antioxidant properties [7]. The Iranian pine. Pinus eldarica Medw. (P. eldarica) belongs to the botanical family pinaceae and is native to Transcaucasian region between Europe and Asia, and has been widely planted in Iran, Afghanistan and Pakistan [8, 9]. Various parts of P. eldarica (e.g., needles, buds, resin and nuts) have been widely used in traditional medicine for the treatment of bronchial asthma, skin wounds, skin irritations, allergic rashes and dermatitis in Russia and the Central Asian countries [10, 11]. In a study, analysis of the *P. eldarica* nut oil indicated several components including: β-caryophyllene, α-pinene, longifolene, α -humulene, δ -3-carene and β -pinene with antioxidant properties [7]. In a previous unpublished study, we found high concentrations of total polyphenols (table 1)

and fatty acids in *P. eldarica* nut indicating its antioxidant properties. Experimental studies in fact, strongly support the efficacy of polyphenols in chronic diseases including diabetes mellitus [12, 13]. In view of the traditional use of *P. eldarica* nuts as food and medicine and its established antioxidant properties, the present study was conducted to evaluate the possible anti-hyperglycemic and anti-hypercholesterolemic effects of *P. eldarica* nut in diabetic hypercholesterolemic rats.

Material and Methods

Plant material

P. eldarica cones were collected from Chitgar forest park (West of Tehran). The cones were collected between June and July of 2010. P. eldarica is preserved in the herbarium of Institute of Medicinal Plants (ACECR). Herbarium code of P. eldarica is 689 and it was identified by M. Ahvazi. Samples were dried in a dark place and at room temperature. The nuts were removed from cones and grouned to a powder by grinder.

Preparation of extract

The hydroalcholic extract of P. eldarica nut powder was prepared using 70% ethanol in water, using percolation method at room temperature. The powdered plant material was soaked initially in a solvent in a percolator and then sufficient amount of the solvent was added to cover material and kept for 24 hours with occasional stirring. The outlet of the percolator then was opened and the liquid contained therein was allowed to drip slowly. The procedure was repeated twice and the combined extractions was clarified filtration and concentrated to dryness on rotatory evaporator at a maximum of 40°C



temperature and under reduced pressure. The extract in the dosages of 50, 100, 200 and 400 mg was suspended in vehicle of 3% Tween 80 in distilled water.

Polyphenols determination

The polyphenols content of *P. eldarica* nut was determined by the High-performance liquid chromatography (HPLC) method developed by Dogan *et al.* [14].

Hypercholesterolemic diet

The hypercholesterolemic diet was prepared according to modified metod of Groot AP et al. In brief 3% cholesterol powder, 1% cholic acid and 6% animal ghee were mixed with 90% rats chow powder [15]. The food mixture was shaped into small pellets by a hand operated device and then dried it in oven at 50°C temperature.

Animals

60 adult male wistar rats aged six months with 200 ± 10 g weight were purchased from the central animal house of institute of medicinal plants. The animals were housed under standard conditions of light and dark cycle (12 hr light and 12 hour dark) with free access to food and water one week before starting the study.

Study protocol

10 rats were kept as normal group and alloxan monohydrate was injected intraperitonially in the doses of 120 mg/kg to the remaining 50 rats [16]. After one week the diabetic rats with fasting blood glucose between 180 to 250 mg/kg were selected and caged in five groups of 10 rats each and fed on hypercholesterolemic diet. One group was kept as control group (untreated diabetic rats) and

P. eldarica nut extract in doses of 50, 100, 200 and 400 mg/kg was gavaged daily to the remaining rats for one month. In control group, extract vehicle was gavaged daily for one month. After one month the fasting blood glucose, cholesterol and triglyceride levels were determined. The experiment protocol was approved by the Institute of Medicinal Plants ethical committee. The above doses were selected according to a previous pilot study in which the anti-hyperglycemic effect of several P. eldarica nut extract doses was tested in diabetic rats for three days.

Blood samples were drawn after 6 hour fasting. Blood glucose levels were determined by the glucose-oxidase method using Beckman Glucose-2 Analyzer. Blood cholesterol and triglyceride levels were measured by the auto analyzer Hitachi 902 using commercially available kits (Pars Azmon).

Statistical analysis

All values are expressed as means \pm SE. Data obtained were analysed using student's t-test to determine the statistical significance. p < 0.05 was considered as significant.

Results

The values for polyphenols contents of the *P. eldarica* nut are presented in Tables 1.

In alloxan induced diabetic rats fed on hypercholesterolemic diet the fasting blood glucose and cholesterol levels were significantly increased compared with normal (non-diabetes) group. In P. eldarica nut extract treated groups in the doses of 200 and 400 mg/kg, fasting blood glucose levels significantly (p=0.000 and P=0.000) decreased compared with control group (Table 2).

Administration of *P. eldarica* nut extract in different doses (50, 100, 200 and 400 mg/kg)

did not producing any significant changes in blood cholesterol and triglyceride levels compared with control group (Table 2).

Table 1- The polyphenols content (mg/1000 g extract) of P. eldarica nut (HPLC method)

Polyphenols	mg/1000g extract
Total phenols	483
Catechin	10.1
Epicatechin	10.3
Gallic acid	1.6
Vanillic acid	_
Para-coumaric acid	1.4
Ferullic acid	1.7
Ortho-coumaric acid	0.12
Tyrosol	29.1
Dimers of catechin and epicatechin	7.5
Unknown	38.18

Table 2- The effects of different doses of *P. eldarica* nut extract on fasting blood parameters of diabetic rats (n=10)

	Fasting blood glucose (mg/dl)	P- value compared to control	Triglyceride (mg/dl)	P- value compared to control	Cholesterol (mg/dl)	P- value compared to control
Normal	94.2 ± 20.7		98.2 ± 11.5		84.1 ± 9.3	
Control	$168.5 \pm 25.3*$	P=0.000	114.1 ± 33.5	P=0.068	98.8 ± 14.0	P=0.042*
P. eldarica nut extract 50 mg/kg	172.2 ± 33.9	P=0.808	112.1 ± 21.9	P=0.196	103.2 ± 15.2	P=0.261
P. eldarica nut extract 100 mg/kg	165.8 ± 22.3	P=0.842	109.5 ± 33.5	P=0.167	104.1 ± 15.3	P=0.496
P. eldarica nut extract 200 mg/kg	115.7 ± 19.5**	P=0.000	106.0 ± 15.6	P=0.112	99.8 ± 14.6	P=0.889
P. eldarica nut extract 400 mg/kg	114.8 ± 13.0**	P=0.000	107.9 ± 12.8	P=0.129	100.8 ± 16.0	P=0.811

Values are given as Mean \pm SD.

**= comparison with control group (student t-test).



^{*=} comparison with normal (non-diabetic) group (student t-test).

Discussion

In the present study we found that P. eldarica nut extract reduced fasting blood glucose in diabetic rats fed on hypercholesterolemic without diet any effects cholesterol significant on and triglyceride. The mechanisms involved in the glucose lowering effect of the P. eldarica nut extract are not yet clearly established. However few theories try to explain the mechanisms involved in its effect on hyperglycemia. It has been reported that fatty acids an important component of *Pinus* nut oil by suppressing appetite may contribute to a successful caloric-restriction regimen and directly influence the blood glucose metabolism [17]. Another suspected antihyperglycemic mechanism may be due to effects of antioxidant components P. eldarica nut extract. Hyperglycemia causes oxidative damage by generation of reactive oxygen species resulting in the acceleration of onset and progression of diabetes [18]. There are increasing evidences that polyphenols offer protection against oxidative stress and positively influence carbohydrate metabolism [19, 20]. P. eldarica nut contains appreciable amounts of phenolic compounds and other chemicals such as α -pinene,

β-pinene and β-caryophyllene with antioxidant properties [7, 21, 22]. The anti-diabetic and antioxidant activities of α -pinene, β-pinene, β-caryophyllene present in Juglans regia L. leaf essential oil have been reported [23, 24]. As these chemicals are important components of *P. eldarica* nut [7], the anti-hyperglycemic effect observed in the present study may be due to effects of these components on body metabolism of diabetic rats.

Conclusion


In conclusion, since *P. eldarica* nut induced significant anti-diabetic effects in hypercholesterolemic diabetic rats, further studies for determination of its active constituents and possible mechanisms responsible for anti-hyperglycemic effects are recommended.

Acknowledgements

This research was supported by a grant from the Institute of Medicinal Plants (ACECR), Karaj, Iran and Endocrinology and Metabolism Research Institute, Tehran University of Medical Sciences Tehran, Iran.

References

- **1.** Gerstein Hc, Miller Me, Byington Rp, Goff Dc Jr and Bigger Jt. Effects of intensive glucose lowering in type 2 diabetes. *N. Engl. J. Med.* 2008; 358: 2545 59.
- 2. Hasani-Ranjbar S, Larijani B and Abdollahi M. A systematic review of Iranian medicinal
- plants useful in diabetes mellitus. *Arch. Med. Sci.* 2008; 4: 285 92.
- **3.** Nahas R and Moher M. Complementary and alternative medicine for the treatment of type 2 diabetes. *Can. Fam. Physician* 2009; 55: 591 6.



- **4.** Montonen J, Knekt P, Järvinen R and Reunanen A, Dietary antioxidant intake and risk of type 2 diabetes. *Diabetes Care* 2004; 27: 362 6.
- **5.** Fallah Huseini H, Larijani B, Fakhrzadeh H, Radjabipour B, Toliat T and Reza M, The efficacy of *Silybum marianum* (L.) Gaertn. (silymarin) in the treatment of type 2 diabetes: a randomized, double-blind, placebocontrolled clinical trial. *Phytotherapy Res.* 2006; 20: 1036 9.
- **6.** Ruhe RC and McDonald BR, Use of antioxidant nutrients in the prevention and treatment of type 2 diabetes. *J. Am. College Nutr.* 2001; 20: 363 9.
- **7.** Afsharypour S and Sanaty F. Essential oil constituents of leaves and fruits of *Pinus eldarica* Medw. *The Journal of Essential oil Res.* 2005; 17: 327 8.
- **8.** Zargary A. Medicinal plants. 5t Edition. Tehran: Tehran University Press, Iran. 1996, pp: 9 12.
- **9.** Mozaffarian W. Tree and shrubs of Iran. First edition, Farhang Moaser Press, Iran. 1383, pp: 563 5.
- **10.** Mamedov N and Craker LE. Medicinal plants used for the treatment of bronchial asthma in Russia and Central Asia. *J. Herbs Spices Med. Plants* 2001; 8: 91 117.
- **11.** Mamedov N, Gardner Z and Craker LE. Medicinal plants used in Russia and Central Asia for the treatment of selected skin conditions. *J. Herbs Spices Med. Plants* 2005; 11: 191 222.
- **12.** Scalbert A, Manach C, Morand C, Remesy C and Jimenez L. Dietary polyphenols and the prevention of diseases. *Crit. Rev. Food Sci.*

- Nutr. 2005; 45: 287 306.
- **13.** Pandey KB and Rizvi SI. Current understanding of dietary polyphenols and their role in health and disease. *Current Nutrition & Food Sci.* 2009; 5: 249 63.
- **14.** Dogan S, Diken ME and Dogan M. Antioxidant, phenolic and protein contents of some medicinal plants. *J. Medicinal Plants Res.* 2010; 4: 2566 73.
- **15.** Groot AP, Luyken R and Pikaar NA. The cholesterol-lowering effects of rolled oats. *The Lancet* 1963; 10: 303 4.
- **16.** El-Demerdash FM, Yousef MI and El-Naga NI. Biochemical study on the hypoglycemic effects of onion and garlic in alloxan-induced diabetic rats. *Food Chem. Toxicol.* 2005; 43: 57 63.
- **17.** Causey JL. Korean pine nut fatty acids induce satiety-producing hormone release in overweight human volunteers. Paper presented at: *American Chemical Society National Meeting & Exposition* 2006; 26 30.
- **18.** Rosen P, Nawroth PP, King G, Moller W, Tritschler HJ and Packer L. The role of oxidative stress in the onset and progression of diabetes and its complications: a summary of a congress series sponsored by UNESCO-MCBN, the American Diabetes Association and the German Diabetes Society. *Diabetes Metab. Res. Rev.* 2001; 17: 189 212.
- **19.** Kumarappan CT, Thilagam E, Vijayakumar M and Mandal SC. Modulatory effect of polyphenolic extracts of Ichnocarpus frutescens on oxidative stress in rats with experimentally induced diabetes. *Indian J. Med. Res.* 2012; 136: 815 21.



- **20.** Hanhineva K, Törrönen R, Bondia-Pons I, Pekkinen J, Kolehmainen M, Mykkänen H and Poutanen K. Impact of dietary polyphenols on carbohydrate metabolism. *Int. J. Mol. Sci.* 2010; 11: 1365 402.
- **21.** Calleja MA, Vieites JM, Montero-Meterdez T, Torres MI, Faus MJ, Gil A and Suárez A. The antioxidant effect of β -caryophyllene protects rat liver from carbon tetrachloride-induced fibrosis by inhibiting hepatic stellate cell activation. *Br. J. Nutr.* 2012; 1: 1 8.
- **22.** Bolling BW, Chen CY, McKay DL and Blumberg JB. Tree nut phytochemicals: composition, antioxidant capacity, bioactivity,

- impact factors. A systematic review of almonds, Brazils, cashews, hazelnuts, macadamias, pecans, pine nuts, pistachios and walnuts. *Nutr. Res. Rev.* 2011; 24: 244 75.
- **23.** Gholamali J, Maleki M and Sirus S. Effect of Walnut Leaf, Coriander and Pomegranate on blood glucose and histopathology of pancreas of alloxan induced diabetic rats. *Afr. J. Tradit. Complement Altern. Med.* 2007; 4: 299 305.
- **24.** Rather MA, Dar BA, Dar MY, Wani BA, Shah WA, Bhat BA, Ganai BA, Bhat KA, Anand R and Qurishi MA. Chemical composition, antioxidant and antibacterial activities of the leaf essential oil of *Juglans regia* L. and its constituents. *Phytomedicine* 2012; 19: 1185 90.