Studies on GC/MS Spectroscopic Analysis of some Bioactive Antimicrobial Compounds from *Cinnamomum zeylanicum*

Uma B (Pharm.D., M.Sc.)¹*, Prabhakar K (M.Sc, Ph.D.)², Rajendran S (M.Sc., Ph.D.)³, Lakshmi Sarayu Y (M.B.B.S, M.D.)⁴

 Research scholar, Division of Microbiology, Rajah Muthiah Medial College and Hospital, Annamalainagar -608001, Tamilnadu, India
Professor, Division of Microbiology, Rajah Muthiah Medial College and Hospital, Annamalai nagar -608001, Tamilnadu, India
Professor, Division of Microbiology, Rajah Muthiah Medial College and Hospital, Annamalai nagar -608001, Tamilnadu, India
Professor, Division of Microbiology, Rajah Muthiah Medial College and Hospital, Annamalai nagar -608001, Tamilnadu, India
Head of the Department, Division of Microbiology, Rajah Muthiah Medial College and Hospital, Annamalainagar -608001, Tamilnadu, India
*Corresponding author: Research scholar, 36/6 Meikaval Street, Chidambaram – 608001, Tamilnadu, India, Tel: +919488791043 E-mail: amu_sri2003@yaho.co.in

Receive: 10 Mar. 2009

Acceptance: 27 Jul. 2009

Abstract

Background: Infectious diarrhoea is an emergent problem in both developing and developed countries for high rates of mortality in infants. In recent years, drug resistance to microbial pathogens has been commonly reported from all over the world. Medicinal plants represent a rich source of antimicrobial agents. Plants are used medicinally in different countries especially in India and are a source of many potent and powerful drugs. This study was aimed to explore the bioactive antimicrobial compounds present in *Cinnamon zeylanicum*, which is used as a spice in Indian foods.

Objective: The aim of this study was to carry out Gas chromatography and Mass spectroscopy studies and to determine the antimicrobial activity of *Cinnamon zeylanicum* extract against some common microbial pathogens isolated from diarrhoeal patients.

Methods: The extract was prepared by the method of maceration using methanol as extraction solvent. The antimicrobial activity was performed by disc diffusion method at the concentration 200, 100, 50, 25 and 10 mg/ml against *E.coli*, *Salmonella typhi*, *Salmonella typhimurium*, *Shigella dysenterae*, *Shigella flexneri*, *Pseudomonas aeruginosa* and *Klebsiella pneumonia* and the fungi *Candida albicans*, *Candida tropicalis*, *Candida krusei* which are isolated from the paediatric diarrhoeal samples. Gas chromatography and Mass spectroscopy studies were performed to identify the bioactive compounds.

Results: The methanol extract was found very sensitive against all the isolated organisms. The preliminary phytochemical screening shows the presence of carbohydrate, glycosides, phenolic compounds, tannins, terpenes and proteins. The GC/MS study on the active principles shows the presence of 38 components.

Conclusion: This study shows that the *Cinnamon zeylanicum* extract exhibit high antifungal activity as compared with antibacterial activity. The most identified compound by GC/MS to have antimicrobial property were monoterpenes, sesquiterpenes, aromatic aldehydes and ketones. Cinnamaldehyde was the major compound responsible for the antimicrobial activity.

Keywords: Antimicrobial activity, *Cinnamon zeylanicum*, Gas chromatography/Mass Spectroscopic Studies



Introduction

In developing countries like India, the majority of people living in rural areas were almost exclusively using traditional medicines in treating all sorts of disease including diarrhea [1]. Infectious diarrhea is one of the principal causes of death in the infants in developing countries particularly in malnourished and unhygienic conditions. It thus becomes important to identify and evaluate commonly available medicinal plant to treat infectious diarrhea as alternative to currently used antimicrobial drugs. Several studies have evaluated the antimicrobial properties in some traditional medicines in treating infectious diarrhea [2]. Cinnamon zeylanicum Blume (Family. Lauraceae) is a small evergreen tree native to tropical southern India and Sri Lanka., growing from sea level to nine hundred meters. Cinnamon barks are widely used as a spice. It has also been used to treat diarrhea and other problems of the digestive system and as a component of compounds used in Indian Avurvedic medicine [3]. The Cinnamon bark tea infusion was used for complaints such as a feeling of distension, flatulence, and mild cramp-like gastrointestinal disorders due to reduced production of gastric juice. Cinnamon bark is also traditionally used to treat symptoms of digestive disorders, functional asthenias, and also to facilitate weight gain [4]. Many documented publications have the antimicrobial activity of Cinnamon against different microbial spices [5-8]. The Cinnamon bark contains volatile oils (14%) of cinnamaldehyde (60%), eugenol (up to 10%) and trans-cinnamic acid (51%); phenolic compounds tannins. (41%). condensed catechins, proanthocyanidins; and monoterpenes and sesquiterpenes, (pinene); calcium-monoterpenes oxalate: gum;

mucilage; resin, starch, sugars, and traces of coumarin [9]. The previous GC/MS studies on Cinnamon bark have shown that the major cinnamaldehyde constituents were and cinnamic acids [10]. The objective of the present study was to determine the antimicrobial activity of Cinnamon bark against various diarrheal pathogens isolated from patients and to Analysis the composition of alcoholic extract of dried fruits of Cinnamon zeylanicum by GC/MS. Although a lot of work has been reported in the literature bv different workers on the chemical composition zevlanicum Cinnamon conventional methods and by GC/MS, yet it is the first time that the characterization has been carried out in Cinnamon zevlanicum by GC/MS in our region.

Material and Methods Plant Materials

Cinnamon zeylanicum dried barks were collected from Kodaikanal area in Tamilnadu, India, and were identified and confirmed the by Department of Botany, Annamalai University, Annamalainagar.

Extraction of plant materials

The barks were shade dried, powered and were extracted with methanol using a maceration process and with occasional shaking for 3 days. The extract was then filtered, dried at 50 to 60 ° and the residue was weighed and percentage yield was calculated and subjected to preliminary phytochemical analysis. The residue was dried and stored in air tight container.

Antibiotic testing of plant extract

The Micro-organism used were *E. coli*, Salmonella typhi, Salmonella typhimurium,

Shigella dysenterae, Shigella flexneri, Pseudomonas and Klesiella aureginosa pneumonia and the fungi Candida albicans, Candida tropicalis, Candida krusei which are isolated from the diarrhoeal samples from the paediatric patients. The dried extracts were reconstituted with 5% aqueous dimethyl sulphoxide (DMSO) at concentration of 500mg/ml. plant extracts were then diluted at a concentration of 200, 100, 50, 25 and 10 mg/ml in 5% aqueous DMSO. The antimicrobial assay was performed by disc diffusion method. Ciprofloxacin (10 mcg/disc) and Fluconazole (10 mcg/disc) were used as positive control for bacteria and Candida respectively and 5% DMSO impregnated disc was used as negative control. The plates were incubated at 37°C for 24 to 48 hours. The results were recorded by measuring the zones of inhibition. The experiment was performed

in triplicate and the mean values of results are given in Table 1.

GC/MS analysis of plant materials

The methanol extract was filtered with sodium sulphate [2g] and concentrated the extract to 1ml by bubbling nitrogen into the solution. The extracted material was taken for GC MS analysis. The Gas chromatography-Mass spectroscopy (Agilant 6890/Hewlettpackard 5975) was fitted with electron impact (EI) mode. The Helium was used as the carrier gas at a flow rate of 1mL/min. The temperature was programmed at 80°C for 5 min then increased to 300°C at the rate of 15°C/min. The temperature of injector and EI detector (70eV) were 280°C and 300°C, respectively. Each plant extract of 29µL was injected with a Hamilton syringe to the GC/MS manually.

	Diameter of zones of inhibition in mm							
Microorganisms	Methanol extracts in mg/ml conc.				Ciprofloxacin	Fluconazole		
	200	100	50	25	10	(5µg/disc)	(10µg/disc)	
K. pneumonia	3+	2+	1+	-	-	26	-	
Ps. aureginosa	3+	2+	1+	-	-	26	-	
E. coli	4+	3+	2+	1+	-	24	-	
S. typhimurium	4+	3+	2+	1+	-	24	-	
S. enteritidis	4+	3+	2+	1+	-	22	-	
S. typhi	4+	3+	2+	1+	-	30	-	
Sh. dysenteriae	4+	3+	2+	1+	-	28	-	
Sh. flexneri	4+	3+	2+	1+	-	24	-	
C. albicans	3+	2+	1+	-		-	20	
C. tropicalis	3+	2+	1+	-		-	20	
C. krusei	3+	2+	1+	-		-	20	

Table 1- Antimicrobial	activity of Cinnamon	i zeylanicum
------------------------	----------------------	--------------

1+ indicates zone of inhibition in average of 7 to 10 mm; **2**+ indicates zone of inhibition in average of 11 to 14 mm; **3**+ indicates zone of inhibition in average of 15 to 18 mm; **4**+ indicates zone of inhibition in average of 19 to 22 mm; **-** No activity



Results and Discussion

The methanol extract exhibits a good antimicrobial activity against all the microorganisms tested, was shown in Table 1. The zone of inhibition exhibited by the alcohol extract was comparable with standard This study shows that antibiotics. the Cinnamon zevlanicum extract exhibit high activity when compare with antifungal antibacterial activity. However the concentration of extract required for activity against Candida, Klebsiella and Pseudomonas *spp.* were very higher than the other pathogens tested.

Many studies support the above findings [5-8]. The percentage yield obtained was preliminary The phytochemical 15.4%. screening shows the presence of carbohydrate, glycosides, phenolic compounds, tannins, flavonoids and proteins. The studies on the content active principles in Cinnamon zeylanicum by GC/MS clearly shows the presence of 38 components (Table 2). The most identified compound to have antimicrobial property were monoterpenes, sesquiterpenes, aromatic aldehydes and ketones. Cinnamaldehyde was the major (68.41%),followed compound by benzaldehyde. The fragrance of plants was mainly due to presence of essential or volatile oil fractions. Cinnamon bark was rich in essential oil which possesses activity against fungi and bacteria due to the presences of antimicrobial compounds. These oils are secondary metabolites that are highly enriched in compounds like cinnamaldehyde, cinnamic acid, benzaldehyde, eugenol, benzoic acid, monoterpenes, triterpenes, and sesquiterpenes [11]. The inhibitory activity by Cinnamon extract could be due to the presence of cinnamaldehyde, benzaldehyde and terpenes. It was reported that 60% of essential oil derivatives examined were inhibitory to fungi while 30% inhibited bacteria [11]. The fungicidal effect of Cinnamon extract is due to the presence of natural products that protects the plant from various diseases and pests. The mechanisms involved cytoplasm are granulation, cytoplasmic membrane rupture inactivation and/or inhibition and of intracellular enzymes. It has been reported that lytic enzymes act on the fungal cell wall, causing breakge of β -1,3glycan, β -1,6glycan and chitin polymers [12].



No	RT	Name of the compound	Molecular	MW	Peak	Compound
		Discula[2,2,1]hantan 2,al 1,7,7	Formula		Area %	Nature
1	6.62	trimethyl-, (1S-endo)-	C ₁₀ H ₁₈ O	154	0.09	alcohol
2	6.89	Naphthalene	C10H8	128	0.04	Aromatic compound
3	7.36	Benzenepropanol	C9H12O	136	0.11	Aromatic alcohol
4	8.00	Cinnamaldehyde, (E)-	C9H8O	132	68.41	Aldehyde
5	8.48	4-(1-Hydroxyethyl)benzaldehyde	C9H10O2	150	1.54	Aromatic Aldehyde
6	9.65	Copaene	C ₁₅ H ₂₄	204	5.27	Sesquiterpene
7	9.76	Cyclohexane, 1-ethenyl-1-methyl- 2,4-bis(1-methylethenyl)-, [1S- (1à,2á,4á)]-	C ₁₅ H ₂₄	204	0.17	Sesquiterpene
8	9.98	1,4-Methano-1H-indene, octahydro- 4-methyl-8-methylene-7-(1- methylethyl)-, [1S- (1à,3aá,4à,7à,7aá)]-	C ₁₅ H ₂₄	204	0.25	Sesquiterpene
9	10.22	2H-1-Benzopyran-2-one	C9H6O2	146	5.27	Ketone compound
10	10.70	à-Caryophyllene	C ₁₅ H ₂₄	204	0.22	Sesquiterpene
11	11.16	Naphthalene, 1,2,4a,5,6,8a- hexahydro-4,7-dimethyl-1-(1- methylethyl)- (1à 4aà 8aà)-	C ₁₅ H ₂₄	204	2.99	Sesquiterpene
12	11.27	2-Propenal, 3-(2-methoxyphenyl)-	$C_{10}H_{10}O_2$	162	0.56	Aldehyde
13	11.41	Naphthalene, 1,2,3,5,6,8a-hexahydro- 4,7-dimethyl-1-(1-methylethyl)-, (1S- cis)-	C ₁₅ H ₂₄	204	3.22	Sesquiterpene
14	11.48	Naphthalene, 1,2,4a,5,8,8a- hexahydro-4,7-dimethyl-1-(1- methylethyl)-, (1à,4aá,8aà)-(ñ)-	C ₁₅ H ₂₄	204	0.38	Sesquiterpene
15	11.58	Naphthalene, 1,2,3,4,4a,7-hexahydro- 1.6-dimethyl-4-(1-methylethyl)-	C ₁₅ H ₂₄	204	1.14	Sesquiterpene
16	11.69	à-Calacorene	C ₁₅ H ₂₀	200	0.31	Hydrocarbon
17	12.24	4aH-Cycloprop[e]azulen-4a-ol, decahydro-1,1,4,7-tetramethyl-, [1aR- (1aà,4á,4aá,7à,7aá,7bà)]- (Palustrol)	C ₁₅ H ₂₆ O	222	0.37	Sesquiterpene alcohol
18	12.38	Caryophyllene oxide	C ₁₅ H ₂₄ O	220	0.11	Sesquiterpene oxide
19	12.59	3,7-Cycloundecadien-1-ol, 1,5,5,8- tetramethyl-	C ₁₅ H ₂₆ O	222	0.17	Sesquiterpene alcohol
20	12.90	Cubenol	C ₁₅ H ₂₆ O	222	0.51	-do-
21	13.06	tauMuurolol	C ₁₅ H ₂₆ O	222	1.35	-do-
22	13.23	à-Cadinol	C ₁₅ H ₂₆ O	222	0.10	-do-
23	13.50	à-Bisabolol	C ₁₅ H ₂₆ O	222	0.09	-do-
24	14.37	Tetradecanoic acid	C14H28O2	228	0.07	Fatty acid
25	14.69	Spiro[tricyclo[4.4.0.0(5,9)]decane- 10,2'-oxirane], 1-methyl-4-isopropyl- 7.8 dibudeauu (85)	C ₁₅ H ₂₄ O ₃	252	0.11	Dihydroxy compound
26	14.77	cis-9-Hexadecenal	C ₁₆ H ₃₀ O	238	0.07	Aldehyde

Table 2- Activity of Phytocomponents identified in the	plant cinnamon zeylanicum (GC-MS Study)
--	---

[Downloaded from jmp.ir on 2025-05-19]



No	RT	Name of the compound	Molecular Formula	MW	Peak Area %	Compound Nature
27	14.87	Isolongifolene, 7,8-dehydro-8a- hydroxy-	C ₁₅ H ₂₄ O	220	0.02	Hydroxy compound
28	15.09	2,5-Octadecadiynoic acid, methyl ester	C ₁₉ H ₃₀ O ₂	290	0.72	Unsaturated fatty acid ester
29	15.85	Pentadecanoic acid	C ₁₅ H ₃₀ O ₂	242	0.09	Saturated fatty
30	17.44	n-Hexadecanoic acid	$C_{16}H_{32}O_2$	256	1.63	Palmitic acid
31	17.58	Palmitic acid vinyl ester	C ₁₈ H ₃₄ O ₂	282	0.42	Ester
32	17.75	Hexadecanoic acid, ethyl ester	C ₁₈ H ₃₆ O ₂	284	0.17	Fatty acid ester
33	19.62	E-2-Octadecadecen-1-ol	C ₁₈ H ₃₆ O	268	0.56	Alkane compound
34	20.15	9,12-Octadecadienoic acid (Z,Z)-	C ₁₈ H ₃₂ O ₂	280	0.80	Linoleic acid
35	20.25	Oleic Acid	C ₁₈ H ₃₄ O ₂	282	1.84	Oleic acid
36	20.57	Octadecanoic acid	C ₁₈ H ₃₆ O ₂	284	0.33	Stearic acid
37	20.74	Stearic acid hydrazide	C ₁₈ H ₃₈ N ₂ O	298	0.14	Nitrogen compound
38	22.81	9-Octadecenoic acid (Z)-, 2,3- dihydroxypropyl ester	C ₂₁ H ₄₀ O ₄	356	0.36	Ester compound

Continue Table 2- Activity of Phytocomponents identified in the plant *cinnamon zeylanicum* (GC-MS Study)

A study shows that the Cinnamaldehyde was completely inhibiting both sensitive and resistant strain of Helicobacter pylori [13]. Terpenes are phenolic compounds that exhibit the antimicrobial activity and mostly mono and sesquiterpenes are active against bacteria, fungi, virus and protozoa [14]. The terpenes observed in this study, consist of one monoterpene (Bicyclo [2.2.1] heptan-2-ol, 1, 7-trimethyl-, (1S-endo)-), 7. and 11 sesquiterpenes. Some of them were Copaene, Cyclohexane,1- ethenyl -1- methyl - 2, 4- bis (1-methyl ethenyl)-, [1S-(1à,2á,4á)]-, and Naphthalene, 1,2,3,4,4a,7-hexahydro-1,6dimethyl-4-(1-methylethyl)-, and that exhibit the antimicrobial activity. Terpenes are found in latex and resins of some plants and physiological function of these compounds is generally believed to be a chemical in defense against certain pathogens causing human and

animal disease [15]. Their activity is a function of the lipophilic properties of the constituents terpenes, the potency of their functional groups and their aqueous solubility [16].

Summarizing these results, we conclude that the antimicrobial activity of the *Cinnamon zeylanicum* was mainly due to presence of cinnamaldehyde, benzaldehyde, and phenolic compounds monoterpenes, triterpenes and sesquiterpenes. These compounds can used as an antidiarrhoeal agent against infective diarrhoeal after successful invivo analysis.

Acknowledgement

The authors thank Mr. D. Kumaravel, senior scientist, Indian institute of crop processing technology, Thanjavur, India, for permitting to perform GC/MS studies.

Downloaded from jmp.ir on 2025-05-19

Journal of Medicinal Plans, Volume 8, No. 31, Summer 2009

References

1. Syder JD, Merson MH. The magnitude of the global problems of acute diarrheal disease a review of active surveillance data. *Bulletin of the World Health Organization* 1982; 60: 605 - 13.

2. Dey AC. Indian Medicinal Plants Used in Ayurvedic Preparations. Bishan Singh Mahendra Pal Singh, 1st ed. Dehradun: Bushan publication; 1980, pp: 136 - 7.

3. Karnick CR. Pharmacopoeial Standards of Herbal Plants, Vol. 1. Delhi: Sri Satguru Publications; 1994, pp: 94 – 5.

4. Angmor JE, Dicks DM, Evans WC, Santra DK. Studies on *Cinnamomum zeylanicum*. *Planta Med.* 1972; 21: 416 - 20.

5. Lima EO, Gompertz OF, Giesbrecht AM, Paulo MQ. In vitro antifungal activity of essential oils obtained from officinal plants against dermatophytes. *Mycoses*, 1993; 36: 333 - 6.

6. Singh HB, Srivastava M, Singh AB, Srivastava AK. *Cinnamon* bark oil, a potent fungitoxicant against fungi causing respiratory tract mycoses. *Allergy*, 1995; 50: 995 - 9.

7. Hsieh PC. Antimicrobial effect of *Cinnamon* extract. *Taiwanese J. Agri. Chem. Food Sci.*, 2000; 38: 184 – 93.

8. Quale JM, Landman D, Zaman MM. Invitro activity of *Cinnamon* zeylanicum against azole resistant and sensitive *Candida* species and pilot study of *Cinnamon* for oral candidasisis. *Am. J. Clin. Med.*, 1996; 24: 103 - 9. **9.** Duke JA. Handbook of medicinal herbs. CRC press, Inc., Boca Ratan, Fla. 1985, pp: 133 - 4.

10. Masada Y. Analysis of Essential Oils by Gas Chromatography and Mass Spectrometry. New York: Halsted (Wiley). 1976.

11. Gupta C, Amar P, Ramesh C, Kumari A. Comparative analysis of the antimicrobial activity of *Cinnamon* oil and *Cinnamon* extract on some food borne microbes. *African J. Microbiol. Res.*, 2008; 2: 247 - 51.

12. Brull S, Coote P. Preservative agents in foods: mode of action and microbial resistance mechanisms. *Int. J. Food Microbiol.*, 1999; 50: 1 - 17.

13. Nir Y, Potasman I, Stermer M, Tabak M, Neeman I. Controlled trail of the effect of *Cinnamon* extract on *Helicobacter pylori*. *Helicobacter*, 2000; 5: 94 - 7.

14. Habtemariam S, Gray AI, Waterman PG. A new antibacterial sesquiterpenes from *Premna aligotricha. J. Nat. Prod.*, 1993; 56: 3864 - 74.

15. Scortichini M, Pia Rossi M. Preliminary *invitro* evaluation of antimicrobial activity of terpenes and terpenoids towards *Erwinia amylovora* (Burrill) Winslow et al. *J. Appl Bacterial*, 1991; 71: 109 - 12.

16. Mahato SB, Sen S. Advances in triterpenoid research, 1990-1994. *Phytochemistry*, 1997; 44: 115 - 1236.

