

Research Article

Naringenin may prevent morphine-induced tolerance via inhibiting glycogen synthase kinase-3beta activity in mice

Samira Shirooie¹, Seyed Kimia Jasemi², Golale Babaei², Mohammad Reza Morovati³, Maryam Ghanbari-Movahed¹, Saman Barzegar², Mohammad Hosein Farzaei^{1,*}

¹ Pharmaceutical Sciences Research Center, Health Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran

² Student Research Committee, Kermanshah University of Medical Sciences, Kermanshah, Iran

³ Persian Medicine Department, Faculty of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

ARTICLE INFO

Keywords:

Naringenin
GSK
p-GSSer⁶⁴⁰
Tolerance to morphine
Opioids
Animal model

ABSTRACT

Background: Opioids are essential for pain treatment, but their long-term usage results in tolerance. Naringenin, a natural flavonoid found in fruit, inhibits the enzyme that causes opioid tolerance, making it effective in treating neurodegenerative diseases. **Objective:** In this study, we evaluate the role of naringenin in morphine-induced tolerance and the glycogen synthase kinase-3beta (GSK-3 β) enzyme. **Methods:** To induce tolerance to morphine in mice, repeated injections of morphine were performed for five days, and on the fifth day, a single dose of morphine was injected intraperitoneally. Pain tests (hot plate and tail flick) were performed on the first, third, and fifth days of injections. To evaluate the impact of naringenin, 45 minutes before each morphine injection, doses of 25, 50, and 100 mg/kg were administrated orally. On the last day, brain tissues were checked for biochemical factors and changes in the phosphorylation of the enzyme by the immunohistochemical method. **Results:** The results indicated that the simultaneous use of naringenin significantly increases the analgesia delay compared to the morphine group ($P < 0.001$) on the third and fifth days. Naringenin at all concentrations decreased the nitrite level caused by morphine. It showed protective effects on morphine tolerance ($P < 0.001$) in the p-GSSer⁶⁴⁰ immunohistochemical assay and reduced the phosphorylation of p-GSSer⁶⁴⁰ by GSK-3 β , activated by chronic morphine administration. **Conclusion:** Based on the results of the present study, the beneficial effect of naringenin on the GSK enzyme in morphine-induced tolerance is confirmed, but more studies are needed to investigate its impact mechanism.

1. Introduction

Opioids are potent painkillers, but they carry risks like addiction and physical dependence.

Long-term use may require higher doses due to tolerance. To manage chronic pain, opioids are used in both cancer and non-cancer patients [1].

Abbreviations: cGMP, cyclic guanosine 3',5'-monophosphate; COX2, cyclooxygenase-2; eNOS, endothelial NOS; GFAP, glial fibrillary acidic protein; GSK3, Glycogen synthase kinase 3; iNOS, inducible NOS; NO, Nitric oxide; NOS, nitric oxide synthase; nNOS, neuronal NOS; NF- κ B, nuclear factor-kappa; p-GSSer⁶⁴⁰, p-glycogen synthase

*Corresponding author: mh_farzaei@kums.ac.ir, mh_farzaei@gmail.com

doi: [10.61186/jmp.23.89.32](https://doi.org/10.61186/jmp.23.89.32)

Received 14 April 2024; Received in revised form 24 June 2024; Accepted 6 July 2024

© 2023. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>)

Prolonged administration of opioids is usually defined as treatment extending more than 3 months [2]. Opioids are drugs that attach to natural opioid receptors in the body, primarily known for pain relief. Their effects vary based on composition and receptors attached [3]. Over the past 30 years, opioid use has increased dramatically in the United States [4]. More than 16 million people worldwide are opioid-dependent [5].

Prolonged opioid use causes tolerance, less effectiveness, addiction, and abuse. This involves receptor downregulation, signaling desensitization, and upregulation of drug metabolism [6]. Glycogen synthase kinase 3 (GSK3) is a serine/ threonine kinase family member. Humans have two subtypes of GSK3, GSK3 α and GSK3 β [7]. GSK3 β regulates glycogen metabolism and cellular physiology and is involved in several cellular processes, such as embryonic development, inflammation, immune response, neuronal plasticity, apoptosis, neurodegeneration, and carcinogenesis [8]. Also, it participates in different pathological conditions, such as cancer, type 2 diabetes, Alzheimer's disease, inflammation, and bipolar disorder [9]. Chronic morphine use activates the GSK3 β enzyme. While inhibiting GSK3 β reduces morphine tolerance [10]. GSK3 β inhibitors were found to reduce the development of morphine tolerance in a dose-dependent way [11]. Liao and his colleagues have indicated frequent morphine administration induced progressive morphine tolerance in analgesic tests such as tail flick test and naloxone withdrawal syndrome in rats which was reversed by co-administration of GSK3 β inhibitors including SB216763 and SB415286 [12]. It has been shown that chronic exposure with morphine reduced Serine 9 phosphorylation of GSK3 β , the inactive form of GSK3 β , in the brainstems of mice and mice revealed morphine tolerance which was reversed

by administration of chaperon for example 4-phenylbutyric acid (PBA) [10].

Morphine use can cause oxidative stress, increasing reactive oxygen species and lead to tolerance and dependence on opioids. Nitric oxide (NO) also contributes to pain pathways and regulates opioid antinociception [13].

Naringenin is an essential phytochemical that belongs to the flavanone group of polyphenols and is mainly found in citrus fruits such as grapefruits, tomatoes, and cherries [14]. Analgesic effects of naringenin have been indicated via induction of anti-inflammatory effects and stimulating antioxidant enzymes [15]. Previous studies have pointed out that naringenin has neuroprotective and analgesic properties in neurodegenerative disorders and neuropathic pain models. It also has therapeutic potential in inflammatory pain models [16]. It has been demonstrated that naringenin improved cognitive functions in ICV-STZ (intracerebroventricular injection of streptozotocin) induced dementia via reducing the activity of GSK3 β IN both cerebral and hippocampus cortex of rats [17].

Therefore, this study examines the impact of naringenin, a citrus flavonoid with antioxidant, anti-inflammatory, and analgesic impacts, on chronic morphine-induced tolerance in mice as a GSK3 β inhibitor.

2. Materials and methods

2.1. Animal preparation

The study was conducted on male mice kept in standard cages one week before the experiment based on Dambisya et al. protocol [18]. The temperature was 22 ± 2 degrees Celsius, and there was a 12-hour light-dark period in the Faculty of Pharmacy, University of Medical Sciences, Kermanshah laboratory. During this period, the animals had free access to food and water (IR.KUMS.REC.1398.1108).

2.2. Drug injections

Morphine HCl and naringenin were purchased from Sigma-Aldrich (Buchs, Switzerland). They both were dissolved in normal saline.

Animals were divided into eight groups (7 mice in each group): (1) control group, which only received normal saline; (2) morphine group; (3) naringenin group (25 mg/kg); (4) naringenin group (50 mg/kg); (5) naringenin group (100 mg/kg); (6) morphine + naringenin group (25 mg/kg); (7) morphine + naringenin group (50 mg/kg); (8) morphine + naringenin group (100 mg/kg). The administration of morphine was done as following:

Three injections per day were done intraperitoneally for four days, including 50 mg/kg at 8:00 and 12:00 o'clock and the third injection 75 mg/kg at 17:00 o'clock. Also, a single dose of 50 mg/kg was administered on the fifth day at 8:00 am [19].

To evaluate the impact of naringenin, 45 minutes before each morphine injection, different doses of naringenin (25, 50, 100 mg/kg dissolved in normal saline) were fed to the animals by gavage. The pain tests were done on the days 1, 3 and 5. At the end of the experiment on the fifth day after pain tests, animals were euthanized, and their brains were examined for biochemical factors. Using the Griess test the amount of nitric oxide metabolites was measured to indicate the opioid tolerance. The number of changes in the downstream target protein of GSK-3 β phosphorylation, p-GS, was also collected via immunohistochemical.

2.3. Nociceptive threshold tests

Pain-related tests (hot-plate and tail-flick) were conducted on the first, third, and fifth days of injections. In the tail-flick test, a timer is started 45 minutes after the first daily morphine injection while an intense light beam is focused on the animal's tail [20]. When the animal wags

its tail, the timer stops. The measured delay time is an indicator of the pain threshold. The cut-off in this test was 10 seconds. The hot-plate device is a plate heated by electric current [21]. When the plate temperature reached 55 degrees, the mice were placed on it. The end time of the test was measured when they licked their hind legs or jumped. The test cutoff was 90 seconds.

2.4. Griess assay

At the end of study, the whole brain of 4 animals of each group were collected and after the homogenization of that in normal saline and centrifugation, the supernatant solution was used for the Griess test [22]. For protein removal with zinc sulfate, 400 μ l of the sample were mixed with 6 mg of zinc sulfate powder. For measuring the concentration of total nitrite and nitrate (Nox) in an ELISA [23] microplate, first 100 μ l of deproteinized serum, then 100 μ l of a mixture (1:1) of sulfonamide and NEDD were added and incubated for 30 minutes at 37°C. After the reaction and color formation, the light absorption resulting from the formation of the color substance was read at 540 nm via the ELISA reader, and the concentration of the samples was measured using the standard curve. Sodium nitrite was used as a standard in concentrations of 50, 25, 12.5, 6.25, and 3.125 μ M/l.

2.5. Immunohistochemistry (IHC) of p-GS^{Ser640}

Prefrontal cortex IHC of 3 animals in each group was carried out to evaluate the levels of phosphorylation of serine 640 GS as a downstream target of GSK3 β . Deparaffinization of samples was done by xylene, then washed with alcohol and rehydration by citrate buffer was done. After washing with the washing solution and then blocking, the primary p-GS^{Ser640} antibody was added to the sample. After the sample was washed the secondary antibody was added and incubated

for one hour at 37°C. Afterwards, we rinse it and add the substrate solution (this step should be done in the dark and away from light); after 5 minutes, we washed it with distilled water. Then it was dehydrated, starting with 70 degrees' alcohol until we reached absolute alcohol. For clarification, we put the sample in Xylol for 5 minutes. We mounted the lamella on the lamella with Entellan™ glue.

2.6. Statistical analyses

Ultimately, the effectiveness of different drug doses was compared to the control group. Mean \pm SEM was used using the software to determine statistical differences with One-way analysis of variance (ANOVAs) followed by post hoc Tukey's tests and two-way ANOVA repeated measure, followed by Bonferroni post hoc test were performed for statistical analysis [24].

3. Results

3.1. Inducing analgesic effect of morphine in mice

As indicated in figure 1 repeated morphine injections over five days decreased analgesic effect in mice in both hot plate test and tail flick compared to day one. There was a significant difference between third day compared to the first day ($P < 0.01$), and the fifth day compared to the first day ($P < 0.001$).

3.2. Effects of naringenin on morphine tolerance

The results depicted in figure 2 (A-F) indicate that administering multiple doses of naringenin (25, 50, and 100 mg/kg) 45 minutes before morphine injection significantly reduces tolerance to the analgesic effect in both the hot

plate and flick tail tests as compared to the control and morphine groups ($P < 0.001$ and $P < 0.001$, respectively). As seen in the figures below, naringenin alone has no analgesic effect in either test group compared to the control group.

3.3. Effects of naringenin on nitrite level

Results of the Griess test show that chronic treatment with morphine significantly ($P < 0.001$) increased the level of nitrite compared to the control group during five days (Figure 3). Naringenin injection 45 minutes before morphine injection significantly reversed the increase in nitrite level in brain tissue compared to the morphine group ($P < 0.01$ for naringenin 25 and $P < 0.001$ for naringenin 50 and 100 mg/kg). Multiple doses of naringenin alone did not produce a significant difference in nitrite levels in comparison to the control group (data not shown).

3.4. Effects of naringenin on p-GS^{Ser640} phosphorylation

The study found that continuous use of morphine resulted in a significant enhancement in phosphorylation of GS at serine 640, which is regulated by GSK-3 β , in comparison to the control group ($P < 0.001$) (Figure 4). However, pretreatment with naringenin at doses of 25, 50, and 100 mg/kg before each morphine administration significantly prevented the increase of p-GSSer640 phosphorylation ($P < 0.001$). These findings indicate that the inhibitory impact of GSK-3 β is responsible for decreasing p-GSSer640 levels after naringenin administration.




Fig. 1. After five-day administration of morphine development of analgesic tolerance was evaluate via (A: tail flick test B: hot plate test). (* $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ compared to the control group and #### $P < 0.001$ compared to the first day of the morphine group).






Fig. 2. The role of different doses of naringenin (25, 50, and 100 mg/kg) in reducing tolerance to the analgesic impact of morphine during five days of treatment. (A-C) the hot plate test and (D-F) the tail flick test. (**P < 0.001 compared to the control group and #P < 0.001 to the morphine group).



D



E




Fig. 2. The role of different doses of naringenin (25, 50, and 100 mg/kg) in reducing tolerance to the analgesic impact of morphine during five days of treatment. (A-C) the hot plate test and (D-F) the tail flick test. (**P < 0.001 compared to the control group and #P < 0.001 to the morphine group).




Fig. 3. Nitrite levels after chronic exposure to morphine and different doses of naringenin (25, 50, and 100 mg/kg) with morphine. Morphine significantly increased the nitrite level compared to the control group, which was reversed by oral administration of different doses of naringenin (25, 50, and 100 mg/kg). (**P < 0.001 vs. the control group, ##P < 0.01, and ###P < 0.001 vs. the morphine group).




Fig. 4. (a) Immunohistochemical analysis of p-GS^{Ser640} expression in the prefrontal cortex of the animal brain (400X). A, control; B, morphine; C, naringenin (25 mg/kg) + morphine; D, naringenin (50 mg/kg) + morphine; E, naringenin (100 mg/kg) + morphine. (b) Change of p-GS^{Ser640} expression in the prefrontal cortex. (***) $P < 0.001$ in comparison to the control group and ### $P < 0.001$ in comparison to the morphine group).

4. Discussion

Our study demonstrates the role of naringenin in preventing morphine-induced tolerance in mice by inhibiting GSK3 β . Intraperitoneal injection of naringenin as a GSK3 inhibitor prior to administration of morphine meaningfully decreased the analgesic tolerance development. Results showed that naringenin has significant anti-tolerant impacts against chronic morphine-induced tolerance. These findings provide strong evidence that GSK3 is involved in modifying the chronic complications induced by morphine and

that naringenin can inhibit morphine-induced tolerance.

Chronic exposure to morphine often results in the development of analgesic tolerance, requiring an increase in dosage to achieve an adequate effect. This can also increase the risk of withdrawal symptoms [25]. It has been reported that the chronic use of morphine triggers the activation of glial cells [26], upregulates inflammatory cytokines [27], and induces morphine tolerance through different mechanisms [26]. Opioids exert their action through G-protein-coupled receptors, including

μ (19), delta [11], and kappa located in the brain and spinal cord (25). Opioid analgesia is linked strongly to the the μ opioid receptors activation located in the CNS [28].

GSK3 β is a regulatory protein kinase [29]. There is a site on the GSK3 β protein known as p-glycogen synthase (p-GS Ser640), and phosphorylation of the p-GS Ser640 residue regulates GSK3 β . It is usually observed that glycogen synthase activity is suppressed via an increase in phosphorylation [30]. The role of GSK3 β in analgesia signaling, opioid tolerance, and dependence has been mentioned in numerous studies. GSK3 β inhibitors were found to reduce chronic morphine tolerance and desensitization [29]. GSK3 is involved in various cellular processes and is linked to neurodegenerative diseases like Alzheimer's [31]. Recently, various types of GSK3 inhibitors have been proposed as possible therapeutic agents for neurodegenerative conditions such as Alzheimer's disease [32]. For instance, Lithium, a non-selective GSK3 inhibitor, has been used to treat bipolar disorder [33].

Various studies have shown that naringenin, a flavonoid found in many sources, interferes with GSK3 pathways and downregulates GSK3 activity [34]. Naringenin has potent neuroprotective and antioxidant properties [35]. It has been demonstrated that citrus juices interfere with the opioids metabolism and maximize their impacts. Furthermore, oral administration of grapefruit juice increases morphine antinociception and prevents tolerance via enhancing the intestinal absorption of this agent [26]. Several studies indicate a correlation between flavonoids and morphine-tolerance. Alifarsangi et al. concluded that naringenin has a dose-dependent anti-tolerant impact toward chronic morphine usage and that the intensity of the impact is essentially dependent upon its concentration [26]. Also, Zhou et al. suggested

that the naringenin contained in grapefruit juice may be useful for chronic pain and might help in the delay of opioid-related unwanted impacts [36].

NO is a gas synthesized from arginine via nitric oxide synthase (NOS). The NOS family comprises three isoforms: endothelial NOS (eNOS), neuronal NOS (nNOS), and inducible NOS (iNOS) [37]. NO has a complex role in pain perception. NO acts as a neuromodulator and neurotransmitter in the peripheral and central nervous systems, and is involved in nociceptive processing. In the spinal cord, NO facilitates pain transmission by enhancing the release of excitatory neurotransmitters like glutamate. In the brain, NO can inhibit pain transmission by activating descending pain inhibitory pathways [38]. NO has a dual role in pain perception - it can facilitate pain transmission peripherally and in the spinal cord, but also inhibit pain at supraspinal sites in the brain. The net effect of NO on pain depends on the specific site of action and the pain modality. Modulating NO signaling may represent a potential therapeutic target for pain management [39].

NO pathways play an essential role in the development of morphine-derived dependence and affect the development of opioid analgesic tolerance and physical dependence [40] [41]. NO/cyclic guanosine 3',5'-monophosphate (cGMP) signaling pathway has been reported to participate in morphine tolerance. Morphine functions by binding to μ -opioid receptors located in the brain and causing an increase in the overexpression of both iNOS and nNOS. [37]. Repeated morphine treatment increases the secretion of NO. Therefore, inhibition of NOS and preventing NO overproduction eliminates morphine tolerance [42]. Previous study have indicated that there was a recognizable relationship between GSK3 β and the levels of NO. In this study it has been shown that GSK3 β

increased the levels of NO production by iNOS via activation of NF- κ B which was banned by GSK3 β inhibitor [43].

Our data indicates that pretreatment with naringenin before each administration of morphine effectively reduced the effect of morphine on p-GS^{Ser640}. Moreover, when morphine was repeatedly administered for five days, a significant reduction in its analgesic effect was observed in the hot plate test and tail flick as compared to the first day of administration. The prefrontal cortex (PFC) phosphorylation level of GS was measured due to the proven role of PFC in pain processing. During acute and chronic pain many changes occurs in PFC such as alterations in gene expression, neuroinflammation, glial cell activity and neurotransmitters [44].

So, in this study we have analyzed the alteration in p-GS in PFC of animals. In the study, it was found that naringenin alone did not have any pain-relieving effects compared to the group receiving morphine. However, chronic administration of naringenin resulted in the loss of tolerance to morphine analgesic effects. All doses of naringenin reduced tolerance to the analgesic impact of morphine almost equally. The thermal hyperalgesia tests showed that naringenin was able to prevent the reduction of the pain threshold compared to the morphine group. Additionally, naringenin was found to significantly decrease the amount of nitric oxide caused by chronic injection of morphine, resulting in a decrease in morphine tolerance.

Also, long-term use of morphine significantly increased p-GS^{Ser640} levels in prefrontal cortex, confirming that morphine increases GSK3 β activity. In contrast, pretreatment with naringenin before each morphine injection

significantly reduces p-GS^{Ser640} and thus reduces the high activity of GSK3 β .

Morphine analgesic dependence and tolerance are essential challenges in clinical settings, and finding novel ways seems necessary. Though, more research are required to discover the neural signaling and exact mechanisms in different brain areas involved in morphine dependence and tolerance.

5. Conclusion

It has been concluded that one of the effective ways to treat complications caused by morphine is to inhibit and modulate the activity of GSK3 β by using inhibitors of this enzyme. Naringenin, a GSK3 β inhibitor, has been found to improve the quality of long-term administration of morphine. Therefore, it can be inferred that naringenin can be used in combination with morphine for preventing tolerance to the morphine analgesic impacts. Yet, further research are needed to investigate the exact impact of naringenin on opioid-tolerance pathways.

Author contributions

SS and MGM were involved in the conception and design of the study; SS acquired and analyzed data; GB, SB and MRM contributed into drafting the article; MRM and MGM contributed to designing of study; MHF designed the study and was involved in the analysis of data and critical revision.

Conflicts of interest

The authors declare that there is no conflict of interest.

Acknowledgment

None.

References

1. Bugada D, Lorini LF, Fumagalli R and Allegri M. Genetics and opioids: towards more appropriate prescription in cancer pain. *Cancers (Basel)*. 2020; 12(7): 1951. doi: 10.3390/cancers12071951.
2. von Oelreich E, Campoccia Jalde F, Rysz S and Eriksson J. Opioid use following cardio-thoracic intensive care: risk factors and outcomes: a cohort study. *Scientific Reports* 2024; 14(1): 20. doi: 0.1038/s41598-023-50508-3.
3. Neumann VE. Comparative Analysis of Opioids as Substrates and Inhibitors of the Human Organic Cation Transporter 1 (OCT1). 2020, Dissertation, Göttingen, Georg-August Universität, 2020. doi: 10.53846/goediss-8125.
4. Singleton JH, Abner EL, Akpunonu PD and Kucharska-Newton AM. Association of nonacute opioid use and cardiovascular diseases: a scoping review of the literature. *JAHA*. 2021; 10(13): e021260. doi: 10.1161/JAHA.121.021260.
5. Dydyk AM JN, Jain NK and Gupta M. Opioid Use Disorder, Dydyk AM, Jain NK, Gupta M. Opioid Use Disorder. [Updated 2023 Apr 29]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK553166/>. 2023.
6. Zhou J, Ma R, Jin Y, Fang J, Du J, Shao X, Liang Y and Fang J. Molecular mechanisms of opioid tolerance: From opioid receptors to inflammatory mediators. *Experimental and Therapeutic Medicine*. 2021; 22(3): 1-8. doi: 10.3892/etm.2021.10437.
7. Armstrong A. Retinoic Acid, Insulin, and Cycloheximide Alter Glycogen Homeostasis in Differentiating L6 Myoblasts. 2023.
8. Kitanaka J, Kitanaka N, Tomita K, Hall FS, Igarashi K, Uhl GR and Sato T. Glycogen synthase kinase-3 inhibitors suppress morphine-induced Straub's tail via a centrally acting mechanism. *Research Square*. 2022. 1-17. doi: 10.21203/rs.3.rs-2278556/v1.
9. Sharma AK, Bhatia S, Al-Harrasi A, Nandave M and Hagar H. Crosstalk between GSK-3 β -actuated molecular cascades and myocardial physiology. *Heart Fail. Rev.* 2021; 26(6): 1495-1504. doi: 10.1007/s10741-020-09961-9.
10. Okuyama Y, Jin H, Kokubun H and Aoe T. Pharmacological chaperones attenuate the development of opioid tolerance. *IJMS*. 2020. 21(20): 7536. doi: 10.3390/ijms21207536.
11. Dorval L. FGF21 Is a Potential Therapeutic for Morphine Preference and Dependence. 2021. University of Rochester.
12. Liao W-W, Tsai S-Y, Liao C-C, Chen K-B, Yeh G-C, Chen J-Y and Wen Y-R. Coadministration of glycogen-synthase kinase 3 inhibitor with morphine attenuates chronic morphine-induced analgesic tolerance and withdrawal syndrome. *Journal of the Chinese Medical Association*. 2014; 77(1): 31-37. doi: 10.1016/j.jcma.2013.09.008.
13. Houshmand GhR, Pourasghar M, Shiran MR, Firozjae AA, Goudarzi M, Manouchehr F, Shirzad Sh, Assadpour S, Nikbakht J and Ghorbanzadeh B. Simvastatin prevents morphine antinociceptive tolerance and withdrawal symptoms through antioxidative effect and nitric oxide pathway in mice. *Behav. Brain Res.* 2021; 402: 113104. doi: 10.1016/j.bbr.2020.113104.
14. Motallebi, M., et al., Naringenin: A potential flavonoid phytochemical for cancer therapy. *Life Sci.* 2022; 305: 120752. doi: 10.1016/j.lfs.2022.120752.
15. Rehman K, Khan II, Akash MSH, Jabeen K and Haider K. Naringenin downregulates inflammation-mediated nitric oxide overproduction and potentiates endogenous antioxidant status during hyperglycemia. *J. Food*

Biochem. 2020; 44(10): e13422. doi: 10.1111/jfbc.13422.

16. Fakhri S, Sabouri S, Kiani A, Farzaei MH, Rashidi Kh, Mohammadi-Farani A, Mohammadi-Noori E and Abbaszadeh F. Intrathecal administration of naringenin improves motor dysfunction and neuropathic pain following compression spinal cord injury in rats: relevance to its antioxidant and anti-inflammatory activities. *Korean J. Pain.* 2022; 35(3): 291-302. doi: 10.3344/kjp.2022.35.3.291.

17. Yang, W., et al., Effect of naringenin on brain insulin signaling and cognitive functions in ICV-STZ induced dementia model of rats. *Neurol. Sci.* 2014; 35(5): 741-751. doi: 10.1007/s10072-013-1594-3.

18. Dambisya YM and Lee T.L. Role of nitric oxide in the induction and expression of morphine tolerance and dependence in mice. *British J. Pharmacol.* 1996; 117(5): 914-918. doi: 10.1111/j.1476-5381.1996.tb15280.x.

19. Shirooie S, Esmaeili J, Sureda A, Esmaeili N, Saffari PM, Yousefi-Manesh H and Dehpour AR. Evaluation of the effects of metformin administration on morphine tolerance in mice. *Neuroscience Letters.* 2020; 716: 134638. doi: 10.1016/j.neulet.2019.134638.

20. Pehlivan DY, Kara AY, Koyu A and Simsek F. Enhancing fentanyl antinociception and preventing tolerance with α -2 adrenoceptor agonists in rats. *Behav. Brain Res.* 2024; 457: 114726. doi: 10.1016/j.bbr.2023.114726.

21. Nakhaee S, Dastjerdi M, Roumi H, Omid Mehrpour and Farrokhfall Kh. N-acetylcysteine dose-dependently improves the analgesic effect of acetaminophen on the rat hot plate test. *BMC Pharmacol and Toxicol.* 2021; 22(4): 1-7. doi: 10.1186/s40360-020-00469-4.

22. Vargas-Maya NI, Padilla-Vaca F, Romero-González OE, Rosales-Castillo EAS, Rangel-Serrano Á, Arias-Negrete S and Franco B. Refinement of the Griess method for measuring nitrite in biological samples. *J. Microbiol. Methods.* 2021; 187: 106260. doi: 10.1016/j.mimet.2021.106260.

23. Van Regenmortel, MHV. and Dubs M.-C. Serological procedures. Diagnosis of plant virus diseases, 2019, p: 159-214.

24. Bertinetto C, Engel J and Jansen J. ANOVA simultaneous component analysis: A tutorial review. *Analytica Chimica Acta: X.* 2020; 6: 100061. doi: 10.1016/j.acax.2020.100061.

25. Fürst S, Zádori ZS, Zádor F, Király K, Balogh M, László SB, Hutka B, Mohammadzadeh A, Calabrese Ch, Galambos AR, Riba P, Romualdi P, Benyhe S, Timár J, Schmidhammer H, Spetea M and Al-Khrasani M. On the role of peripheral sensory and gut mu opioid receptors: Peripheral analgesia and tolerance. *Molecules.* 2020; 25(11): 2473. doi: 10.3390/molecules25112473.

26. Alifarsangi A, Esmaeili-Mahani S, Sheibani V and Abbasnejad M. The citrus flavanone naringenin prevents the development of morphine analgesic tolerance and conditioned place preference in male rats. *Am. J. Drug Alcohol Abuse.* 2021; 47(1): 43-51. doi: 10.1080/00952990.2020.1813296 .

27. Liu DQ, Zhou YQ and Feng Gao F. Targeting Cytokines for Morphine tolerance: A narrative review. *Curr. Neuropharmacol.* 2019; 17(4): 366-376. Doi: 10.2174/1570159X15666171128144441.

28. Listos J, Łupina M, Talarek S, Mazur A, Orzelska-Górka J and Kotlińska J. The mechanisms involved in morphine addiction: an overview. *Int. J. Mol. Sci.* 2019; 20(17): 4302. doi: 10.3390/ijms20174302.

29. Nezamoleslami Sadaf, Sheibani M, Mumtaz F, Esmaeili J, Shafaroodi H and Dehpour AR. Lithium reverses the effect of opioids on eNOS/nitric oxide pathway in human umbilical

vein endothelial cells. *Molecular Biology Reports*. 2020; 47(9): 6829-6840. doi: 10.1007/s11033-020-05740-9.

30. Jall S, Angelis MD, Lundsgaard AM, Fritzen AM, Nicolaisen TS, Klein AB, Novikoff A, Sachs S, Richter EA, Kiens B, Schramm KW, Tschöp MH, Stemmer K, Clemmensen Ch, Müller TD and Kleinert M. Pharmacological targeting of $\alpha 3\beta 4$ nicotinic receptors improves peripheral insulin sensitivity in mice with diet-induced obesity. *Diabetologia*. 2020; 63(6): 1236-1247. doi: 0.1007/s00125-020-05117-4.

31. Lin ChH, Hsieh YS, Sun YC, Huang WH, Chen ShL, Weng ZK, Lin TH, Wu YR, Chang KH, Huang HJ, Lee GC, Hsieh-Li HM and Lee-Chen GJ. Virtual screening and testing of GSK-3 inhibitors using human SH-SY5Y cells expressing Tau folding reporter and mouse hippocampal primary culture under Tau cytotoxicity. *Biomol. Ther. (Seoul)*. 2023; 31(1): 127-138. doi: 10.4062/biomolther.2022.035 .

32. Chauhan N, Paliwal S, Jain S, Verma K, Paliwal S and Sharma S. GSK-3 β and its Inhibitors in Alzheimer's Disease: A Recent Update. *Mini Rev. Med. Chem.* 2022; 22(22): 2881-2895. doi: 10.2174/138955752266220420094317.

33. Mohamadian M, Fallah H, Ghofrani-Jahromi Z, Rahimi-Danesh M, Shokouhi Qare Saadlou MS and Vaseghi S. Mood and behavior regulation: interaction of lithium and dopaminergic system. *Naunyn-Schmiedeberg's Archives of Pharmacology*. 2023; 396 (Suppl 3): 1-21. doi: 10.1007/s00210-023-02437-1.

34. Atoki AV, Aja PM, Shinkafi TS, Ondari EN and Awuchi ChG. Naringenin: its chemistry and roles in neuroprotection. *Nutr. Neurosci.* 2023; 637-666. doi: 10.1080/ 1028415X.2023.2243089.

35. Li L, Lin Zh, Yuan J, Li P, Wang Q, Cho N, Wang Y and Lin Z. The neuroprotective mechanisms of naringenin: Inhibition of apoptosis through the PI3K/AKT pathway after hypoxic-ischemic brain damage. *J. Ethnopharmacol.* 2024; 318: 116941. doi: 10.1016/j.jep.2023.116941.

36. Zhou Y, Cai S, Moutal A, Yu J, Gómez K, Madura CL, Shan Z, Pham NYN, Serafini MJ, Dorame A, Scott DD, François-Moutal L, Perez-Miller S, Patek M, Khanna M and Khanna R. The natural flavonoid naringenin elicits analgesia through inhibition of NaV1. 8 voltage-gated sodium channels. *ACS Chem. Neurosci.* 2019; 10(12): 4834-4846. doi: 10.1021/acschemneuro.9b00547.

37. Li DY, Gao ShJ, Sun J, Zhang LQ, Wu JY, Song FH, Liu DQ, Zhou YQ and Mei W. Targeting the nitric oxide/cGMP signaling pathway to treat chronic pain. *Neural Regen. Res.* 2023; 18(5): 996-1003. doi: 10.4103/1673-5374.355748.

38. Anbar M and Gratt BM. Role of nitric oxide in the physiopathology of pain. *JPSM*. 1997; 14(4): p. 225-254. doi: 10.1016/S0885-3924(97)00178-4.

39. Afzidi B, Khan H, Akkol EK and Aschner M. Pain perception and management: where do we stand? *Curr. Mol. Pharmacol.* 2021; 14(5): 678-688. doi: 10.2174/1874467213666200611142438.

40. Zamanian G, Shayan M, Rahimi N, Bahremand T, Shafaroodi H, Ejtemaei-Mehr Sh, Aghaei I, Dehpour AR. Interaction of morphine tolerance with pentylenetetrazole-induced seizure threshold in mice: The role of NMDA-receptor/NO pathway. *Epilepsy Behav.* 2020; 112: 107343. doi: 10.1016/j.yebeh.2020.107343.

41. Wolińska R, Kleczkowska P, de Cordé-Skurska A, Poznański P, Sacharczuk M, Mika J and Bujalska-Zadrożny M. Nitric oxide modulates tapentadol antinociceptive tolerance and physical dependence. *Eur. J. Pharmacol.*

2021; 907: 174245. doi: 10.1016/j.ejphar.2021.174245.


42. Hassanipour M, Rahimi N, Rajai N, Amini-Khoei H, Ejtemaei Mehr Sh, Momeny M, Heidari M and Dehpour AR. The Expression and Function of Nitric Oxide Synthase Enzyme in Atorvastatin Effects on Morphine-Induced Dependence in Mice. *Archives of Neuroscience*, 2021; 8(3): e117122. doi: 10.5812/ans.117122.

43. Cuzzocrea S, Crisafulli C, Mazzon E, Esposito E, Muià C, Abdelrahman M, Di Paola R and Thiemermann C. Inhibition of glycogen synthase kinase-3 β attenuates the development of carrageenan-induced lung injury in mice. *Br. J. Pharmacol.* 2006; 149(6): 687-702. doi:

10.1038/sj.bjp.0706902.

44. Ong, WY, Stohler CS and Herr DR. Role of the prefrontal cortex in pain processing. *Mol. Neurobiol.* 2019; 56(2): 1137-1166. doi: 10.1007/s12035-018-1130-9.

How to cite this article: Shirooie S, Jasemi SK, Babaei G, Morovati MR, Ghanbari-Movahed M, Barzegar S, Farzaei MH. Naringenin may prevent morphine-induced tolerance via inhibiting glycogen synthase kinase-3 β activity in mice. *Journal of Medicinal Plants* 2023; 23(89): 32-46. doi: 10.61186/jmp.23.89.32



مقاله تحقیقاتی

بررسی اثر احتمالی ناریتینین بر تحمل ناشی از مورفین توسط مهار گلیکوژن ستاز کیناز-۳-بتا در موش سوری سمیرا شیروئی^۱، سیده کیمیا جاسمی^۲، گلاله بابائی^۲، محمد رضا مروتی^۳، مریم قبری موحد^۱، سامان بزرگر^۲، محمد حسین فرزائی^{۱*}

^۱ مرکز تحقیقات علوم دارویی، دانشگاه علوم پزشکی کرمانشاه، کرمانشاه، ایران

^۲ کمیته تحقیقات دانشجویی، دانشگاه علوم پزشکی کرمانشاه، کرمانشاه، ایران

^۳ گروه پژوهشکی ایرانی، دانشکده پژوهشکی، دانشگاه علوم پزشکی کرمانشاه، کرمانشاه، ایران

چکیده

اطلاعات مقاله

گل و ازگان:

مقدمه: مواد افیونی برای درمان درد ضروری هستند، اما استفاده طولانی مدت از آنها منجر به تحمل دارویی می‌شود. ناریتینین، یک فلاونوئید طبیعی موجود در میوه، آنزیمی را که باعث تحمل مواد افیونی می‌شود، مهار می‌کند و آن را در درمان بیماری‌های عصبی موثر می‌سازد. هدف: در این مطالعه نقش ناریتینین در تحمل ناشی از مورفین و آنزیم گلیکوژن ستاز کیناز-۳-بتا مورد بررسی قرار گرفت. روش بررسی: برای القای تحمل مورفین، تزریق‌های مکرر مورفین به مدت پنج روز انجام شد و در روز پنجم، یک دوز مورفین به صورت داخل صافاقی تزریق شد. تست درد (هات پلیت و تیل فلیک) در روزهای اول، سوم و پنجم تزریق انجام شد. برای بررسی اثر ناریتینین، ۴۵ دقیقه پیش از هر تزریق مورفین، دوزهای ۲۵، ۵۰ و ۱۰۰ میلی گرم بر کیلوگرم به صورت خوراکی گواز شد. در روز آخر بافت‌های مغز از نظر فاکتورهای بیوشیمیایی و تغییرات فسفوریل‌اسیون آنزیم به روش ایمونو‌هیستوشیمی بررسی شدند. نتایج: نتایج نشان داد که مصرف همزمان ناریتینین به طور معنی‌داری تاخیر بی‌دردی را نسبت به گروه مورفین در روز سوم و پنجم افزایش می‌دهد ($P < 0.001$). ناریتینین در تمام غلظت‌ها باعث کاهش سطح نیتریت ناشی از مورفین شد. در روش ایمونو‌هیستوشیمی p -GS^{Ser640} اثرات محافظتی بر تحمل مورفین نشان داد ($P < 0.001$) و فسفوریل‌اسیون p -GS^{Ser640} توسط آنزیم گلیکوژن ستاز کیناز-۳-بتا را کاهش داد که با تجویز مزمن مورفین فعال می‌شود. نتیجه‌گیری: با توجه به نتایج مطالعه حاضر، تأثیر مفید ناریتینین بر آنزیم گلیکوژن ستاز کیناز در تحمل ناشی از مورفین تأیید می‌شود، اما برای بررسی مکانیسم تأثیر آن به مطالعات بیشتری نیاز است.

ناریتینین:

GSK
p-GS^{Ser640}

تحمل به مرفین:

مواد افیونی:

مدل حیوانی:

مخلف‌ها: cGMP، گوانوزین مونوفسفات حلقوی؛ COX2، سیکلواکسیژناز-۲؛ eNOS، نیتریک اکساید ستاز اندوتیالی؛ GFAP، پروتئین اسیدی رشته‌ای گلیالی؛ GSK3، پروتئین کیناز فعال شده با میتوژن؛ NOS، نیتریک اکسید ستاز القایی؛ NO، نیتریک اکسید؛ NOS، نیتریک اکسید ستاز؛ nNOS، نیتریک اکسید ستاز عصبی؛ B- κ B، فاکتور هسته‌ای تقویت‌کننده زنجیره سبک کاپا از لنفوسیت‌های B فعال شده؛ p- p -GS^{Ser640}-گلیکوژن ستاز

* نویسنده مسؤول: mh.farzaei@gmail.com, mh_farzaei@kums.ac.ir

تاریخ دریافت: ۲۶ فوریه ۱۴۰۳؛ تاریخ دریافت اصلاحات: ۴ تیر ۱۴۰۳؛ تاریخ پذیرش: ۱۶ تیر ۱۴۰۳

doi: [10.61186/jmp.23.89.32](https://doi.org/10.61186/jmp.23.89.32)

© 2023. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>)