




Institute of
Medicinal Plants

Journal of Medicinal Plants

Journal homepage: www.jmp.ir



Research Article

Antioxidant and acetylcholinesterase inhibitory activities of Indonesian cultivated medicinal plants

Sumi Wijaya*, Henry Kurnia Setiawan

Faculty of Pharmacy Widya Mandala Surabaya Catholic University Indonesia, Jalan Kalisari Selatan 1 Surabaya 60112 East Java Indonesia

ARTICLE INFO

Keywords:

Plant medicines
Acetylcholinesterase
inhibitor
Antioxidant
Persea Americana
Piper cubeba

ABSTRACT

Background: Throughout many decades, researchers all around the world still extensively search for treatment strategies for Alzheimer's disease. Dispute the small number of medicines that can be used for the treatment, the side effects, and the only symptomatic treatments of the drug, which makes the necessity of finding new sources of drugs from plant sources a critical movement. **Objective:** In this research, we investigate the potential of 33 traditional medicines plants from Indonesia for their acetylcholinesterase inhibitor and antioxidant properties. **Methods:** The inhibitory potential of acetylcholinesterase was carried out using spectrophotometry using acetylcholine as a substrate and antioxidant activity was measured using the FRAP method. **Results:** The results indicate that among the 33 plants, only seven plants with an acetylcholinesterase inhibitor (AChEI). Those plants are *Chromolaena odorata*, *Mikania scandes*, *Piper cubeba*, *Peperomia pellucida*, *Persea americana*, *Lycium barbarum* and *Phyllanthus niruri*. All the plant samples showed remarkable antioxidant potency with the range of value 4.11 – 52.65 mg/mL. **Conclusions:** *Persea americana* (Lauraceae) had the greatest AChEI with a value of 4.11 mg/mL, meanwhile *Piper cubeba* (Piperaceae) has the highest potency in scavenging free radicals with an FRAP Value of 10.89 mg/mL. There was no correlation between AChEI and Antioxidant potency.

1. Introduction

Alzheimer's disease (AD) is a neurological disorder characterized by reduced numbers of enzymes, almost 90% in the brain, used to decompose acetylcholine, shown by the gradual decline in cognitive function and memory [1, 2]. This disease is caused by the appearance of

neurofibrillary tangles and plaque neuritis [3, 4]. Currently, it is not clear what causes damage and death of neurons in AD patients. Still, several causes, such as free radicals, excessive stimulation of neurotransmitter receptors, and increased Ca²⁺, are reported to play a role in AD patients. Nowadays, about 44 million people

Abbreviations: AChEI, Acetylcholinesterase inhibitor; AD, Alzheimer's disease; AChE, Acetylcholinesterase; CO, *Chromolaena odorata*; EO, Essential Oils; IC₅₀, Inhibition Concentration; LB, *Lycium barbarum*; MS, *Mikania scandes*; PA, *Persea Americana*; PC, *Piper cubeba*; PN, *Phyllanthus niruri*; PP, *Peperomia pellucida*; TPC, Total Phenolic Content; TFC, Total Flavanoid Content

*Corresponding author: sumi@ukwms.ac.id

doi:

Received 24 June 2024; Received in revised form 30 November 2024; Accepted 30 April 2025

© 2023. Open access. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<https://creativecommons.org/licenses/by-nc/4.0/>)

suffer from AD, and the prevalence of this disease increases exponentially with age, reaching approximately 10% at age 65 and reaching nearly 50% at age 85 [3, 5, 6]. This disease is predicted to become one of the leading causes of death in the world, and it is estimated that 115.4 million people in 2050 will be affected by this disease [7, 8, 9]. Recent studies have designated that women endure AD more compared to men [3]. The previous research also makes a relation or link between AD and some other neurodegenerative diseases, such as dilated cardiomyopathy, stroke, Parkinson's disease, traumatic brain injury, and dementia caused by neuronal degradation [10, 11, 12].

Many researchers link AD to Acetylcholinesterase (AChE). AChE is an enzyme in the hydrolase group responsible for breaking down acetylcholine through carboxylic acid ester bonds. AChE also takes responsibility for cell apoptosis, catecholaminergic–cholinergic balance, inflammation, and oxidative stress [13, 14, 15]. Through the fact that AChE inhibitors such as protamine, memantine, and tacrine fail to cure the disease, only improve cognitive and memory symptoms in AD, and cause side effects such as dizziness, vomiting, gastrointestinal disturbance, nausea, diarrhea, and hepatotoxicity [15-23], this trigger the urgency to obtain candidates for acetylcholine inhibitor drugs derived from plants. Interestingly, for the last decade, a study reported that searching for AChE inhibitors from plants recorded as much as 15% of all ACh research [24]. Huperzine A, physostigmine, and galantamine are some of the AChE inhibitors derived from plants [25]. Physostigmine is isolated from *Physostigma venenosum*, a plant belonging to the Leguminosae tribe. Galanthamine is a bioactive compound that is widely found in the Amarylidaceae family.

Huperzine A, an alkaloid derived from *Huperzia serrate* (Lycopodiaceae) [3, 4, 16]. The AChE inhibitors from plant generality are alkaloid compounds such as isoquinoline, indole, quinolizidine, steroidal alkaloid, and piperine. Phenolic compounds, flavonoids, and some terpenoids were also potent AChE inhibitors (AChEI) [19, 25, 26].

In this research, a preliminary study will be conducted of several plants from families known to contain an abundance of alkaloids and polyphenols in order to obtain the candidate samples that can inhibit acetylcholine and have antioxidant activity. Several preliminary studies have shown a link between acetylcholinesterase inhibitory activity and antioxidant activity, where the presence of antioxidant compounds can protect nerves from oxidative stress so that they can slow the development of AD [16, 27].

2. Materials and methods

2.1. Sample Collection and Preparation

The selection of plant parts and the type of solvent are adjusted to the regulations for making traditional medicines in Indonesia. The plant parts collected are healthy and not mouldy. The leaves used are leaves that are neither old nor too young. The criteria for the fruit used are ripe fruit. The rhizome part is approximately three months old. The plants used were determined and stored for herbarium samples at the Widya Mandala Pharmaceutical Botany Laboratory, Surabaya. The extraction method was maceration with the solvent used is 96% ethanol or a mixture of ethanol-water (70:30). Samples were stored at -20°C until testing.

2.2. Evaluation of AChEI

The AChEI activity was tested according to the research method developed by Ali-Shtayeh *et al.* (2014). Each plant extract was prepared at a

concentrations range of 100 - 0.78 mg/ml). β -naphthyl acetate (Sigma Aldrich, Germany), AChE solution (Sigma Aldrich, Germany), and fast blue b salt (Sigma Aldrich, Germany) were used as a reagent for this method. Galanthamine hydrobromide (Sigma Aldrich, Germany) served as a positive control. The linear regression between sample concentration and the % of inhibition will be used to get IC_{50} values [28, 29].

2.3. Antioxidant assay using FRAP Method

The antioxidant activity was carried out according to the method used by Wijaya et al. (2020). The plant extract concentration used was 1.56-100 mg/mL. Trolox, Quercetin, Rutin, and Vitamin C were used as a reference antioxidant compound. The FRAP value was used to express the antioxidant activity of the sample, representing a sample concentration with an absorbance value equivalent to 1 mM of $FeSO_4$. The IC_{50} was determined by plotting the sample concentration against the value of absorbance at 593 nm [30].

2.4. Total Phenolic Content

This assay was executed according to Wijaya et al. (2020) using Folin-Ciocalteu (Sigma, Germany) and sodium carbonate (Brataco, Indonesia). The absorbance of the prepared sample mixture was read at 765 nm. Gallic acid (Sigma, Germany) was used for the standard curve, and the total phenolic content expressed of mg Gallic acid equivalents/g of the sample [30].

2.5. Total Flavonoid Content

Total flavonoid contents (TFC) in the sample extract were analyzed with aluminum chloride reagent using the colorimetric method adapted from Nurcholis et al. (2021). The amount of flavonoid was calculated using a quercetin

(Sigma, Germany) calibration curve and was expressed as mg QE/g sample [31].

2.6. Statistical Analysis

All results in this study are the average of three repetitions expressed in values \pm standard deviation (SD). Correlations between antioxidant and AChEI activity were examined using Pearson's correlation coefficient (r).

3. Results

Neurodegenerative diseases such as Alzheimer's disease have become a big concern worldwide in the last decade. Acetylcholinesterase (AChE) inhibition is one therapy to reduce the symptoms of AD, as this inhibition increases cholinergic activity in the brain. In this study, 33 plants in Indonesia were examined for their ability as acetylcholinesterase inhibitors and their antioxidant activity. The plant parts used in research vary, depending on the experiences of traditional use for those plants. Maceration with ethanol or a mixture of ethanol-water (70:30) was used, depending on the solubility and stability of the active compounds. The list of plants and the extraction method are shown in Table 1.

Table 2 shows the plant extract activity for antioxidants and AChEI. In this research, the total amount of phenols and flavonoids was also carried out to determine the correlation between antioxidant activity with the total amount of phenols and antioxidant activity with the total amount of flavonoids (Table 2 & Figure 1). Several studies have correlated the existence of a linear relationship between antioxidant and anti-cholinesterase activity. Figure 2-3 illustrates the relationship between the two in this research. Correlation data analysis was carried out using only the seven plants that provided data on antioxidant and anti-cholinesterase activity.

Table 1. The list of the plants, part of used, specimen numbers and the extraction method

No	Plants	Common Name	Part of used	Family	Specimen Number	Extraction Method
1	<i>Persea americana</i>	Avocado	Leaves	Lauraceae	WM-P-005	Maceration-Ethanol 96%
2	<i>Muntingia calabura</i>	Jamaica cherry	Leaves	Tiliaceae	WM-M-007	Maceration-Ethanol 96%
3	<i>Acalypha indica</i>	Indian acalypha	Aerial part	Euphorbiaceae	WM-A-125	Maceration-Ethanol 96%
4	<i>Anredera cordifolia</i>	Madeira vine	Leaves	Basellaceae	WM-A-126	Maceration-Ethanol 96%
5	<i>Mangifera indica</i>	Mango	Leaves	Anacardiaceae	WM-M-008	Maceration-Ethanol 96%
6	<i>Sansiviera trifasciata</i>	Snake plant	Leaves	Asparagaceae	WM-S-051	Maceration-Ethanol 96%
7	<i>Smallanthus sonchifolius</i>	Yacon	Leaves	Asteraceae	WM-S-052	Maceration-Ethanol 96%
8	<i>Phaseolus vulgaris</i>	Bean	Fructus	Fabaceae	WM-P-006	Maceration-Ethanol 96%
9	<i>Mikania scandens</i>	Climbing hempvine	Leaves	Asteraceae	WM-M-009	Maceration-Ethanol 96%
10	<i>Emilia sonchifolia</i>	Lilac tassel flower	Aerial part	Asteraceae	WM-E-023	Maceration-Ethanol 96%
11	<i>Lopholaena coriifolia</i>	Leather-leaved Fluff-bush	Aerial part	Asteraceae	WM-L-012	Maceration-Ethanol 96%
12	<i>Crassocephalum crepidiodes</i>	Fireweed	Leaves	Asteraceae	WM-C-008	Maceration-Ethanol 96%
13	<i>Synedrella nodiflora</i>	Cinderella weed	Leaves	Asteraceae	WM-S-053	Maceration-Ethanol 96%
14	<i>Chromolaena odorata</i>	Siam weed	Leaves	Asteraceae	WM-C-009	Maceration-Ethanol 96%
15	<i>Camellia sinensis</i>	Tea plant	Leaves	Theaceae	WM-C-010	Maceration – Ethanol:Water (70:30)
16	<i>Piper betle</i>	Betel pepper	Leaves	Piperaceae	WM-P-007	Maceration – Ethanol:Water (70:30)
17	<i>Guazuma ulmifolia</i>	West indian elm	Leaves	Sterculiaceae	WM-P-007	Maceration – Ethanol:Water (70:30)
18	<i>Aloe vera</i>	Aloe	Leaves	Asphodelaceae	WM-A-127	Maceration – Ethanol:Water (70:30)
19	<i>Murraya paniculata</i>	Orange jasmine	Leaves	Myrtaceae	WM-M-010	Maceration – Ethanol:Water (70:30)
20	<i>Rheum palmatum</i>	Turkish rhubarb	Root	Polygonaceae	WM-R-034	Maceration – Ethanol:Water (70:30)
21	<i>Tamarindus indica</i>	Tamarind	Pulp	Leguminosae	WM-T-005	Maceration – Ethanol:Water (70:30)
22	<i>Cassia angustifolia</i>	Senna	Leaves	Fabaceae	WM-C-011	Maceration – Ethanol:Water (70:30)
23	<i>Psidium guajava</i>	Guava	Leaves	Myrtaceae	WM-P-008	Maceration-Ethanol 96%
24	<i>Syzygium polyanthum</i>	Bay leaves	Leaves	Myrtaceae	WM-S-054	Maceration-Ethanol 96%
25	<i>Lycium barbarum</i>	Chinese wolfberry	Fruits	Solanaceae	WM-L-013	Maceration-Ethanol 96%
26	<i>Cordyline fruticosa</i>	Ti plant	Leaves	Asparagaceae	WM-C-012	Maceration-Ethanol 96%

Table 1. The list of the plants, part of used, specimen numbers and the extraction method (Continued)

No	Plants	Common Name	Part of used	Family	Specimen Number	Extraction Method
27	<i>Clerodendrum serratum</i>	Bag plant	Leaves	Verbenaceae	WM-C-013	Maceration-Ethanol 96%
28	<i>Curcuma domestica</i>	Tumeric	Rhizome	Zingiberaceae	WM-C-014	Maceration-Ethanol 96%
29	<i>Piper cubeba</i>	Tailed pepper	Leaves	Piperaceae	WM-P-009	Maceration-Ethanol 96%
30	<i>Pluchea indica</i>	Indian camphorweed	Leaves	Asteraceae	WM-P-010	Maceration-Ethanol 96%
31	<i>Phyllanthus niruri</i>	Gale of the wind	Aerial part	Euphorbiaceae	WM-P-011	Maceration-Ethanol 96%
32	<i>Peperomia pellucida</i>	Pepper elder	Aerial part	Piperaceae	WM-P-012	Maceration-Ethanol 96%
33	<i>Elephantopus scaber</i>	Elephant's foot	Leaves	Asteraceae	WM-E-024	Maceration-Ethanol 96%

Table 2. Total phenolic content (TPC), total flavonoid content (TFC), antioxidant activity and anti-cholinesterase activity

No	Samples	TPC (mg/mL)	TFC (mg/mL)	FRAP Value ¹	% IC ₅₀ ²
1	<i>Persea americana</i>	3.79 ± 0.06	2.82 ± 0.05	43.26 ± 0.05	4.11 ± 0.13
2	<i>Muntingia calabura</i>	5.57 ± 0.09	3.22 ± 0.12	14.04 ± 0.06	n.d
3	<i>Acalypha indica</i>	4.15 ± 0.11	3.01 ± 0.06	15.23 ± 0.09	n.d
4	<i>Anredera cordifolia</i>	2.10 ± 0.06	2.27 ± 0.20	n.d	n.d
5	<i>Mangifera indica</i>	5.51 ± 0.17	2.61 ± 0.06	14.53 ± 0.11	n.d
6	<i>Sansiviera trifasciata</i>	1.96 ± 0.04	2.57 ± 0.03	n.d	n.d
7	<i>Smallanthus sonchifolius</i>	2.21 ± 0.05	2.84 ± 0.05	15.37 ± 0.05	n.d
8	<i>Phaseolus vulgaris</i>	1.35 ± 0.01	1.43 ± 0.28	114.77 ± 0.24	n.d
9	<i>Mikania scandens</i>	3.69 ± 0.12	2.77 ± 0.04	24.29 ± 0.03	69.18 ± 0.09
10	<i>Emilia sonchifolia</i>	3.40 ± 0.06	2.90 ± 0.04	17.81 ± 0.04	n.d
11	<i>Lopholaena coriifolia</i>	1.08 ± 0.09	0.76 ± 0.17	53.07 ± 0.07	n.d
12	<i>Crassocephalum crepidiodes</i>	3.55 ± 0.13	2.72 ± 0.04	31.92 ± 0.05	n.d
13	<i>Synedrella nodiflora</i>	5.62 ± 0.14	2.79 ± 0.03	83.58 ± 0.16	n.d
14	<i>Chromolaena odorata</i>	7.70 ± 0.12	3.01 ± 0.03	16.82 ± 0.01	52.14 ± 0.05
15	<i>Camellia sinensis</i>	0.72 ± 0.07	1.06 ± 0.17	19.59 ± 0.17	n.d
16	<i>Piper betle</i>	0.04 ± 0.01	0.01 ± 0.00	n.d	n.d
17	<i>Guazuma ulmifolia</i>	0.03 ± 0.00	0.23 ± 0.07	n.d	n.d
18	<i>Aloe vera</i>	n.d	n.d	n.d	n.d
19	<i>Murraya paniculata</i>	n.d	0.03 ± 0.01	16.44 ± 0.01	n.d
20	<i>Rheum palmatum</i>	0.83 ± 0.01	0.10 ± 0.01	n.d	n.d
21	<i>Tamarindus indica</i>	0.74 ± 0.06	0.62 ± 0.01	n.d	n.d
22	<i>Cassia angustifolia</i>	1.65 ± 0.10	1.20 ± 0.17	14.85 ± 0.17	n.d
23	<i>Psidium guajava</i>	2.30 ± 0.05	1.99 ± 0.21	28.89 ± 0.21	n.d

Table 2. Total phenolic content (TPC), total flavonoid content (TFC), antioxidant activity and anti-cholinesterase activity (Continued)

No	Samples	TPC (mg/mL)	TFC (mg/mL)	FRAP Value ¹	% IC ₅₀ ²
24	<i>Syzygium polyanthum</i>	4.89 ± 0.06	2.41 ± 0.16	22.85 ± 0.09	n.d
25	<i>Lycium barbarum</i>	1.04 ± 0.03	1.45 ± 0.21	36.57 ± 0.04	91.27±0.07
26	<i>Cordyline fruticosa</i>	1.37 ± 0.04	2.25 ± 0.27	25.29 ± 0.03	n.d
27	<i>Clerodendrum serratum</i>	3.95 ± 0.01	2.82 ± 0.08	38.33 ± 0.03	n.d
28	<i>Curcuma domestica</i>	8.77 ± 0.04	3.06 ± 0.03	38.33 ± 0.05	n.d
29	<i>Piper cubeba</i>	5.50 ± 0.08	0.57 ± 0.09	10.89 ± 0.06	44.08 ± 0.14
30	<i>Pluchea indica</i>	2.90 ± 0.02	2.91 ± 0.05	15.78 ± 0.11	n.d
31	<i>Phyllanthus niruri</i>	3.77 ± 0.05	2.96 ± 0.02	13.16 ± 0.06	76.93 ± 0.08
32	<i>Peperomia pellucida</i>	1.88 ± 0.06	1.62 ± 0.24	52.65 ± 0.14	86.33 ± 0.15
33	<i>Elephantopus scaber</i>	6.08 ± 0.09	3.22 ± 0.07	25.42 ± 0.09	n.d
34	Galanthamine	-	0.11 ± 0.07	-	-
35	Quercetine	3.03 ± 0.04	-	-	-
36	Rutin	26.60 ± 0.13	-	-	-
37	Trolox	47.16 ± 0.09	-	-	-
38	Ascorbic Acid	1.00 ± 0.03	-	-	-

n.d: not determined. Data were acquired from three data-independent experiments, represented in value ±SD, each performed in three replicates (n = 9).

¹ FRAP value represents the concentration of samples that have an absorbance value equal to that of 1 mM Fe₂SO₄.

² IC₅₀ was the concentration of the sample that gives 50% inhibition of acetylcholinesterase




Figure 1. Correlation between (1) Total Phenolic Content and Antioxidant Activity (left); (2) Total Flavonoid Content and Antioxidant Activity (right)




Figure 2. Correlation between Antioxidant Activity and Acetylcholinesterase Activity




Figure 3. Comparison between Plant samples that have Acetylcholine esterase activity vs their Antioxidant activity (PA: *Persea americana*; MS: *Mikania scandens*; CO: *Chromolaena odorata*; LB: *Lycium barbarum*; PC: *Piper cubeba*; PN: *Phyllanthus niruri*; PP: *Peperomia pellucida*)

4. Discussion

4.1. Acetylcholinesterase inhibitory activity

Medicinal plants are widely known to play many roles in maintaining health and treating diseases in many countries. More than 200 plants (from 70 families) are estimated to be used for neurodegenerative diseases. Species from the Fabaceae, Asteraceae, and Lamiaceae families were recorded as the most contributing to this activity [24]. Ellman method was used in this research to measure acetylcholinesterase potency from 33 plants that are usually used for traditional medicines in Indonesia. The mechanism of this method is based on the ability of the sample to hydrolyze β -naphthyl acetate into naphthol and acetate, where naphthol reacts with fast blue B salts to produce a purple color. From 33 plants, there are only seven plants that give acetyl cholinesterase activity (Table 2), there are two plants from Asteraceae family (*Chromolaena odorata* (CO) and *Mikania scandes* (MS)), two plants from Piperaceae (*Piper cubeba* (PC) and *Peperomia pellucida* (PP)), and the others from Lauraceae (*Persea americana* (PA)), Solanaceae (*Lycium*

barbarum (LB)) and Euphorbiaceae (*Phyllanthus niruri* (PN)). The sequences from the highest activity to low activity were PA>PC>CO>MS>PN>PP>LB with the IC₅₀ range to 4.1099 – 91.2688 mg/mL, respectively.

Asteraceae is one of the most prominent plant families, consisting of approximately 1700 genera with more than 24,000 species. Almost all species from this family are spread worldwide, including in Indonesia [32]. Plants from this family are being used to treat various diseases, including neurodegenerative diseases. *Silybum marianum*, *Phagnalon saxatile*, *Pulicaria stephanocarpa*, *Calendula officinalis*, *Chamaemelum nobile*, and *Artemisia annua* proved to inhibit AChE [3, 33-35].

Phenolic, essential oil and terpenoids are active compounds thought to have the potential to inhibit the action of acetylcholinesterase [33, 36, 37]. Plants from the Piperaceae family, such as *Piper cubeba* and *Peperomia pellucida*, have been proven to inhibit acetylcholinesterase. Previous research reported five essential oils (EO) obtained from Piper species in inhibiting

acetylcholinesterase (AChE) with IC_{50} values range of 1.51 - 14.0 mg/mL(6). Major constituents of EO piper are caryophyllene (sesquiterpenes) and asaricin (phenylpropanoids), which are also contained in *Piper cubeba* and *Peperomia pellucida*(38,39). Piperine, piperettine, piperettyline, bavinol A, ampelopsin, and violanthin isolated from Piper species also exhibited AChE inhibitory properties(3,40–43).

Phyllanthus niruri is one of the plants belonging to the Euphorbiaceae Family. Several species belonging to this family have been investigated for the potential to inhibit acetylcholinesterase. *Jatropha gossypifolia* (leaves, stem bark, and roots), extracted in different solvents, showed that the ethyl acetate of leaf extracts exhibited the highest AChE with a value of $80.46 \pm 0.44\%$. Another study reported that *Croton socotranus* leaves and flowers extracted with ethanol showed inhibition of AChE. Another plant, *Phyllanthus acidus* (methanol extract), was reported to inhibit AChE with an IC_{50} value of 449.51 μ g/mL(3,44–46). Diterpenoid isolated from *Euphorbia fischeriana* and 4-O-methyl gallic acid from *Euphorbia schimperiana* and *Euphorbia balsamifera* are some of the promising AChE inhibitors(47,48). Previous research reported that *Persea americana* has potential as an AChEI. In this study, the parts used were leaves and seeds, where the inhibitory activity of seeds was higher when compared to leaf extract. This research also stated that secondary metabolites that play a role in this inhibitory process are saponins, alkaloids, and terpenoids. In our finding, ethanol extract of *Persea americana* (Lauraceae) inhibits the greatest activity for acetylcholinesterase compared to other plants. Another researcher stated that isolated alkaloids from three *Cryptocarya* species, which belong to this family,

can inhibit AChE. In a separate study, *Beilschmiedia pulverulenta* essential oils exhibited inhibition of around 56.5% against AChE [49]. Much research in this decade concluded that alkaloids are the most potent secondary metabolites for AChE inhibitors, and one of the families containing most alkaloids is Solanaceae. Several experiments have reported alkaloids such as indole alkaloid (uleine) and isoquinoline alkaloid as cholinesterase inhibitors [50, 51]. *Lycium barbarum*, one of the plants belonging to the Solanaceae family, exhibits weaker AChE inhibitors compared to other plants in this research. This potency is probably due to less alkaloid content in the fruits compared to the leaves [52, 53].

4.2. Antioxidant Activity

The antioxidant activity was evaluated using the FRAP method. The results (Table 2) disclose that *Piper cubeba* has the highest antioxidant activity with the FRAP Value of 4.11 mg/mL, which is even better than standard Rutin and Trolox. The order of antioxidant activity for plants that have activity as AChEI was PC>PN>CO>MS>LB>PA>PP with the range of value 4.11 – 52.65 mg/mL. According to Muchandi *et al.* (2018), the antioxidant potency of the plants used for this research indicates potent antioxidant action [54]. Literature study states that almost all plants have antioxidant potency due to the plant's activity in facing photo-oxidative stress [55]. Oxidative chain reaction termination, free radical scavenging, pro-oxidant metal ions binding, and reducing capacity are believed to be involved in the antioxidant capacity of plant extracts [56]. The active substances involved in antioxidant action were phenolic compounds, including flavonoids and tannins, and some alkaloids. The antioxidant capacity of polyphenols is due to their ability to

bind ions that trigger the formation of reactive oxygen species [57]. Existing research shows that flavonoids not only function as antioxidants by capturing free radicals or activating the secretion of superoxide dismutase and glutathione but can also contribute to treating neurodegenerative diseases [58]. Through research by Cui et al. (2020), catechin tannins and their derivatives can be used to treat neurodegenerative diseases, where this ability is related to the way and position of the binding of the hydroxyl group [58]. In this research no correlation between total phenolic content/total flavonoid content and antioxidant activity (Figure 1). Alkaloids are strong antioxidants; some of the alkaloids, such as caffeine and nicotine, have already been proved to protect neurons from oxidative stress and prevent memory loss [58, 59].

4.3. The correlation between Antioxidant Activity and Acetyl cholinesterase Activity

It has been observed in the brain of AD patients that the presence of oxidative stress and the amount of Cu^{2+} and Fe^{2+} gradually increase [16]. This condition indicated that metal accumulation can trigger AD's emergence [60]. Due to the large number of free radical products in the brains of AD patients, antioxidants are considered effective in treating this disease [4, 61]. Phenolic and flavonoid compounds are well known for their great antioxidant activities, and this activity is due to their potency to scavenge free radicals or chelating metal ions [35, 60, 62]. Despite their antioxidant abilities, the phenylchroman backbone served as the minimum structural requirement for phenol to inhibit AChE, and their specific structure can influence their inhibition strength [4]. Flavonoids have been well-known as powerful antioxidants that can reduce oxidative stress in AD patients [4]. This ability is attributed to the presence of an $-\text{OH}$ group on the side phenyl ring of the

compound [16]. Unsaturation of the C-ring number and position of hydroxyl groups, especially A5-OH, A6-OH, and A7-OH, determine the potential of flavonoids as AChEI [4, 63-65]. AChEI flavonoids are concluded to have several mechanisms of action: (1) increasing nerve transmission [64, 66]; (2) increasing the amount of acetylcholine [67]; (3) inhibition of acetylcholine hydrolysis [19]. However, the efficacy of phenolic and flavonoids as AChE inhibitor agents would depend on their bio accessibility and bioavailability since most of them are hydrophilic substances. This condition might also be one of the reasons that there is no correlation between antioxidant activity and anticholinesterase inhibitor activity (Figures 1 & 2) in this research. This fact is also supported by data showing that plants with high antioxidant activity do not always have good acetylcholine inhibitory activity (Figure 3). Even though research may not correlate terpenoids with their antioxidant activity, there is a straight relationship between monoterpenoid structure and their AChE inhibitory potency. The position of the double bond of allylic methyl group - bicyclic monoterpene hydrocarbon influences their AChEI activity [64, 65]. Since Acetylcholinesterase takes an important role in AD and has even proven to have the potency to lower stroke risk [66], the importance of continuing the research to find out the active compounds from 7 plants that have the potential as an AChEI becomes the next step project.

5. Conclusion

The research results showed that of the 33 traditional medicinal plants generally used in Indonesia, seven plants were found to have acetylcholinesterase inhibitory activity. Of these seven plants, *Persea americana* proved to have higher AChEI activity, while *Piper cubeba* had

the highest antioxidant activity. The statistical results showed no correlation between acetylcholinesterase inhibitor activity and antioxidant potency.

Author contributions

All authors have the same contribution in conducting literature studies, research and journal writing.

References

1. Moreta MPG, Burgos-Alonso N, Torrecilla M, Marco-Contelles J and Bruzos-Cidón C. Efficacy of acetylcholinesterase inhibitors on cognitive function in alzheimer's disease. Review of Reviews. *Biomedicines*. 2021; 9(11): 1689. doi: 10.3390/biomedicines9111689.
2. World Health Organization. *Towards a dementia plan: a WHO guide* [Internet]. France: WHO; 2018, 178.
3. Ahmed S, Khan ST, Zargaham MK, Khan AU, Khan S, Hussain A, Uddin J, Khan A and Al-Harrasi A. Potential therapeutic natural products against Alzheimer's disease with reference of acetylcholinesterase. *Biomedicine & Pharmacotherapy*. 2021; 139: 111609. doi: 10.1016/j.biopha.2021.111609.
4. Balkis A, Tran K, Lee YZ and Ng K. Screening flavonoids for inhibition of acetylcholinesterase identified baicalein as the most potent inhibitor. *Journal of Agricultural Science*. 2015; 7(9): 26-35. doi: 10.5539/jas.v7n9p26.
5. Niu H, Álvarez-Álvarez I, Guillén-Grima F, Aguinaga-Ontoso I. Prevalence and incidence of alzheimer's disease in Europe: A meta-analysis. *Neurología (English Edition)*. 2017; 32(8): 523-32. doi: 10.1016/j.nrleng.2016.02.009.
6. Xiang CP, Han JX, Li XC, Li YH, Zhang Y, Chen L, Qu Y, Hao C-Y, Li H-Z, Yang C-R, Zhao S-J and Xu M. Chemical composition and acetylcholinesterase inhibitory activity of essential oils from Piper species. *J. Agric. Food Chem.* 2017; 65(18): 3702-10. doi: 10.1021/acs.jafc.7b01350.
7. Fiest KM, Roberts JI, Maxwell CJ, Hogan DB, Smith EE, Frolkis A, Cohen A, Kirk A, Pearson D, Pringsheim T, Venegas-Torres A and Jetté N. The prevalence and incidence of dementia due to alzheimer's disease: a systematic review and meta-analysis. *C.J.N.S.* 2016; 43(Suppl 1): S51-82. doi: 10.1017/cjn.2016.36.
8. Prince M, Bryce R, Albanese E, Wimo A, Ribeiro W and Ferri CP. The global prevalence of dementia: A systematic review and meta-analysis. *Alzheimer's & Dementia*. 2013; 9(1): 63-75.e2. doi: 10.1016/j.jalz.2012.11.007.
9. Santos TC dos, Gomes TM, Pinto BAS, Camara AL, Paes AM de A. Naturally occurring Acetylcholinesterase inhibitors and their potential use for Alzheimer's disease therapy. *Frontiers in Pharmacology*. 2018; 9(1192): 1-14. doi: 10.3389/fphar.2018.01192.
10. Fricker M, Tolkovsky AM, Borutaite V, Coleman M and Brown GC. Neuronal cell death. *Physiol. Rev.* 2018; 98(2): 813-80. doi: 10.1152/physrev.00011.2017.
11. Ramos-Cejudo J, Wisniewski T, Marmar C, Zetterberg H, Blennow K, Leon MJ de Fossati S. Traumatic brain injury and alzheimer's disease: The Cerebrovascular link. *eBioMedicine*. 2018; 28: 21-30. doi: 10.1016/j.ebiom.2018.01.021.

Acknowledgments

This research work is fully supported by Widya Mandala Surabaya Catholic University.

Conflict of interest

The authors declare no conflict of interest.

12. Tublin JM, Adelstein JM, del Monte F, Combs CK, Wold LE. Getting to the heart of Alzheimer disease. *Circulation Res.* 2019; 124(1): 142-9. doi: 10.1161/CIRCRESAHA.118.313563.
13. Ruz C, Alcantud JL, Vives Montero F, Duran R, Bandres-Ciga S. Proteotoxicity and neurodegenerative diseases. *Int. J. Mol. Sci.* 2020; 21(16): 5646. doi: 10.3390/ijms21165646.
14. van Enkhuizen J, Janowsky DS, Olivier B, Minassian A, Perry W, Young JW and Geyer MA. The catecholaminergic–cholinergic balance hypothesis of bipolar disorder revisited. *Eur. J. Pharmacol.* 2015; 753: 114-26. doi: 10.1016/j.ejphar.2014.05.063.
15. Walczak-Nowicka ŁJ and Herbet M. Acetylcholinesterase inhibitors in the treatment of neurodegenerative diseases and the role of acetylcholinesterase in their pathogenesis. *IJMS.* 2021; 22(17). 9290. doi: 10.3390/ijms22179290.
16. Ademosun AO, Oboh G, Bello F and Ayeni PO. Antioxidative properties and effect of quercetin and its glycosylated form (Rutin) on acetylcholinesterase and butyrylcholinesterase activities. *J. Evid-Based Integr. Med.* 2015; 21(4): NP11-7. doi: 10.1177/2156587215610032.
17. Ehret MJ and Chamberlin KW. Current practices in the treatment of alzheimer disease: where is the evidence after the phase III trials? *Clin. Ther.* 2015; 37(8): 1604-16. doi: 10.1016/j.clinthera.2015.05.510.
18. Kaufmann D, Kaur Dogra A, Tahrani A, Herrmann F and Wink M. Extracts from traditional chinese medicinal plants inhibit acetylcholinesterase, a known alzheimer's disease target. *Molecules.* 2016; 21(9): 1161. doi: 10.3390/molecules21091161.
19. Khan H, Marya, Amin S, Kamal MA and Patel S. Flavonoids as acetylcholinesterase inhibitors: Current therapeutic standing and future prospects. *Biomed. Pharmacother.* 2018; 101: 860-70. doi: 10.1016/j.biopha.2018.03.007.
20. Marucci G, Buccioni M, Ben DD, Lambertucci C, Volpini R and Amenta F. Efficacy of acetylcholinesterase inhibitors in alzheimer's disease. *Neuropharmacol.* 2021; 190: 108352. doi: 10.1016/j.neuropharm.2020.108352.
21. Murata K, Matsumura S, Yoshioka Y, Ueno Y and Matsuda H. Screening of β -secretase and acetylcholinesterase inhibitors from plant resources. *J. Nat. Med.* 2015; 69(1): 123-9. doi: 10.1007/s11418-014-0859-3.
22. Yan X, Tang J, Passos C dos S, Nurisso A, Simões-Pires CA, Ji M, Lou H and Fan P. Characterization of lignanamides from Hemp (*Cannabis sativa* L.) seed and their antioxidant and acetylcholinesterase inhibitory activities. *JAFc.* 2015; 63(49): 10611-9. doi: 10.1021/acs.jafc.5b05282.
23. Zemek F, Drtinova L, Nepovimova E, Sepsova V, Korabecny J, Klimes J and Kuca L. Outcomes of alzheimer's disease therapy with acetylcholinesterase inhibitors and memantine. *Expert. Opin Drug Saf.* 2014; 13(6): 759-74. doi: 10.1517/14740338.2014.914168.
24. Masondo NA, Stafford GI, Aremu AO and Makunga NP. Acetylcholinesterase inhibitors from southern African plants: An overview of ethnobotanical, pharmacological potential and phytochemical research including and beyond Alzheimer's disease treatment. *South African Journal of Botany.* 2019; 120: 39-64. doi: 10.1016/j.sajb.2018.09.011.
25. Ranjan N and Kumari M. Acetylcholinesterase inhibition by medicinal plants: A review. *Annals of Plant Sciences.* 2017; 6(6): 1640-4. doi: 10.21746/aps.2017.06.003.
26. Ajayi OS, Aderogba MA, Obuotor EM and Majinda RRT. Acetylcholinesterase inhibitor from *Anthocleista vogelii* leaf extracts. *Journal*

- of *Ethnopharmacol.* 2019; 231: 503-6. doi: 10.1016/j.jep.2018.11.009.
- 27.** Ali Hassan H, E. Allam A, H. Abu-Baih D, A. Mohamed MF, Ramadan Abdelmohsen U, Shimizu K, Y. Desoukey S, M. Hayallah A, A. Elrehany M, M. Mohamed Kh and S. Kamel M. Isolation and characterization of novel acetylcholinesterase inhibitors from *Ficus benghalensis* L. leaves. *RSC Advances*. 2020; 10(60): 36920-9. doi: 10.1039/d0ra06565j.
- 28.** Ali-Shtayeh MS, Jamous RM, Zaitoun SYA and Qasem IB. In-vitro screening of acetylcholinesterase inhibitory activity of extracts from Palestinian indigenous flora in relation to the treatment of Alzheimer's disease. *Functional Foods in Health and Disease*. 2014; 4(9): 381. doi: 10.31989/ffhd.v4i9.149.
- 29.** Cortes N, Posada-Duque RA, Alvarez R, Alzate F, Berkov S, Cardona-Gómez GP and Osorio E. Neuroprotective activity and acetylcholinesterase inhibition of five Amaryllidaceae species: A comparative study. *Life Sciences*. 2015; 122: 42-50. doi: 10.1016/j.lfs.2014.12.011.
- 30.** Wijaya S, Nee TK, Jin KT and Wiart C. Antibacterial, Antioxidant, Anti-inflammatory, and Anti-acetylcholinesterase Activity of *Mikania scandens* (L.) Willd (Climbing Hempvine). *Asian J. Pharmacog.* 2020; 4(1): 15-24.
- 31.** Nurcholis W, Sya'bani Putri DN, Husnawati H, Aisyah SI and Priosoeryanto BP. Total flavonoid content and antioxidant activity of ethanol and ethyl acetate extracts from accessions of *Amomum compactum* fruits. *Annals of Agricultural Sciences*. 2021; 66(1): 58-62. doi: 10.1016/j.aos.2021.04.001.
- 32.** Michel J, Abd Rani NZ and Husain K. A review on the potential use of medicinal plants from asteraceae and Lamiaceae plant family in cardiovascular diseases. *Front. Pharmacol.* 2020; 11. doi: 10.3389/fphar.2020.00852.
- 33.** Barral-Martinez M, Garcia-Oliveira P, Nuñez-Estevez B, Silva A, Finimundy TC, Calhella R, Nenadic M, Sokovic M, Barroso F, Simal-Gandara J, R. Ferreira ICF, Barros L and Prieto MA. Plants of the family Asteraceae: evaluation of biological properties and identification of phenolic compounds. *Chemistry Proceedings*. 2021; 5(1): 51. doi: 10.3390/CSAC2021-10486.
- 34.** Duan S, Guan X, Lin R, Liu X, Yan Y, Lin R, Zhang T, Chen X, Huang J, Sun X, Li Q, Fang Sh, Xu J, Yao Zh and Gu H. Silibinin inhibits acetylcholinesterase activity and amyloid β peptide aggregation: a dual-target drug for the treatment of alzheimer's disease. *Neurobiology of Aging*. 2015; 36(5): 1792-807. doi: 10.1016/j.neurobiolaging.2015.02.002.
- 35.** Orhan IE, Gulyurdu F, Kupeli Akkol E, Senol FS, Arabaci Anul S, Tatli II. Anticholinesterase, antioxidant, analgesic and anti-inflammatory activity assessment of *Xeranthemum annuum* L. and isolation of two cyanogenic compounds. *Pharmaceutical Biology*. 2016; 54(11): 2643-51. doi: 10.1080/13880209.2016.1177092.
- 36.** Liu J, Hua J, Qu B, Guo X, Wang Y, Shao M and Luo Sh. Insecticidal Terpenes from the essential oils of *Artemisia nakaii* and their inhibitory effects on acetylcholinesterase. *Front. Plant Sci.* 2021; 12. 2021. doi: 10.3389/fpls.2021.720816.
- 37.** Trendafilova A, Ivanova V, Rangelov M, Todorova M, Ozek G, Yur S, Ozek T, Aneva I, Veleva R, Moskova-Doumanova V, Doumanov J and Topouzova-Hristova T. Caffeoylquinic acids, cytotoxic, antioxidant, acetylcholinesterase and tyrosinase enzyme inhibitory activities of six *Inula* species from Bulgaria. *Chemistry & Biodiversity*. 2020; 17(4): e2000051. doi: 10.1002/cbdv.202000051.

38. Marques AM and Kaplan MAC. Active metabolites of the genus *Piper* against *Aedes aegypti*: Natural alternative sources for dengue vector control. *Universitas Scientiarum*. 2015; 20(1): 61-82. doi: 10.11144/Javeriana.SC20-1.amgp.
39. Ruslan NB, Amin IM, Hasani NAH, Ahmad VN, Aris F, Khor GH. In vitro cytotoxic evaluation and apoptosis effects of dillapiole in human nasal squamous cell carcinoma. *Journal Teknologi*. 2021; 83(5): 93-9. doi: 10.11113/jurnalteknologi.v83.17077.
40. Choi SJ, Oh SS, Kim CR, Kwon YK, Suh SH, Kim JK, Park GG, Son S-Y and Shin D-H. *Perilla frutescens* extract ameliorates acetylcholinesterase and trimethyltin chloride-induced neurotoxicity. *J. Med. Food*. 2016; 19(3): 281-9. doi: 10.1089/jmf.2015.3540.
41. Dung HV, Cuong TD, Chinh NM, Quyen D, Kim JA, Byeon JS, Woo MH, Choi JS and Min BS. Compounds from the aerial parts of *Piper bavinum* and their anti-cholinesterase activity. *Arch. Pharm. Res.* 2015; 38(5): 677-82. doi: 10.1007/s12272-014-0432-3.
42. Senol FS, Ślusarczyk S, Matkowski A, Pérez-Garrido A, Girón-Rodríguez F, Cerón-Carrasco JP, den-Haan H, Peña-García J, Pérez-Sánchez H, Domaradzki K, Orhan IE. Selective in vitro and in silico butyrylcholinesterase inhibitory activity of diterpenes and rosmarinic acid isolated from *Perovskia atriplicifolia* Benth. and *Salvia glutinosa* L. *Phytochem.* 2017; 133: 33-44. doi: 10.1016/j.phytochem.2016.10.012
43. Tu Y, Zhong Y, Du H, Luo W, Wen Y, Li Q, Zhu C and Li Y. Anticholinesterases and antioxidant alkamides from *Piper nigrum* fruits. *Nat. Prod. Res.* 2016; 30(17): 1945-9. doi: 10.1080/14786419.2015.1089243.
44. Moniruzzaman Md, Asaduzzaman Md, Hossain MdS, Sarker J, Rahman SMA, Rashid M and Rahman MM. In vitro antioxidant and cholinesterase inhibitory activities of methanolic fruit extract of *Phyllanthus acidus*. *BMC Complementary and Alternative Medicine*. 2015; 15(1): 403. doi: 10.1186/s12906-015-0930-y.
45. Pisano MB, Cosentino S, Viale S, Spanò D, Corona A, Esposito F, Tramontano E, Montoro P, Tuberoso CIG, Medda R and Pintus F. Biological activities of aerial parts extracts of *Euphorbia characias*. *BioMed Res. Int.* 2016; 2016: e1538703. doi: 10.1155/2016/1538703.
46. Saleem H, Ahmad I, Shahid MN, Gill MSA, Nadeem MF, Mahmood W and Rashid I. In vitro acetylcholinesterase and butyrylcholinesterase inhibitory potentials of *Jatropha gossypifolia* plants extract. *Acta Pol. Pharm.* 2016; 73(2): 419-23.
47. Aljubiri SM, Elsalam EA, Hady FKA, Radwan MO, Almansour AI and Shaker KH. In vitro acetylcholinesterase, tyrosinase inhibitory potentials of secondary metabolites from *Euphorbia schimperiana* and *Euphorbia balsamifera*. *Zeitschrift für Naturforschung C*. 2023; 78(5-6): 209-16. doi: 10.1515/znc-2021-0178.
48. Wei JC, Zhang XY, Gao YN, Wang DD, He XL, Gao XX, Hu G-S, Wang A-H and Jia J-M. Euphorfinoids E-L: Diterpenoids from the roots of *Euphorbia fischeriana* with acetylcholinesterase inhibitory activity. *Phytochem.* 2021; 190: 112867. doi: 10.1016/j.phytochem.2021.112867.
49. Wan Othman WNN, Liew SY, Khaw KY, Murugaiyah V, Litaudon M and Awang K. Cholinesterase inhibitory activity of isoquinoline alkaloids from three *Cryptocarya* species (Lauraceae). *Bioorg. Med. Chem.* 2016; 24(18): 4464-9. doi: 10.1016/j.bmc.2016.07.043.
50. Selly JB, Abdurrouf A and Juswono UP. Effect of *Sterculia quadrifida* extract R.Br. against free radical in liver organs *Oreochromis niloticus* due to heavy metal pollution. *Natural B,*

- Journal of Health and Environmental Sciences*. 2015; 3(2): 175-81. doi: 10.21776/ub.natural-b.2015.003.02.11.
- 51.** Tuzimski T and Petruczynik A. Application of HPLC-DAD for In vitro investigation of acetylcholinesterase inhibition activity of selected isoquinoline alkaloids from *Sanguinaria canadensis* extracts. *Molecules*. 2021; 26(1): 230. doi: 10.3390/molecules26010230.
- 52.** Bendjedou H, Barboni L, Maggi F, Bennaceur M and Benamar H. Alkaloids and sesquiterpenes from roots and leaves of *Lycium europaeum* L. (Solanaceae) with antioxidant and anti-acetylcholinesterase activities. *Natural Product Research*. 2021; 35(16): 2784-8. doi: 10.1080/14786419.2019.1666386.
- 53.** Teixeira F, Silva AM, Delerue-Matos C, Rodrigues F. *Lycium barbarum* berries (Solanaceae) as source of bioactive compounds for healthy purposes: A review. *IJMS*. 2023; 24(5): 4777. doi: 10.3390/ijms24054777.
- 54.** Muchandi AA, Jadhav AS, Patil SB and Jadhav NB. Antioxidant and in vitro antidiabetic activity of methanol extract of *Piper cubeba* L. *IRJPMS*. 2018; 1(3): 1-4.
- 55.** Agati G, Brunetti C, Fini A, Gori A, Guidi L, Landi M, Sebastiani F and Tattini M. Are flavonoids effective antioxidants in plants? Twenty years of our investigation. *Antioxidants*. 2020; 9(11): 1098. doi: 10.3390/antiox9111098.
- 56.** Kindl M, Blažeković B, Bucar F and Vladimir-Knežević S. Antioxidant and anticholinesterase potential of six *Thymus* species. *Evidence-Based Complementary and Alternative Medicine*. 2015; 2015:e403950. doi: 10.1155/2015/403950.
- 57.** Mervić M, Bival Štefan M, Kindl M, Blažeković B, Marijan M and Vladimir-Knežević S. Comparative antioxidant, anti-acetylcholinesterase and anti- α -glucosidase activities of mediterranean *Salvia* species. *Plants*. 2022; 11(5): 625. doi: 10.3390/plants11050625.
- 58.** Cui X, Lin Q and Liang Y. Plant-derived antioxidants protect the nervous system from aging by inhibiting oxidative stress. *Frontiers in Aging Neuroscience*. 2020; 12. doi: 10.3389/fnagi.2020.00209.
- 59.** Kaster MP, Machado NJ, Silva HB, Nunes A, Ardais AP, Santana M, Baqi Y, E Müller C, Lúcia S Rodrigues A, O Porciúncula L, Fan Chen J, R Tomé Â, Agostinho P, M Canas P and A Cunha R. Caffeine acts through neuronal adenosine A2A receptors to prevent mood and memory dysfunction triggered by chronic stress. *Proc. Natl. Acad. Sci. USA*. 2015; 112(25): 7833-8. doi: 10.1073/pnas.1423088112.
- 60.** Bush AI. The metal theory of Alzheimer's disease. *J. Alzheimers Dis*. 2013; 33(Suppl 1): S277-281. doi: 10.3233/JAD-2012-129011.
- 61.** Lee JP, Kang MG, Lee JY, Oh JM, Baek SC, Leem HH, Park D, Cho M-L and Kim H. Potent inhibition of acetylcholinesterase by sargachromanol I from *Sargassum siliquastrum* and by selected natural compounds. *Bioorganic Chem*. 2019; 89: 103043. doi: 10.1016/j.bioorg.2019.103043.
- 62.** Kumar S and Pandey A. Phenolic content, reducing power and membrane protective activities of *Solanum xanthocarpum* root extracts. *Vegetos: An International Journal of Plant Research*. 2013; 26(1): 1-7. doi: 10.5958/j.2229-4473.26.1.043.
- 63.** Cao J, Xia X, Chen X, Xiao J and Wang Q. Characterization of flavonoids from *Dryopteris erythrosora* and evaluation of their antioxidant, anticancer and acetylcholinesterase inhibition activities. *Food and Chem. Toxicol*. 2013; 51: 242-50. doi: 10.1016/j.bioorg.2019.103043.
- 64.** Ding X, Ouyang MA, Liu X and Wang RZ. Acetylcholinesterase inhibitory activities of flavonoids from the leaves of *Ginkgo biloba*

- against *Brown planthopper*. *J. Chem.* 2013; 2013(1): e645086. doi: 10.1155/2013/645086.
- 65.** Dzoyem JP, Nkuete AHL, Ngameni B and Eloff JN. Anti-inflammatory and anticholinesterase activity of six flavonoids isolated from *Polygonum* and *Dorstenia* species. *Arch. Pharm. Res.* 2017; 40(10): 1129-34. doi: 10.1007/s12272-015-0612-9.
- 66.** Rhee JS, Kim BM, Jeong CB, Park HG, Leung KMY, Lee YM and Lee J-S. Effect of pharmaceuticals exposure on acetylcholinesterase (AChE) activity and on the expression of AChE gene in the monogonont rotifer, *Brachionus koreanus*. *Comp. Biochem. Physiol. C: Toxicol. Pharmacol.* 2013; 158(4): 216-24. doi: 10.1016/j.cbpc.2013.08.005.
- 67.** Chandar NB and Ganguly B. A first principles investigation of aging processes in soman conjugated AChE. *Chemico-Biological Interactions.* 2013; 204(3): 185-90. doi: 10.1016/j.cbi.2013.05.013.

How to cite this article: Wijaya S, Setiawan HK. Antioxidant and acetylcholinesterase inhibitory activities of Indonesian cultivated medicinal plants. *Journal of Medicinal Plants* 2025; 24(93): 12-26.

doi: