# A Review on Pharmacological, Cultivation and Biotechnology Aspects of Milk Thistle (Silybum marianum (L.) Gaertn.)

Qavami N (Ph.D. Student), Naghdi Badi H (Ph.D.), Labbafi MR (Ph.D.), Mehrafarin A (Ph.D.)\*

Cultivation & Development Department of Medicinal Plants Research Center, Institute of Medicinal Plants, ACECR, Karaj, Iran

\* Corresponding author: Medicinal Plants Research Center, Institute of Medicinal Plants, Iranian Academic Center for Education, Culture and Research (ACECR),

P.O.Box: 33651/66571, Karaj, Iran

Tel: +98-26-34764010-9, Fax: +98-26-34764021

Email: A.Mehrafarin@gmail.com

Received: 4 May 2013 Accepted: 9 July 2013

#### **Abstract**

Milk thistle (Silybum marianum (L.) Gaertn.) is one of the valuable medicinal plants which used in the treatment of liver disorders. The major active constituents in this plant are flavonolignans, collectively known as silymarin which is a mixture of three isomer silybin, silydianin and silycristin. Its therapeutic properties are due to the presence of silymarin. The seeds contain the highest amount of silymarin, but the other plant parts have less amount of this compound. The silymarin content in fruits depends on milk thistle variety and geographic and climatic condition. In this review, we summarized the accomplished investigations on aspects of medicinal, cultivation, biology and biotechnology of milk thistle.

Keywords: Silybum marianum, Silymarin, Silybin, Flavonolignan, Cultivation and Medicinal properties



### Introduction

thistle (Silybum marianum (L.) Gaetrn) has been used since the time of ancient to treat a range of liver and gallbladder disorders (hepatitis, cirrhosis and jaundice) and to protect liver damage due to alcoholism, pharmaceutical drugs, and chemical pollutants [1, 2, 3]. Historically, milk thistle was grown in Europe as a food source. The roots can be eaten (soaked overnight to remove the bitterness); flower receptacle eaten like artichokes; the leaves eaten as a spinach substitute: the stalks eaten like asparagus [4, 5] and roasted fruits eaten as a coffee substitute [1, 4] or mixed with sea salt for use as a condiment [2, 4]. In parts of England and Scotland, the leaves were used extensively as food for cattle and horses. In some places, it cultivated as a garden ornamental [6] on rocky or sandy soils [7].

The ancient documents indicated that milk thistle was used in the Mediterranean area about 2000 years ago [1, 3]. Ancient Greek and Roman physician and herbalists were among the earliest people to use and write about milk thistle [3, 8]. Theophrastus called it under the name of "Pternix", Pliny the Elder called it sillybum and expressed that the juice of this plant mixed with honey was desirable for "carrying off bile" and Dioscorides called "Sillybon". Historical references particularly abundant in herbals of the middle ages [1, 3] including commendation of John Gerard (1545-1612) for expelling melancholy (depression, black bile), usage of roots, herbs and leaves of this plant for swelling and erysipelas [3, 7, 9]. By the 19<sup>th</sup> century, American Eclectic physicians used milk thistle for varicose veins and liver, spleen and kidney disorders [2] and in the mid-19<sup>th</sup> century, use of the fruit for treatment of liver diseases was revitalized by the German physician Rademacher [1, 7, 9, 10].

The biological and pharmacological properties of milk thistle are attributed to a flavanolignan complex, silymarin, which was first isolated from the fruits (achenes), (sometimes mistakenly called seeds) in 1968 [7, 10, 11, 12]. The silymarin mixture is predominantly composed of silvbin (also called silibinin) [13] (30–50 %) with varying percentages of isosilybin, silychristin and silydianin [1, 2, 3, 14, 15]. Apart from silymarin and other flavonolignans, 20 – 30% of the fruit is composed of fatty acids; 25 -30% protein; 0.038% tocopherol; 0.63% sterol and some compounds such deoxyflavanolignans mucilage [1, 2, 15, 16, 17, 18].

# Name of the Herb Common Names

Milk thistle is the most well-known English common name for this species and other names including Holy thistle (not to be confused with blessed thistle, *Cnicus benedictus*) [2], Mary thistle, St. Mary's thistle, Marian thistle, Lady's thistle, Christ's crown, Venus thistle, Heal thistle, Variegated thistle, Pig leaves, Royal thistle, Snake milk, Sow thistle and Wild artichoke [1, 3, 7]. In different parts of the world, the plant is known by various regional names such as:


**Persian:** Mary thiqal [19], Khar mariam [20, 21]

**Arabic:** Akub, Shawk ed diman, Shawk en nassara and Hharshaf barri [22, 23]

French: Chardon-Marie, Chardon argente,

Artichaut sauvage [22]

**Germany:** Mariendistel [1, 22]



Chinese: Shui Fei Ji [5]

Spanish: Cardo mariano and Cardo lechero [7]

#### **Botanical name**

Silybum marianum (L.) Gaertn (Syn. Carduus marianum L.)

Silybum is the name Dioscorides gave to edible thistle and marianum comes from the legend that the white veins running through the plant leaves were caused by a drop of the Virgin Mary's milk [3, 17].

Pharmacopeia name: Cardui mariae fructus [24]

#### Taxonomy

Kingdom Plantae – Plants

Subkingdom Tracheobionta – Vascular plants Superdivision Spermatophyta – Seed plants Division Magnoliophyta – Flowering plants

Class	Magnoliopsida – Dicotyledons
Subclass	Asteridae
Order	Asterales
Family	Asteraceae - Aster family
Genus	Silybum Adans Milk thistle
Species	Silybum marianum (L.) Gaertn.
	[6, 24]

Milk thistle is one of the most important medicinal members of Asteracae family [3]. The genus contains two species: *S. marianum* (L.) Gaertn, with variegated leaves, and *S. eburneum* Coss. Et Durieux, with totally green leaves, but genetic investigations of these two species showed that they are only variants [7, 25]. A morphological description of the plant from different references is presented in Table 1 and Figure 1.

Table 1- Morphological characteristics of milk thistle

Row	Morphological characteristics	<b>Description, Color and Texture</b>	Dimensions
1	Plant habit	High, erect [32, 33]	-
2	Stem	Stout, rigid, glabrous or slightly downy and not spiny, branched or unbranched [29, 32, 36]	3 m [34], 40 – 200 cm [28, 32], 200 – 250 cm [35]
3	Leaf	Alternate, glossy, dark green with milk-white veins running throughout [2, 3] Basal leaves: alternate, large, deeply lobed [2] and glabrous with spiny margins [28] Stem leaves: alternate and smaller, clasp the stem [32], not quite as lobed [2]	Length: 75cm Width: Up to 30 cm [6]
4	Spines	Woody [35]	Spines of leaves: 3 - 4.5 cm, Spines of bracts: 1.9 - 5 cm [3]
5	Root	One long taproot [3]	_
6	Receptacles	Including rows of broad, leathery bracts that are tipped with very stiff spines (1.9 - 5 cm) long and fringed with smaller spines [2]	1.5 - 1.9 cm



**Table 1- Continued** 


Row	Morphological	Description, Color and Texture	Dimensions
Row	characteristics	Description, color and Texture	Difficusions
7	Inflorescence	Large and round capitula, solitary at the apex of the stem or its branches, surrounded by thorny bracts [2, 36]	Diameter of flower head: about 5 cm
8	Florets	Tubular, hermaphrodite [37]	13 - 25 mm [4]
9	Number of achenes	<del>-</del>	150 per capitulum, 6000 per plant [3]
10	Seed	Heavy, flat, smooth, and shiny, achene, with a white, silky pappus and color ranging from black to brown [33], glossy brown to black, with a cocoalike odor and oily taste [3]	Broad: 3 mm and thickness: 1.5 mm, length: 6 – 8 mm [26], pappus scales 15 –20 mm [4]



Figure 1- Milk thistle plant [30]

In general, two types of *S. marianum* occur in some area of Asia including purple flowers and white flowers [26]. A single capitulum can produce up to 200 florets with color ranging from magenta to purple [26]. In cut transversely, the fruit shows a narrow, brown outer area and two large, dense, white oily cotyledons [27]. Also the fruits are an

elaiosome, fleshy structure on fruit that is rich in lipid and is attractive to ants and thus aids seeds dispersal [28]. A significant dependence between the shapeliness of the fruits and the mass of 1000 fruits was indicated that biggest fruits had the highest mass and best vigour [29]. Seeds in the secondary flower heads had



a conspicuously lower weight, when compared to the primary ones [28].

The plant size at the first bloom was reduced by postponing the sowing period from October to February. The same decreasing trend was observed in the number of flower heads and in the number of days required for the first bloom [28].

#### **Origin**

Milk thistle is indigenous to the Mediterranean region and is widespread in Central Europe, Central and Western Asia, North Africa, North and South America and Southern Australia [2, 27]. The plant was carried to North America by European colonists during the 19<sup>th</sup> century and is now naturalized in the United States and South America, Australia, China and Central Europe [7].

#### Milk thistle in Iran

Milk thistle is commonly found in the provinces of Mazandaran, Gilan, West and East Azarbaijan, Kermanshah, Khuzestan, Fars and Bushehr [21]. In one of the experiments, Shokrpour et al. [31] compared milk thistle accessions coming from some provinces of Iran for quantitative and qualitative features (Table 2). These results showed considerable variation among the studied genotypes for

the measured attributes.

#### **Ecology**

Milk thistle adapts with different conditions climates. It can be cultivated in northern climates such as Canada as well as in southern and arid conditions [11], (Table 3) because it is hardy and adaptable [6]. Morazzoni and Bombardelli (1995), claimed that the highest content of silybin, the main component of silymarin, is found in plants from subtropical climates and not from moderate ones, because higher temperatures seem to enhance the accumulation of that compound [37, 38]. Milk thistle grows well and has good yield on different soil types [37, 38, 39]. Milk thistle is normally considered to be a weed that grows along roadsides and on wastelands [32] and reported as being a noxious weed in several countries because it competes with crops both for water and for nutrients [3]. Milk thistle proliferates best in nitrogen-rich media such as dairy yards, chicken coop waste, garbage dumps and abandoned agricultural fields [38].

The silymarin content in fruits depends on the milk thistle variety and geographic and climatic conditions in which they grow. However, the highest content of silybinin as the main component of silymarin, is found in subtropical climates rather than from temperate climates [21, 38].

Table 2- Some characters of Iranian milk thistle accessions [31]

Table 2- Some characters of framan milk thistie accessions [31]		
Characters	Lowest - Highest	Unit
Capitulum per plant	11.85 (Dezfoul) - 26.9 (Ramhormoz)	Number
1000 seed weight	15.131 (Gharaghieh) - 22.73 (Dezfoul)	g
Stem height	131.8 (Hamidieh) - 166.475 (Gharaghieh)	cm
Capitulum diameter	3.775 (Andimeshk) - 4.48 (Jolgeh khalaj)	cm
Seed yield	961.9 (Hamidieh) - 2239.7 (Parsabad)	kg ha <sup>-1</sup>
Seed weight per capitulum	0.970 (Hamidieh) - 1.89 (Dezfoul)	g
Seed per capitulum	51.64 (Hamidieh) - 95.89 (Naharkhoran)	Number



Table 3 - Some ecological factors of milk thistle growth

Factor	Descriptions and reports
Climate	<ul> <li>Adapts perfectly to different conditions of moderate subtropical climates</li> <li>Cool wetter conditions probably prolong the vegetative phase, resulting in fewer flowers in the seed dehiscing development stage at harvest [41]</li> </ul>
Altitude	<ul> <li>Hilly environments produced a higher number of secondary flower heads compared with the plane site [28]</li> <li>700-1100 m [38]</li> <li>In India in 1800 - 2400 m [6]</li> <li>250-2400 m [36]</li> <li>In Iran, 0-420 m [40]</li> </ul>
Soil	<ul> <li>Grows well and has good yield on different soil types [39, 41, 42]</li> <li>Deeper clay soil and very good supply of nutrients [43, 44]</li> <li>Sandy soils to much heavier clay soil [33]</li> <li>Growing well in soils with a pH of 5.5 – 7.6 [33]</li> </ul>


Milk thistle is a nitrate accumulator and can be lethal when livestock ingest the plant, particularly in the early wilting stage [40].

#### **Growing period**

Milk thistle grows as a winter annual or biennial herb, depending on climate. Germination occurs in autumn and spring [7]. Studies have shown that milk thistle seed germination is affected by light temperature conditions [32]. Fresh milk thistle seeds seem to need an after-ripening period and germinate better at low temperatures compare to high temperatures. The seeds remain viable for 9 years or more. Young (1978) expressed that the incubated seeds in higher temperatures need for the longer afterripening period [45]. Mel'nikova (1983) reported that the minimum and maximum germination temperatures constant S. marianum are 10 and 35°C, and optimum germination occurs at 20 - 25°C [45]. Ghavami and Ramin (2007) reported that the percentage of germination at 15°C was higher than in 25 and 35°C [46].

After seedling establishment, milk thistle overwinters as a rosette and in the meanwhile basal leaf number were increasing. In late winter and early spring, milk thistle enters into the flowering stage when it receives a stimulus from low temperatures. Flowers anthesis occure from April - May. The achenes (fruits) are ripe in July. The whole growth period of milk thistle was 125 - 140 day, which could be divided into seedling stage (15 – 20 day), vegetative stage including two stages: rosette stage and stem elongation (45 - 60 day), flowering stage, fruit bearing stages and withering stages [7].

Within a capitulum, anthesis usually lasted five days. Ripe fruits were released about 17 days later. Individual plants had the potential to produce an average of 55 capitulum [47]. A single seed head can produce around 100 - 190 seeds [39]. Flavonolignan accumulation in seeds depends on the stage of flower development and is maximum at the late flowering time [39].



# **Cultivation practices**

Cultivation practices such as sowing date, fertilization and irrigation are important management factors in the production of all medicinal plants. In milk thistle, increased yield of the fruits along high silymarin content is an important aim for its cultivation. Omer and Ibrahim (1995) reported the nitrogen and especially potassium fertilization and plant spacing affected fruit yield and content of flavonolignans milk thistle of Andrzejewska and Sadowska (2007), showed that soil fertilization positively affected milk thistle fruit yield, silymarin content and proportion of unsaturated fatty acids [49]. However, soil rich in chemical fertilizers (nitrogen in particular) may have an adverse effect on the plant [41, 50]. Omidbaige (1991), showed that nitrogen fertilization had a significant effect on growth parameters of milk thistle. Nitrogen fertilization stimulates the growth of this plant but had significant negative impact on the silymarin and silybin content of the fruits [35]. In areas with high rainfall, nitrogen fertilizer can be applied regularly throughout the life cycle of the crop because it leaches from the root zone [33]. Milk thistle responds highly to moisture conditions. Both water excess and deficit inhibit silymarin accumulation. The highest silymarin level was recorded in plants grown at 60% field capacity [51]. Hassan et al. (1999) stated that irrigation and nitrogen fertilization (360 kg ha<sup>-1</sup>) produced the most vegetation and the greatest number of flower heads per plant, but the maximum yield of fruits occurred when the plants had a moderate level of irrigation (60% of field capacity) combined with a suitable level of nitrogen fertilization [48].

The number of lateral branches produced on milk thistle plant depends on the density of plants and climate conditions. Milk thistle planted at the low density of 10 plants per square meter, formed 10 – 16 lateral branches. Also, silymarin content was affected by row spacing. Narrow row spacing of 25 cm increased the friut yield, but reduced oil and silymarin content compared to plants grown in rows 50 cm apart. Delayed sowing date causes the plant to reach the reproductive phase higher temperatures, quickly at decreases the fruit yield but content of silymarin increases [52, 53, 54, 55, 56]. In summary, agronomic practices of milk thistle references and relevant for additional information reported in Table 4.

# **Biotechnology**

Silybum marianum (L.) Gaertn. is a diploid species with 2n = 34 chromosomes. Five of these chromosomes are of types "metacentric", "sub-metacentric" and "acrocentric" [57, 58, 59]. Genetic improvements in milk thistle can only be achieved through clear understanding of the plant's behavior and the amount of variability presented in wild populations (ecotypes). [28, 33]. Silymarin samples from native ecotypes had lower quantities of silvbinin as compared to that of silymarin from cultivated ones, but they had higher amounts of other compounds such as silychristin, silydianin and isosilybinin [60, 61].

The breeding studies for this plant are very low due to strong thorny stem, spiked leaves, flowers tipped with stiff spines [62]. In some countries such as Germany, a few varieties of milk thistle have been bred. Cultivar Silyb, was bred in the early 80's [41]. In the 1990s,



Table 4 - Some of milk thistle cultivation practices

Practice	Descriptions and reports
Soil preparation	The soil usually plows to 25 – 30 cm depth [36]. Before the sowing, rotary hoe
	cultivation is used to prepare a good seed-bed [33]
Sowing depth	1 – 1.5 cm [33], 3 cm [36]. Plant emergence was strongly reduced from a depth of
	more than 3 cm [32]
Methods of sowing	Directly seeded in soils [33]
Spacing of sowing	Row spacing is usually $40 - 75$ cm, with $20 - 30$ cm between plants in the row [33, 37]
	Rows 60 cm [36]
	In Egypt 50 cm between hills [49]
	In Spain, $40 - 50$ plants per square meter [37]
	In Germany, 20 – 30 plants per square meter [55]
Plant density	In warmer climates it is usually recommended that plants be planted at a much
	lower density than in moderate climates [49]
	Row spacing is usually $40 - 75$ cm, with $20 - 30$ cm between plants in the row [49]
Seed rate	In Germany at 20 – 30 plants per square meter [38]
	In Poland, $15 - 20 \text{ kg ha}^{-1}$ of seed [36]
	Increasing the sowing rate of seed from 12 to 24 kg ha <sup>-1</sup> resulted significant
	increase in achene yield [36]
Seed yield	Seed yield of plants cultivated in clay soil was superior to that of plants on sandy
	soil in both seasons [56]
	In Poland, especially on good soils the average yields are $1000 - 1700 \text{ kg ha}^{-1}$ [49]
Time of Sowing	In autumn and spring [29]
	In Italy sown in late autumn [41]
	In Iran, September [34]
	In New Zealand, in late summer [25]
Irrigation	Drought resistant and normal rainfall will often suffice [36]
	In a Mediterranean environment, under severe drought conditions, the crops should
	be irrigated during seed growth and filling [33]



**Table 4 - Continued** 

Practice	Descriptions and reports
Fertilizer	Low to moderate [33]
	Potassium and phosphorus are generally applied pre-sowing
	In Poland: 50 kg ha <sup>-1</sup> N, 30.5 kg ha <sup>-1</sup> P and 58 kg ha <sup>-1</sup> K before sowing [36]
	In Bulgaria, 49.5 kg ha <sup>-1</sup> N, 138 kg ha <sup>-1</sup> P <sub>2</sub> O <sub>5</sub> and 150 kg ha <sup>-1</sup> K <sub>2</sub> O to the soil before sowing [43]
	In Czech Republic: 45 - 60 kg ha <sup>-1</sup> N, 17.5 kg ha-1 P and 33.2 kg ha <sup>-1</sup> K
	In Egypt: 120 - 240 kg ha <sup>-1</sup> N, and 62 kg ha <sup>-1</sup> P [50]
Growth regulating	Treatment of milk thistle with plant-growth regulators in combination with soil or foliar mineral fertilizers increased the total amount of silymarin by increasing seed yield per hectare [43]
Weed control	Pendimethalin and metribuzin herbicides are safe for weed control, both alone and in combination [33]
Rotation	The milk thistle is recommended for incorporation into arable crop rotation as forecrop of maize [50]
	The rotation could be planned preferring spring-summer crop. During these seasons, milk thistle is smaller and therefore less competitive [36]
	Yields of milk thistle grown in an monoculture were about 40% lower than the
	yields obtained in crop rotation that the yield decrease was mainly due to root damage caused by larvae insect ( <i>Cleonus</i> piger) [52]
Harvesting	Harvest with the use of combined harvester, when 40-50% of inflorescences have pappus [51]
	In July – August after blooming [43]

the Silma cultivar was bred and registered in Poland [37]. Ram et al. (2005) analyzed the genetic variability of 15 accessions of milk thistle based on morphological characteristics silymarin content and and reported considerable variation silymarin in concentration [43]. Hetz (1995) reported that silybum is predominantly a self-pollinator, since the cross-pollination rate on average is only about 2% [25].

In vitro cultures have been used as an alternative source of biologically active compounds [60, 61]. Studies on culture of milk thistle in vitro began in the late 1970s [7]. Tissue culture protocol have been established for milk thistle from cotyledons and sterile plantlets to induce callus on Murashige and Skoog medium (MS) supplemented with different hormones. Optimum results obtained with the use of naphthalenee acetic acid and



kinetin [62, 63]. Liu and Cai referred to the isolation and culture of protoplasts [64]. Cotyledon cultures have employed to produce organogenesis and somatic embryogenesis in milk thistle [65]. Production of flavonolignans from transformed (hairy) and untransformed root cultures of Silybum marianum also was reported [66]. Elicitors have been widely employed to increase the formation of secondary metabolites in plant cell cultures and this strategy has also been effective in stimulating the production of silymarin in cell cultures derived from milk thistle [67]. Production of silymarin and the effect of the elicitor, methyl jasmonate (MeJA), were studied in cell cultures of Silybum marianum [68].

Treatment of S. marianum suspensions with a crude extract of yeast elicitor (YE) improved the production of silymarin and caused the release of silymarin into the culture medium to a level about threefold higher than that of the extract and control. The yeast methyl iasmonate (iasmonic acid derivatives), strongly promoted the accumulation silymarin [67, 68]. Hansaloo et al. (2006), reported the effects of jasmonic acid on silymarin production in milk thistle cultures [69]. Yeast extracts stimulated flavonolignan production in hairy root cultures two fold higher than the control cultures. Moreover, it was reported that the yeast extract treatment induced the activity of lipoxygenase for production of the jasmonate. It was concluded that jasmonate signaling is an integral part of the yeast extract signal transduction for the production of flavonolignans [70]. Besides improving production, elicitation allows the study of signal transduction pathways which regulate the expression of biosynthetic genes

involved in plant secondary metabolism [7, 70]. Rahimi et al. (2011) studied the effect of different concentrations of L-phenylalanine (0, 1, 10 and 100 µM) as the precursor, on thephenylalanine ammonia-lyase activity, naringenin content, root biomass and silymarin production in Silybum marianum hairy roots. All concentrations of added phenylalanine stimulated phenylalanine ammonia-lyase activity [71]. In a study the effect of abiotic elicitors, two newly synthesized substituted amides of pyrazine-2-carboxylic acids on the flavonolignan accumulation in callus and suspension culture of Silybum marianum (L.) Gaertn. was investigated. The compounds markedly influenced the production flavonolignans in an in vitro culture [72].

# Pharmacological applications

The pharmacological active ingredients present in milk thistle fruits collectively are silymarin. Silymarin known contains flavonolignan isomers: structural silybin (silibinin) (50 - 60%), isosilibinin (5%), silicristin (20%), silidianin (10%) and other components such as taxifolin (5%) (Figure 2) [60]. Other constituents include apigenin, silvbonol, dehydrosilybin, deoxysilycistin, deoxysilydianin, silandrin. silvbinome. silyhermin, and neosilyhermin [2, 3, 73]; It has been reported that milk thistle fruits have a relatively high amount of oil (20-31%) [74, 75, 76] that contains fatty acids such as linoleic acid, oleic acid, linolenic acid, palmitic acid, stearic acid. Silybin is the major biologically active constituent of milk thistle and responsible for its pharmacological activity. It has been used for centuries to selftreat liver disorders [3, 74].




Figure 2- Chemical compositions of milk thistle [7]

Milk thistle is used for multiple medicinal purposes, due to its various physiological characteristics. Research has confirmed that silymarin extracted from milk thistle fruits can protect healthy liver cells from deterioration, helping cleanse and detoxification, as well as contributing to regeneration of damaged cells [3, 77].

Various components of milk thistle (silymarin, silybin, etc.) have multiple mechanisms of action that may be hepatoprotective, including anti-inflammatory

activity, antioxidant activity, toxin blockade, enhanced protein synthesis and anti-fibrotic activity [77, 78]. As mentioned, the main action of the active principles of *S. marianum* is hepatoprotective. A summary of the effective pharmacological applications milk thistle included hepatoprotective action, alcoholic liver diseases, viral hepatitis, liver cirrhosis [79, 80], toxic and drug diseases of the liver, mushroom poisoning, diabetes patients with chronic liver disease and hypocholesterolemic action [3, 79].



#### **Antioxidant properties**

The antioxidant properties of milk thistle were evaluated by studing the ability to react with relevant biological ROS or oxidants such as superoxide anion radical  $(O_2)$ , hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), hydroxyl radical (OH) and hypochlorous acid (HOCL). Kiruthiga et al [81] have shown that administration of silymarin increases the activities of antioxidant enzymes like superoxide dismutase (SOD), glutathione catalase, peroxidase glutathione reductase (GR) and glutathione-stransferase (GST) together with a decrease in the levels of malondialdehyde (MDA), a marker for lipid peroxidation, in erythrocytes exposed to  $H_2O_2$  [82].

#### **Anti-inflammatory Activity**

Anti-inflammatory effects of silymarin are related to inhibition of the transcription factor nuclear factor-κB (NF-κB), which regulates and coordinates the expression of various genes involved in inflammation, cell survival, differentiation and growth [82].

#### **Mushroom Poisoning**

The most remarkable use of silymarin is in the treatment of Amanita mushroom poisoning. Amanita mushrooms possess two extremely powerful hepatotoxins, amanitin and phalloidin. Severe liver damage (and death) was avoided if silymarin was administered within 24 hours [83].

#### Alcoholic liver diseases

Research on *In Vitro* and *In Vivo* animal models suggested that silymarin has ability to protect liver cells from toxins. In alcoholic liver diseases (ALD), silymarin was found to exert hepatoprotective effects by attenuating the tumor necrosis factor (TNF) production

Journal of Medicinal Plants, Volume 12, No. 47, Summer 2013 along with decreasing the serum alanine aminotransferase (ALT) activity, inhibiting lipid peroxidation, and increasing intracellular reduced glutathione content in mouse model of ALD [84]. A double-blind study achieved on patients suffered from chronic alcoholic liver disease. The result demonstrated serum bilirubin. that aminotransferase values and gamma glutamil transferase (GGT) activity were normalized in the silymarin group [85].

#### **Anti-cancer Activity**

Carcinogenesis is a multistep process that is activated by altered expression of transcriptional factors and proteins involved in proliferation. cell cycle regulation, differentiation, apoptosis, angiogenesis, invasion and metastasis. Silymarin and silybin modulate imbalance between cell survival and apoptosis through interference with expressions of cell cycle regulators and proteins involved in apoptosis [82].

Anti-cancer activity of silymarin has been demonstrated in human breast cancer, skin cancer, androgen-dependent and -independent prostate cancer, cervical cancer, colon cancer, ovarian cancer, hepatocellular carcinoma, bladder cancer, and lung cancer cells [82, 86, 87, 88].

#### **Anti-diabetic Activity**

The property of silymarin in reducing fasting glycaemia and insulin level have supported its use as an antihyperglycaemic compound. The potent hypoglycaemic and antihyperglycaemic activities of an aqueous extract of milk thistle have also been demonstrated in experimental animal models of diabetes [89, 90].

#### **Hepatoprotective Activity**

Silymarin has been used for centuries as a hepatoprotectant [91]. This effects have been attributed to direct and/or indirect anti-oxidant capacity of silymarin, such as being scavenger of reactive oxygen species, scavenger of phenylglyoxylic ketyl radicals, chain breaking antioxidant [83].

### **Hypocholesterolaemic Activity**

Survey the influence of silymarin and its polyphenolic fraction on rats fed with a high-cholesterol diet showed that silymarin reduced cholesterol levels in the liver and plasma of rats [92].

The hypocholesterolaemic activity of silymarin on the basis of experimental evidence showing that silybin inhibits HMG-CoA reductase activity in vitro; and silymarin improved the binding of low lipoproteins (LDL) to rat hepatyocytes, decreased the liver cholesterol content in rabbits fed with a high-cholesterol diet, decreased the plasma-cholesterol and LDLcholesterol levels in hyperlipaemic rats. The influence of silymarin and polyphenolic fraction (PF) of silymarin on cholesterol absorption in rats fed on high cholesterol diet (HCD) was studied. Silymarin and PF significantly reduced cholesterol absorption in rats fed on HCD and caused significant decreases in content of cholesterol and

triacylglycerol (TAG) in the liver. These results suggest that the inhibition of cholesterol absorption caused by silymarin and its polyphenolic fraction could be a mechanism contributing to the positive changes in plasma cholesterol lipoprotein profile and in lipid content in liver [93].

# Conclusion

Milk thistle is one of the most important medicinal plants grown in the world. Silymarin is the pharmacological active principle of the fruit of this plant. The seeds contain the highest amount of silymarin, but the whole plant is used medicinally. Despite the world demand of silymarin, there is a lack of research efforts on the domestication and improvement of this plant especially in Iran. A few varieties of milk thistle have been developed. Our knowledge of agronomic practices and genotype-environment interactions which could promote high quality and quantity of milk thistle production under Iranian growth conditions is very limited. It is essential to define to what extent agrotechnical practices can affect the quality and quantity of silymarin.

# Acknowledgment

This work was supported by the Cultivation and Development Department of Medicinal Plants Research Center, Institute of Medicinal Plants, ACECR in Karaj-Iran.

#### References

- **1.** Leung AY and Foster S. Encyclopedia of Common Natural Ingredients Used in Food, Drugs & Cosmetic. John Wiley & Sons. 1996, pp: 366 8.
- **2.** Libster M. Delmar's integrative herb guide for nurses. Thamson Learning. 2002, pp. 669 77.



- **3.** Abenavoli L, Spagnuolo R, Luppino I and Luzza F. Recent Progress in Medicinal Plants. Spllc Press. 2010, pp: 387 409.
- **4.** Tumova L, Rimakova J, Tuma J and Dusek J. *Silybum marianum* in vitro flavonolignan production. *Plant Soil Environ*. 2006; 52 (10): 454 –8.
- **5.** Murphy JM, Caban M and Kemper KJ. Longwood Herbal Task Force: http://www.mcp.edu/herbal/default.htm.
- **6.** Cwalina-Ambroziak B, Wierzbowska J, Damszel M and eresa Bowszys T. The effect of mineral fertilization on achenes yield and fungal communities isolated from the stems of milk thistle *Silybum marianum* (L.) Gaertner. *Acta Sci. Pol., Hortorum Cultus* 2012; 11 (4): 157 68.
- 7. Das SK, Mukherjee S and Vasudevan DM. Medicinal properties of milk thistle with special reference to silymarin an overview. *Natural Product Radiance* 2008; 7 (2): 182 92.
- **8.** Corchete P. Bioactive Molecules and Medicinal Plants. Springer-Verlag Berlin Heidelberg. Chapter 6, 2008, pp. 123 42.
- **9.** *Susan* G. Wynn, DVM. *Silybum marianum*: Milk Thistle. wendyblount.com/etvma2/milk-thistle.pdf.
- **10.** Wu JW, LinL C and Tsai TH. Drug–drug interactions of silymarin on the perspective of pharmacokinetics. *Journal of Ethnopharmacol.* 2009; 121: 185 93.
- 11. Wallace S, Vaughn K, Stewart BW, Viswanathan T, Clausen E, Nagarajan S and Carrier DJ. Milk Thistle Extracts Inhibit the Oxidation of Low-Density Lipoprotein (LDL) and Subsequent Scavenger Receptor-

- Dependent Monocyte Adhesion. *J. Agric. Food Chem.* 2008; 56: 3966 72.
- **12.** El-Mallah MH, El-Shami SM and Hassanein MM. Detailed studies on some lipids of *Silybum marianum* (L.) seed oil. *Grasasy Aceites*. 2003; 54 (4): 397 402.
- **13.** Mayer KE, Myers RP and Lee SS. Silymarin treatment of viral hepatitis: a systematic review. *Journal of Viral Hepatitis* 2005; 12: 559 67.
- **14.** Zheng X, Xin W, Yubin L, Shic J, Jun XuecS and Liua Ch. Application of response surface methodology to optimize microwave-assisted extraction of silymarin from milk thistle seeds. *Separation and Purification Technol.* 2009; 70: 34 40.
- **15.** Lee DYW and Liu Y. Molecular Structure and Stereochemistry of Silybin A, Silybin B, Isosilybin A, and Isosilybin B, Isolated from *Silybum marianum* (Milk Thistle). *J. Nat. Prod.* 2003; 66: 1171 4.
- **16.** Lee J I, Hsub BH, Wua D and Barrett JS. Separation and haracterization of silybin, isosilybin, silydianin and silychristin in milk thistle extract by liquid chromatography–electrospray tandem mass spectrometry. *J. Chromatography* 2006; 1116: 57 –68.
- **17.** Brindaa BJ, Zhua H and Markowitza JS. A sensitive LC–MS/MS assay for the simultaneous analysis of the major active components of silymarin in human plasma. *J. Chromatography* 2012; 902: 1 9.
- **18.** Abourashed Ehab A. Julie R. Mikell b, Ikhlas A. Khan. Bioconversion of silybin to phase I and II microbial metabolites with retained antioxidant activity. *Bioorganic & Medicinal Chem.* 2012; 20: 2784 8.

- **19.** Omidbaigi R. Approaches to production and processing of medicinal plants. Tarrahan Nashr. Iran. Vol. 2. 1997, p: 319.
- **20.** Mozaffarian V. A dictionary of Iranian Plant Names (3*ed.*). Farhang Moaser Press. Iran. 2003, pp: 59 60.
- **21.** Ghahreman A. Flora of IRAN. Research Institute of Forests Rangelands, Iran, 1999, 3, pp: 587.
- **22.** Hammouda SI, Ismail SI, Abdel-Azim NS and Shams KA. *Silybum marianum* (L.) Gaertn. A Guide to Medicinal Plants in North Africa http://www.uicnmed.org/nabp/database/HTM/PDF/p61.pdf.
- 23. Dini M. Investigation of Various Common Names of Plants Used in Traditional Medicine, Research Institute of Forests Rangelands, Iran, 2006, pp. 44.
- **24.** Anonomys. WHO monographs on selected medicinal plants. World Health organization. 2007, Vol. 2.
- **25.** Hetz E, Liersch R and Schieder O. Genetic investigations on *Silybum marianum* and *S. eburneum* with respect to leaf colour, outcrossing ratio, and flavonolignan composition. *Planta Med.* 1995; 61: 54 7.
- **26.** Vaknin Y, Hadas R, Schafferman D, Murkhovsky D and Bashan N. The potential of milk thistle (*Silybum marianum* L.), an Israeli native, as a source of edible sprouts rich in antioxidants. *International Journal of Food Sciences and Nutrition* 2008; 59 (4): 339 46.
- **27.** British Pharmacopoeia. Herbal Drugs and Herbal Drug Preparations Milk-thistle Fruit. 2009, Volume III. pp: 7173.
- **28.** Gresta F, Avola G and Guarnaccia P. Agronomic characterization of some


- spontaneous genotypes of milk thistle (*Silybum marianum* L. Gaertn.) in Mediterranean environment. *Journal of Herbs, Spices & Medicinal Plants* 2006; 12 (4): 51 60.
- **29.** Najda A and Dyduch J. The evaluation of the quality of the milk thistle (*Silybum marianum* L. Gaertn.) fruits depending on their shapeliness. *Herba Polonica*. 2007; 53 (3): 337 42.
- 30. Flora von Deutschland Österreich und der Schweiz (1885) Otto Wilhelm Thomé from site. Index: Lateinische botanische Namen. 2013. http://caliban.mpiz-koeln.mpg.de/thome/index.html.
- **31.** Shokrpour M, Torabi GiglooM, li Asghari A and Bahrampour Sh. Study of some agronomic attributes in milk thistle (*Silybum marianum* Gaertn.) ecotypes from Iran. *Journal of Medicinal Plants Res.* 2011; 5 (11): 2169 74.
- **32.** Montemurro P, Fracchiolla M and Lonigro A. Effects of Some Environmental Factors on Seed Germination and Spreading Potentials of *Silybum marianum* Gaertn. *Ital. J. Agron. / Riv. Agron.* 2007; 3: 315 20.
- **33.** Karkanis A, Bilalis D and Efthimiadou A. Cultivation of milk thistle (*Silybum marianum* L. Gaertn.), a medicinal weed. *Industrial Crops and Products* 2011; 34 (1): 825 30.
- **34.** Belitz AR and Sams CE. The effect of population density on growth, yield and flavonolignan content in milk thistle (*Silybum marianum*). *Acta Horticulture* 2007; 756: 251 8.
- **35.** Omidbaigi R and Nobakht A. Nitrogen fertilizer affecting growth, seed yield and



active substances of milk thistle (*Silybum marianum*). *Pak. J. Biol. Sci.* 2001; 4: 1345 – 9.

- **36.** Kozera W, Nowak K and Techniczno-Rolnicza A. The effect of fertilization on milk thistle (*Silybum marianum*) yield and its chosen features. *Annales Universitatis Mariae Curie-Sklodowska*. 2004; 59 (1): 369 74.
- **37.** Andrzejewska J, Sadowska K and Mielcarek S. Effect of sowing date and rate on the yield and flavonolignan content of the fruits of milk thistle (*Silybum marianum* L. Gaertn.) grown on light soil in a moderate climate. *Industrial Crops and Products* 2011; 33 (2): 462 8.
- **38.** Morazzoni P and Bombardelli E. Silybum marianum (*Carduus marianus*). *Fitoterapia* 1995; 66: 3 42.
- **39.** Carrier DJ, Crowe T, Sokhansanj S, Wahab J and Barl B. Milk thistle, *Silybum marianum* L. Gaertn., flower head development and associated marker compound profile. *J. Herbs Spices Med. Plants* 2002; 10: 65 74.
- **40.** Azim Khan M, Blackshaw R E and Marwat K B. Biology of milk thistle (*Silybum marianum*) and the management options for growers in North-Western Pakistan. *Weed Biology and Management* 2009; 9: 99 105.
- **41.** Carrubba A and la Torre R. Cultivation trials of milk thistle *Silybum marianum* Gaertn. into the semiarid Mediterranean environment. *Agric. Med.* 2003; 133: 14 9.
- **42.** Haban M, Otepka P, Kobida L and Habanova M. Production and quality of milk thistle (*Silybum marianum* (L.) Gaertn.) cultivated in cultural conditions of warm agri-

- climatic macroregion. *Horti. Sci.* 2009; 36 (2): 25 30.
- **43.** Ram G, Bhan MK, Gupta KK, Thaker B, Jamwal U and Pal S. Variability pattern and correlation studies in *Silybum marianum* Gaertn. *Fitoterapia* 2005; 76: 143 7.
- **44.** Ryant P. Nutrition and fertilization and of alternative oil plants for non-food purpose. 2003, pp. 1 15.
- **45.** Mel'nktov T. M. Morphological-biological characteristics of *Silybum marianum* seeds as sowing material. *Khimiko Farmatsevticheskii Zhurnal*. 1983; 17 (8): 958 63.
- **46.** Ghavami N and Ramin A.A. Salinity and temperature effects on seed germination of milk thistle. *Commun. Soil. Sci. Plant Anal.* 2007; 38: 2681 91.
- **47.** Dodd J. Phenology and seed production of variegated thistle, *Silybum marianum* (L.) Gaertn., in Australia in relation to mechanical and biological control. *Weed Res.* 1989; 29 (4): 255 63.
- **48.** Omer EA, Ibrahim ME, Razin AM and Ahmed SS. Effect of spacing, nitrogen and potassium fertilization of *Silybum marianum* L. cultivated in newly reclaimed lands. *Egyptian Journal of Horticulture* 1996; 22 (1): 97 101.
- **49.** Andrzejewska J and Sadowska K. Effect of the sowing date on the content and composition of flavonolignans and nutrients in milk thistle (*Silybum marianum* (L.) Gaertn.) fruit. *Herba Pol.* 2007; 3: 273 8.
- **50.** Purohit SM, Iqbal M and Srivastava PS. Seed germination studies on an important medicinal plant, *Silybum marianum* (Linn.) *Hamdard Medicus* 1997; 40 (2): 31 3.




- **51.** Andrzejewska J and Sadowska K. Effect of cultivation conditions on the variability and interrelation of yield and raw material quality in milk thistle (*Silybum marianum* (L.) Gaertn.). *Acta Sci. Pol.*, *Agricultura*. 2008; 7 (3): 3 11.
- **2.** Andrzejewska J and Skinder Z. Yield and quality of raw material of milk thistle [*Silybum marianum* (L.) Gaertn.] grown in monoculture and in crop rotation. Part I. Reaction of milk thistle to the sowing date. *Herba Pol.* 2006; 4: 11 7.
- **33.** Rahimi A and Kamali M. Different planting date and fertilizing system effects on the seed yield, essential oil and nutrition uptake of milk thistle (*Silybum marianum* (L.) Gaertn.). *Advances in Environmental Biol*. 2012; 6 (5): 1789 96.
- **54.** Hammouda FM, Ismail SI, Hassan NM and Zaki AK. Comparative studies of the oil from *Silybum marianum* cultivated in Egypt using GLC. *Qatar University Sci. J.* 1994; 14: 154 7.
- **55.** Andrzejewska J and Skinder Z. Yield and quality of milk thistle (*Silybum marianum* (L). Gaertn.) raw material grown in monoculture and in crop rotation. Part 2. Milk thistle reaction to potassium fertilization. *Herba Polonica*. 2007; 53 (1): 5 10.
- **56.** Omer E A. Effect of different nitrogen sources on Romanian *Silybum marianum* cultivated in sandy and clay soils. *Egyptian Journal of Horticulture* 1996; 23 (1): 63 76.
- **57.** Kazmierczak K, Seidler-Łoz K. Silma the Polish variety of milk thistle (*Silybum marianum* L. Gaertn.). *Herba Polonica*. 1997; 3: 195 7.

- **58.** Asghari-Zakaria R, Panahi AR and Sadeghizadeh M. Comparative study of chromosome morphology in *Silybum marianum* L. *Cytologia*. 2008; 73: 327 32.
- **59.** Sarrami M, Zeinali H and Bakhshi Khaniki GH. Population diversity and analysis of cytogenetic affinities between different *Silybum marianum* L. populations from Iran. *Nucleus* 2012; 1 6.
- **60.** Radjabian T, Rezazadeh SH and Fallah-Huseini H. Analysis of silymarin components in the seed extracts of some milk thistle ecotypes from Iran by HPLC. *Iranian Journal of Science & Technol.* 2008; 32 (A2): 141 6.
- **61.** Ghavami N and Ramin A.A. Grain yield and active substances of milk thistle as affected by soil salinity. *Commun. Soil Sci. Plant Anal.* 2008; 39: 2608 18.
- **2.** Gabay R, Plitmann U and Danin A. Factors affecting the dominance of *Silybum marianum* L. (Asteraceae) in its specific habitats. *Flora* 1994; 189: 201 6.
- **63.** Verpoorte R, Heijden VR, Hoopen H and Memelink G.J. Metabolic engineering of plant secondary metabolite pathways for the production of fine chemicals. *Biotechnol. Lett.* 1999; 21: 467 79.
- **64.** Liu S and Cai QG. Callus Formation from Protoplasts and Plant Regeneration from Tissue Culture of *Silybum marianum* Gaertn. *J. Integrative Plant Biol.* 1990; 32 (1): 19 25.
- **65.** Radice S, Caso OH. Somatic embryogenesis and organogenesis in cultured cotyledons of *Silybum marianum* (L.) Gaertn., *Biocell*. 1997; 21 (3): 59 64.
- 66. Alikaridis F, Papadakis D, Pantelia K and



- Kephalas T. Flavonolignan production from *Silybum marianum* transformed and untransformed root cultures. *Fitoterapia* 2000; 71 (4): 379 84.
- **67.** Sanchez-Sampedro MA, Fernandez-Tarrago J and Corchete P. Yeast extract and methyl jasmonate induced silymarin production in cell cultures of *Silybum marianum* (L.) Gaertn. *J. Biotechnol*. 2005; 119: 60 9.
- **68.** Sanchez-Sampedro MA, Fernandez-Tarrago J, Corchete P. Elicitation of silymarin in cell cultures of *Silybum marianum*: effect of subculture and repeated addition of methyl jasmonate. *J. Biotechnol.* 2009; 31: 1633 7.
- **69.** Hasanloo T, Khavari-Nejad RA, Majidi E, Shams Ardakani MR. *Iran J. Pharm. Sci.* 2006; 2: 206.
- **70.** Hasanloo T, Khavari-Nejad RA, Majidi E and Shams Ardakani MR. Flavonolignan Production in Cell Suspension Culture of *Silybum marianum*. *Pharmaceutical Biol*. 2008; 46 (12): 876 82.
- 71. Rahimi Sh, Hasanloo T, Najafi F, Khavari-Nejad R. Enhancement of silymarin accumulation using precursor feeding in *Silybum marianum* hairy root cultures. *POJ*. 2011; 4 (1): 34 9.
- **72.** Tumova L, Gallova K, Rimakova J. *Silybum marianum*. In vitro. *Ceska Slov Farmacy* 2004; 53: 135 40.
- **73.** Kvasnicka F, Bida B, Sevcik R, Voldrich M, Kratka, JM. Analysis of the active components of silymarin. *J. Chromatogr*. 1990; 1: 239 –245.
- **74.** Hadolin M, Skerget M, Knez Z, Bauman D. High pressure extraction of vitamin E-rich

- oil from *Silybum marianum*. *Food Chem*. 2001; 74: 355 64.
- **75.** Fathi-Achachlouei B and Azadmard-Damirchi S. Milk Thistle Seed Oil Constituents from Different Varieties Grown in Iran. *J. Am. Oil. Chem. Soc.* 2009; 86: 643 9.
- **76.** Qavami N, Labbafi MR, Dehghani-Meshkani MR and Mehrafarin A. Determination of Seed and Oil Yield and Yield Components in Two Variety of Milk Thistle (*Silybum marianum* Gaertn.) Based on Path Analysis and Regression. Medicinal plants. *Medicinal Plants* 2012; 11 (44): 78 84.
- 77. Davis-Searles PR, NakanishiY, Kim N, Graf TN, Oberlies NH, Wani MC, Wall ME, Agarwal R and Kroll DJ. Milk Thistle and Prostate Cancer: Differential Effects of Pure Flavonolignans from *Silybum marianum* on Antiproliferative end Points in Human Prostate Carcinoma Cells. *Cancer Res.* 2005; 65 (10): 4448 57.
- **78.** Johnn S, Scaler F, Sonnenbichler I and Weyhenmeyerr. Stimulatory Effects of Silibinin and Silicristin from the Milk Thistle *Silybum marianum* on Kidney Cells. *JPET*. 1999; 290 (3): 1375 83.
- **79.** Ball KR and Kowdley KV. A review of *Silybum marianum* (Milk Thistle) as a Treatment for alcoholic liver disease. *Journal of Clinical Gastroenterol*. 2005; 39 (6): 520 8.
- **80.** Fallah Huseini H, Alavian SM, Toliat T, Jamshidi AH, Heshmat R, Naghdi Badi H and Khani M. The efficacy of herbal medicine khar maryam (*Silybum marianum* (L.) Gaertn.) on liver cirrhosis in chronic hepatitis B patients. *J. Medicinal Plants* 1994; 8: 1 6.



- 81. Kiruthiga PV, Shafreen RB, Pandian SK, Devi KP. Silymarin protection against major reactive species released oxygen by environmental toxins:  $H_2O_2$ exogenous erythrocytes. Clin. exposure in **Basic** *Pharmacol. Toxicol.* 2007; 100: 414 – 9.
- **82.** Ramasamy K and Agarwal R. Multitargeted therapy of cancer by silymarin. *Cancer Lett.* 2008; 269 (2): 352 62.
- **83.** Luper S. A Review of Plants Used in the Treatment of Liver Disease: part 1. *Alternative Medicine Review 3*. 1998; 410 21.
- 84. Ladas EJ, Kroll DJ, Oberlies NH, Cheng B and Ndao DH. A Randomized, Controlled, Double-Blind, Pilot Study of Milk Thistle for the Treatment of Hepatotoxicity in Childhood Acute Lymphoblastic Leukemia (ALL). *Cancer* 2010; 116: 506 13.
- **85.** Feher J, Deak G, Muzes G, Lang I, Niederland V, Nekam K and Karteszi M. Liver-protective action of silymarin therapy in chronic alcoholic liver diseases. *Orvosi Hetilap.* 1989; 130: 2723 7.
- **86.** Singh RP and Agarwal R. Mechanisms and preclinical efficacy of silibinin in preventing skin cancer. *Eur. J. Cancer.* 2005; 41: 1969 79.
- **87.** Singh RP and Agarwal R. Prostate cancer chemoprevention by silibinin: bench to bedside. *Mol. Carcinog.* 2006; 45 (6): 436 42.
- **88.** Tyagi A, Singh RP, Ramasamy K, Raina K, Redente EF, Dwyer- Nield LD, Radcliffe RA, Malkinson AM and Agarwal R. Growth

- inhibition and regression of lung tumors by silibinin: modulation of angiogenesis by macrophage-associated cytokines and nuclear factor-kappaB and signal transducers and activators of transcription 3. *Cancer Prev. Res.* 2009; 2: 74 83.
- **89.** Soto CP, Perez BL, Favari LP and Reyes JL. Prevention of alloxan-induced diabetes mellitus in the rat by silymarin, *Comp. Biochem. Physiol. C. Pharmacol. Toxicol. Endocrinol.* 1998; 119: 125 9.
- **90.** Maghrani M, Zeggwagh NA, Lemhadri A, El Amraoui M, Michel JB and Eddouks M. Study of the hypoglycaemic activity of Fraxinus excelsior and *Silybum marianum* in an animal model of type 1 diabetes mellitus. *J. Ethnopharmacol.* 2004; 91: 309 16.
- **91.** Polyak SJ, Morishima C, Lohmann V, Pala S, Lee D YW, Liu Y, Graf T N and Oberlies N H. Identification of hepatoprotective favonolignans from silymarin. *PNAS*. 2010; 107 (13): 5995–5999.
- **92.** Skottova N, Vecera R, Urbanek K, Vana P, Walterova D and Cvak L. Effects of polyphenolic fraction of silymarin on lipoprotein profile in rats fed cholesterol-rich diets. *Pharmacol. Res.* 2003; 47 (1): 17 26.
- **93.** Sobolova L, Skottova N, Vecera R and Urbanek K. Effect of silymarin and its polyphenolic fraction on cholesterol absorption in rats. *Pharmacol Res.* 2006; 53 (2): 104 12.

